

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
6 December 2001 (06.12.2001)

PCT

(10) International Publication Number
WO 01/92525 A2

- (51) International Patent Classification: **C12N 15/12**, 15/62, C07K 14/47, 16/30, C12Q 1/68, G01N 33/53, 33/574, A61K 39/00, 39/395, A61P 35/00
- (74) Agents: **VERNA, James, V.**; Seed Intellectual Property Law Group PLLC, Suite 6300, 701 Fifth Avenue, Seattle, WA 98104-7092 et al. (US).
- (21) International Application Number: PCT/US01/17066
- (22) International Filing Date: 25 May 2001 (25.05.2001)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/207,485 26 May 2000 (26.05.2000) US
60/230,475 6 September 2000 (06.09.2000) US
- (71) Applicant (for all designated States except US): **CORIXA CORPORATION** [US/US]; 1124 Columbia Street, Suite 200, Seattle, WA 98104 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **HARLOCKER, Susan, L.** [US/US]; 7522 13th Avenue W., Seattle, WA 98117 (US). **WANG, Tongtong** [US/US]; 8049 N.E. 28th Street, Medina, WA 98039 (US). **BANGUR, Chaitanya, S.** [IN/US]; 2102 N. 105th Street, Apt. 1101, Seattle, WA 98133 (US). **KLEE, Jennifer, Irmela** [US/US]; 5255 19th Avenue N.E., Seattle, WA 98105 (US). **SWITZER, Ann** [US/US]; 3838 N.E. 85th Street, Seattle, WA 98115 (US).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: COMPOSITIONS AND METHODS FOR THE THERAPY AND DIAGNOSIS OF LUNG CANCER

(57) Abstract: Compositions and methods for the therapy and diagnosis of cancer, particularly lung cancer, are disclosed. Illustrative compositions comprise one or more lung tumor polypeptides, immunogenic portions thereof, polynucleotides that encode such polypeptides, antigen presenting cell that expresses such polypeptides, and T cells that are specific for cells expressing such polypeptides. The disclosed compositions are useful, for example, in the diagnosis, prevention and/or treatment of diseases, particularly lung cancer.



WO 01/92525 A2

COMPOSITIONS AND METHODS FOR THE THERAPY AND DIAGNOSIS OF LUNG CANCER

TECHNICAL FIELD OF THE INVENTION

The present invention relates generally to therapy and diagnosis of cancer, such as lung cancer. The invention is more specifically related to polypeptides, comprising at least a portion of a lung tumor protein, and to polynucleotides encoding such polypeptides. Such polypeptides and polynucleotides are useful in pharmaceutical compositions, *e.g.*, vaccines, and other compositions for the diagnosis and treatment of lung cancer.

10 BACKGROUND OF THE INVENTION

Cancer is a significant health problem throughout the world. Although advances have been made in detection and therapy of cancer, no vaccine or other universally successful method for prevention and/or treatment is currently available. Current therapies, which are generally based on a combination of chemotherapy or surgery and radiation, continue to prove inadequate in many patients.

Lung cancer is a significant health problem throughout the world. In the U.S., lung cancer is the primary cause of cancer death among both men and women, with an estimated 172,000 new cases being reported in 1994. The five-year survival rate among all lung cancer patients, regardless of the stage of disease at diagnosis, is only 13%. This contrasts with a five-year survival rate of 46% among cases detected while the disease is still localized. However, early detection of lung cancer is difficult since clinical symptoms are often not seen until the disease has reached an advanced stage, and only 16% of lung cancers are discovered before the disease has spread.

In spite of considerable research into therapies for these and other cancers, lung cancer remains difficult to diagnose and treat effectively. Accordingly, there is a need in the art for improved methods for detecting and treating such cancers. The present invention fulfills these needs and further provides other related advantages.

SUMMARY OF THE INVENTION

In one aspect, the present invention provides polynucleotide compositions comprising a sequence selected from the group consisting of:

- (a) sequences provided in SEQ ID NO:1-35, 42-55, 58-60, 63-91 and 93-95;
- (b) complements of the sequences provided in SEQ ID NO:1-35, 42-55, 58-60, 63-91 and 93-95;
- (c) sequences consisting of at least 20, 25, 30, 35, 40, 45, 50, 75 and 100 contiguous residues of a sequence provided in SEQ ID NO: 1-35, 42-55, 58-60, 63-91 and 93-95;
- (d) sequences that hybridize to a sequence provided in SEQ ID NO:1-35, 42-55, 58-60, 63-91 and 93-95, under moderate or highly stringent conditions;
- (e) sequences having at least 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98% or 99% identity to a sequence of SEQ ID NO:1-35, 42-55, 58-60, 63-91 and 93-95; and
- (f) degenerate variants of a sequence provided in SEQ ID NO:1-35, 42-55, 58-60, 63-91 and 93-95.

In one preferred embodiment, the polynucleotide compositions of the invention are expressed in at least about 20%, more preferably in at least about 30%, and most preferably in at least about 50% of lung tumors samples tested, at a level that is at least about 2-fold, preferably at least about 5-fold, and most preferably at least about 10-fold higher than that for normal tissues.

The present invention, in another aspect, provides polypeptide compositions comprising an amino acid sequence that is encoded by a polynucleotide sequence described above.

The present invention further provides polypeptide compositions comprising an amino acid sequence selected from the group consisting of sequences recited in SEQ ID NO:36-41, 56, 57, 61, 62, 92 and 96.

In certain preferred embodiments, the polypeptides and/or polynucleotides of the present invention are immunogenic, *i.e.*, they are capable of

eliciting an immune response, particularly a humoral and/or cellular immune response, as further described herein.

The present invention further provides fragments, variants and/or derivatives of the disclosed polypeptide and/or polynucleotide sequences, wherein the fragments, variants and/or derivatives preferably have a level of immunogenic activity of at least about 50%, preferably at least about 70% and more preferably at least about 90% of the level of immunogenic activity of a polypeptide sequence set forth in SEQ ID NO:36-41, 56, 57, 61, 62, 92 and 96 or a polypeptide sequence encoded by a polynucleotide sequence set forth in SEQ ID NO:1-35, 42-55, 58-60, 63-91 and 93-95.

10 The present invention further provides polynucleotides that encode a polypeptide described above, expression vectors comprising such polynucleotides and host cells transformed or transfected with such expression vectors.

Within other aspects, the present invention provides pharmaceutical compositions comprising a polypeptide or polynucleotide as described above and a physiologically acceptable carrier.

Within a related aspect of the present invention, the pharmaceutical compositions, *e.g.*, vaccine compositions, are provided for prophylactic or therapeutic applications. Such compositions generally comprise an immunogenic polypeptide or polynucleotide of the invention and an immunostimulant, such as an adjuvant.

20 The present invention further provides pharmaceutical compositions that comprise: (a) an antibody or antigen-binding fragment thereof that specifically binds to a polypeptide of the present invention, or a fragment thereof; and (b) a physiologically acceptable carrier.

Within further aspects, the present invention provides pharmaceutical compositions comprising: (a) an antigen presenting cell that expresses a polypeptide as described above and (b) a pharmaceutically acceptable carrier or excipient. Illustrative antigen presenting cells include dendritic cells, macrophages, monocytes, fibroblasts and B cells.

Within related aspects, pharmaceutical compositions are provided that comprise: (a) an antigen presenting cell that expresses a polypeptide as described above and (b) an immunostimulant.

The present invention further provides, in other aspects, fusion proteins that comprise at least one polypeptide as described above, as well as polynucleotides encoding such fusion proteins, typically in the form of pharmaceutical compositions, *e.g.*, vaccine compositions, comprising a physiologically acceptable carrier and/or an immunostimulant. The fusions proteins may comprise multiple immunogenic polypeptides or portions/variants thereof, as described herein, and may further comprise one or more polypeptide segments for facilitating the expression, purification and/or immunogenicity of the polypeptide(s).

Within further aspects, the present invention provides methods for stimulating an immune response in a patient, preferably a T cell response in a human patient, comprising administering a pharmaceutical composition described herein. The patient may be afflicted with lung cancer, in which case the methods provide treatment for the disease, or patient considered at risk for such a disease may be treated prophylactically.

Within further aspects, the present invention provides methods for inhibiting the development of a cancer in a patient, comprising administering to a patient a pharmaceutical composition as recited above. The patient may be afflicted with lung cancer, in which case the methods provide treatment for the disease, or patient considered at risk for such a disease may be treated prophylactically.

The present invention further provides, within other aspects, methods for removing tumor cells from a biological sample, comprising contacting a biological sample with T cells that specifically react with a polypeptide of the present invention, wherein the step of contacting is performed under conditions and for a time sufficient to permit the removal of cells expressing the protein from the sample.

Within related aspects, methods are provided for inhibiting the development of a cancer in a patient, comprising administering to a patient a biological sample treated as described above.

Methods are further provided, within other aspects, for stimulating and/or expanding T cells specific for a polypeptide of the present invention, comprising contacting T cells with one or more of: (i) a polypeptide as described above; (ii) a polynucleotide encoding such a polypeptide; and/or (iii) an antigen presenting cell that

expresses such a polypeptide; under conditions and for a time sufficient to permit the stimulation and/or expansion of T cells. Isolated T cell populations comprising T cells prepared as described above are also provided.

Within further aspects, the present invention provides methods for
5 inhibiting the development of a cancer in a patient, comprising administering to a patient an effective amount of a T cell population as described above.

The present invention further provides methods for inhibiting the development of a cancer in a patient, comprising the steps of: (a) incubating CD4⁺ and/or CD8⁺ T cells isolated from a patient with one or more of: (i) a polypeptide
10 comprising at least an immunogenic portion of polypeptide disclosed herein; (ii) a polynucleotide encoding such a polypeptide; and (iii) an antigen-presenting cell that expressed such a polypeptide; and (b) administering to the patient an effective amount of the proliferated T cells, and thereby inhibiting the development of a cancer in the patient. Proliferated cells may, but need not, be cloned prior to administration to the
15 patient.

Within further aspects, the present invention provides methods for determining the presence or absence of a cancer, preferably a lung cancer, in a patient comprising: (a) contacting a biological sample obtained from a patient with a binding agent that binds to a polypeptide as recited above; (b) detecting in the sample an amount
20 of polypeptide that binds to the binding agent; and (c) comparing the amount of polypeptide with a predetermined cut-off value, and therefrom determining the presence or absence of a cancer in the patient. Within preferred embodiments, the binding agent is an antibody, more preferably a monoclonal antibody.

The present invention also provides, within other aspects, methods for
25 monitoring the progression of a cancer in a patient. Such methods comprise the steps of: (a) contacting a biological sample obtained from a patient at a first point in time with a binding agent that binds to a polypeptide as recited above; (b) detecting in the sample an amount of polypeptide that binds to the binding agent; (c) repeating steps (a) and (b) using a biological sample obtained from the patient at a subsequent point in
30 time; and (d) comparing the amount of polypeptide detected in step (c) with the amount

detected in step (b) and therefrom monitoring the progression of the cancer in the patient.

The present invention further provides, within other aspects, methods for determining the presence or absence of a cancer in a patient, comprising the steps of: (a) contacting a biological sample, e.g., tumor sample, serum sample, etc., obtained from a patient with an oligonucleotide that hybridizes to a polynucleotide that encodes a polypeptide of the present invention; (b) detecting in the sample a level of a polynucleotide, preferably mRNA, that hybridizes to the oligonucleotide; and (c) comparing the level of polynucleotide that hybridizes to the oligonucleotide with a predetermined cut-off value, and therefrom determining the presence or absence of a cancer in the patient. Within certain embodiments, the amount of mRNA is detected via polymerase chain reaction using, for example, at least one oligonucleotide primer that hybridizes to a polynucleotide encoding a polypeptide as recited above, or a complement of such a polynucleotide. Within other embodiments, the amount of mRNA is detected using a hybridization technique, employing an oligonucleotide probe that hybridizes to a polynucleotide that encodes a polypeptide as recited above, or a complement of such a polynucleotide.

In related aspects, methods are provided for monitoring the progression of a cancer in a patient, comprising the steps of: (a) contacting a biological sample obtained from a patient with an oligonucleotide that hybridizes to a polynucleotide that encodes a polypeptide of the present invention; (b) detecting in the sample an amount of a polynucleotide that hybridizes to the oligonucleotide; (c) repeating steps (a) and (b) using a biological sample obtained from the patient at a subsequent point in time; and (d) comparing the amount of polynucleotide detected in step (c) with the amount detected in step (b) and therefrom monitoring the progression of the cancer in the patient.

Within further aspects, the present invention provides antibodies, such as monoclonal antibodies, that bind to a polypeptide as described above, as well as diagnostic kits comprising such antibodies. Diagnostic kits comprising one or more oligonucleotide probes or primers as described above are also provided.

These and other aspects of the present invention will become apparent upon reference to the following detailed description and attached drawings. All references disclosed herein are hereby incorporated by reference in their entirety as if each was incorporated individually.

5 SEQUENCE IDENTIFIERS

SEQ ID NO:1 is the cDNA sequence for Clone ID # 55964 which is named clone L1040C, and is the same sequence as SEQ ID NO:2337 from U.S. Provisional Application 60/207,485.

10 SEQ ID NO:2 is an extended cDNA sequence for L1040C (Clone ID # 55964).

SEQ ID NO:3 is the cDNA sequence for Clone ID # 58269 which is named clone L1039C, and is the same sequence as SEQ ID NO:7264 from U.S. Provisional Application 60/207,485.

15 SEQ ID NO:4 is an extended cDNA sequence for L1039C (Clone ID # 58269), and which corresponds to the fbx5 F-box gene.

SEQ ID NO:5 is the cDNA sequence for Clone ID # 58267 which is named clone L1037C, and is the same sequence as SEQ ID NO:4978 from U.S. Provisional Application 60/207,485.

20 SEQ ID NO:6 is an extended cDNA sequence for L1037C (Clone # 58267), and which corresponds to the mitotic checkpoint kinase mad3-like gene.

SEQ ID NO:7 is the cDNA sequence for Clone ID # 58245 which is named clone L1038C, and is the same sequence as SEQ ID NO:1796 from U.S. Provisional Application 60/207,485.

25 SEQ ID NO:8 is an extended cDNA sequence for L1038C (Clone ID # 58245), and which corresponds to a neuronal ER localized gene.

SEQ ID NO:9 is the cDNA sequence for Clone ID # 55571 which is named clone L1027C, and is the same sequence as SEQ ID NO:4538 from U.S. Provisional Application 60/207,485.

30 SEQ ID NO:10 is an extended cDNA sequence for L1027C (Clone ID # 55571).

- SEQ ID NO:11 is the cDNA sequence for Clone ID # 55978.
SEQ ID NO:12 is an extended cDNA sequence for Clone ID # 55978.
SEQ ID NO:13 is the cDNA sequence for Clone ID # 55980.
SEQ ID NO:14 is an extended cDNA sequence for Clone ID # 55980.
5 SEQ ID NO:15 is the cDNA sequence for Clone ID # 58346.
SEQ ID NO:16 is an extended cDNA sequence for Clone ID # 58346.
SEQ ID NO:17 is the cDNA sequence for Clone ID # 55561.
SEQ ID NO:18 is an extended cDNA sequence for Clone ID # 55561.
SEQ ID NO:19 is the cDNA sequence for Clone ID # 55984.
10 SEQ ID NO:20 is an extended cDNA sequence for Clone ID # 55984,
and which corresponds to a gt mismatch glycosylase gene.
SEQ ID NO:21 is the cDNA sequence for Clone ID # 58261.
SEQ ID NO:22 is an extended cDNA sequence for Clone ID # 58261,
and which corresponds to a phosphoserine aminotransferase gene.
15 SEQ ID NO:23 is the cDNA sequence for Clone ID # 58348.
SEQ ID NO:24 is an extended cDNA sequence for Clone ID # 58348,
and which corresponds to a hCAP gene.
SEQ ID NO:25 is the cDNA sequence for Clone ID # 56016.
SEQ ID NO:26 is an extended cDNA sequence for Clone ID # 56016.
20 SEQ ID NO:27 is the cDNA sequence for Clone ID # 55987.
SEQ ID NO:28 is an extended cDNA sequence for Clone ID # 55987.
SEQ ID NO:29 is the cDNA sequence for Clone ID # 55956.
SEQ ID NO:30 is an extended cDNA sequence for Clone ID # 55956.
SEQ ID NO:31 is the cDNA sequence for Clone ID # 55952.
25 SEQ ID NO:32 is the cDNA sequence for Clone ID # 55957.
SEQ ID NO:33 is an extended cDNA sequence for Clone ID # 55957.
SEQ ID NO:34 is the cDNA sequence for Clone ID # 55559.
SEQ ID NO:35 is an extended cDNA sequence for Clone ID # 55559.
30 SEQ ID NO:36 is an amino acid sequence of an ORF for L1027C,
encoded by the polynucleotide of SEQ ID NO: 10.

SEQ ID NO:37 is an amino acid sequence of the F-box protein Fbx5 encoded by SEQ ID NO:4.

SEQ ID NO:38 is an amino acid sequence of the mitotic checkpoint kinase MAD3-like protein encoded by SEQ ID NO:6.

- 5 SEQ ID NO:39 is an amino acid sequence of the neuronal olfactomedin-related ER localized protein encoded by SEQ ID NO:8.

SEQ ID NO:40 is an amino acid sequence of the phosphoserine aminotransferase encoded by SEQ ID NO:22.

- 10 SEQ ID NO:41 is an amino acid sequence of the gt mismatch glycosylase encoded by SEQ ID NO:20.

SEQ ID NO:42 is the determined cDNA sequence for Clone ID # 63575 which is named clone L1053C.

SEQ ID NO:43 is the determined cDNA sequence for Clone ID # 63582 which is named clone L1054C.

- 15 SEQ ID NO:44 is the determined cDNA sequence for Clone ID # 63598 which is named clone L1055C.

SEQ ID NO:45 is the determined cDNA sequence for Clone ID # 64963 which is named clone L1056C.

- 20 SEQ ID NO:46 is the determined cDNA sequence for Clone ID # 64988 which is named clone L1058C.

SEQ ID NO:47 is the determined cDNA sequence for Clone ID # 63485.

SEQ ID NO:48 is the determined cDNA sequence for Clone ID # 65010.

- 25 SEQ ID NO:49 is a predicted full-length cDNA sequence for SEQ ID NO:42 which is a full-length sequence from Genbank for an insulinoma-associated 1 mRNA.

SEQ ID NO:50 is a predicted full-length cDNA sequence for SEQ ID NO:43 which is a full-length sequence from Genbank for KIAA0535.

SEQ ID NO:51 is a predicted extended cDNA sequence for SEQ ID NO:44.

- 30 SEQ ID NO:52 is a a predicted full-length cDNA sequence for SEQ ID NO:45 which is a full-length sequence from genbank for a human DAZ mRNA 3'UTR.

- SEQ ID NO:53 is a predicted extended cDNA sequence for SEQ ID NO:46.
- SEQ ID NO:54 is a predicted extended cDNA sequence for SEQ ID NO:47.
- 5 SEQ ID NO:55 is a predicted extended cDNA sequence for SEQ ID NO:48.
- SEQ ID NO:56 is the deduced amino acid sequence encoded by SEQ ID NO:49.
- 10 SEQ ID NO:57 is the deduced amino acid sequence encoded by SEQ ID NO:50.
- SEQ ID NO:58 is the determined full-length cDNA sequence for clone L1058C (sequence of the originally isolated clone is given in SEQ ID NO:46 and the predicted extended cDNA sequence in SEQ ID NO:53).
- SEQ ID NO:59 is a first predicted ORF of SEQ ID NO:58.
- 15 SEQ ID NO:60 is a second predicted ORF of SEQ ID NO:58.
- SEQ ID NO:61 is the deduced amino acid sequence encoded by SEQ ID NO:59.
- SEQ ID NO:62 is the deduced amino acid sequence encoded by SEQ ID NO:60.
- 20 SEQ ID NO:63 is the determined cDNA sequence for Clone ID # 72761.
- SEQ ID NO:64 is the determined cDNA sequence for Clone ID # 72762.
- SEQ ID NO:65 is the determined cDNA sequence for Clone ID # 72763.
- SEQ ID NO:66 is the determined cDNA sequence for Clone ID # 72764.
- SEQ ID NO:67 is the determined cDNA sequence for Clone ID # 72765.
- 25 SEQ ID NO:68 is the determined cDNA sequence for Clone ID # 72766.
- SEQ ID NO:69 is the determined cDNA sequence for Clone ID # 72772.
- SEQ ID NO:70 is the determined cDNA sequence for Clone ID # 72775.
- SEQ ID NO:71 is the determined cDNA sequence for Clone ID # 72776.
- SEQ ID NO:72 is the determined cDNA sequence for Clone ID # 72779.
- 30 SEQ ID NO:73 is the determined cDNA sequence for Clone ID # 72781.
- SEQ ID NO:74 is the determined cDNA sequence for Clone ID # 72784.

- SEQ ID NO:75 is the determined cDNA sequence for Clone ID # 72788.
SEQ ID NO:76 is the determined cDNA sequence for Clone ID # 72789.
SEQ ID NO:77 is the determined cDNA sequence for Clone ID # 72790.
SEQ ID NO:78 is the determined cDNA sequence for Clone ID # 72791.
5 SEQ ID NO:79 is the determined cDNA sequence for Clone ID # 72792.
SEQ ID NO:80 is the determined cDNA sequence for Clone ID # 72794.
SEQ ID NO:81 is the determined cDNA sequence for Clone ID # 72795.
SEQ ID NO:82 is the determined cDNA sequence for Clone ID # 72797.
SEQ ID NO:83 is the determined cDNA sequence for Clone ID # 72798.
10 SEQ ID NO:84 is the determined cDNA sequence for Clone ID # 72804.
SEQ ID NO:85 is the determined cDNA sequence for Clone ID # 72805.
SEQ ID NO:86 is the determined cDNA sequence for Clone ID # 72806.
SEQ ID NO:87 is the determined cDNA sequence for Clone ID # 72807.
SEQ ID NO:88 is the determined cDNA sequence for Clone ID # 72808.
15 SEQ ID NO:89 is the determined cDNA sequence for Clone ID # 72809.
SEQ ID NO:90 is the determined cDNA sequence for Clone ID # 72811.
SEQ ID NO:91 is the determined full-length cDNA sequence for Clone
ID # 72813 which is named clone L1080C.
SEQ ID NO:92 is the deduced amino acid sequence encoded by SEQ ID
20 NO:91.
SEQ ID NO:93 is the ORF for L1027C.
SEQ ID NO:94 is a first determined full-length cDNA sequence for
L1027C.
SEQ ID NO:95 is a second determined full-length cDNA sequence for
25 L1027C.
SEQ ID NO:96 is the deduced amino acid sequence encoded by SEQ ID
NO:93.

DETAILED DESCRIPTION OF THE INVENTION

- The present invention is directed generally to compositions and their use
30 in the therapy and diagnosis of cancer, particularly lung cancer. As described further

below, illustrative compositions of the present invention include, but are not restricted to, polypeptides, particularly immunogenic polypeptides, polynucleotides encoding such polypeptides, antibodies and other binding agents, antigen presenting cells (APCs) and immune system cells (*e.g.*, T cells).

- 5 The practice of the present invention will employ, unless indicated specifically to the contrary, conventional methods of virology, immunology, microbiology, molecular biology and recombinant DNA techniques within the skill of the art, many of which are described below for the purpose of illustration. Such techniques are explained fully in the literature. See, *e.g.*, Sambrook, et al. Molecular Cloning: A Laboratory Manual (2nd Edition, 1989); Maniatis et al. Molecular Cloning: A Laboratory Manual (1982); DNA Cloning: A Practical Approach, vol. I & II (D. Glover, ed.); Oligonucleotide Synthesis (N. Gait, ed., 1984); Nucleic Acid Hybridization (B. Hames & S. Higgins, eds., 1985); Transcription and Translation (B. Hames & S. Higgins, eds., 1984); Animal Cell Culture (R. Freshney, ed., 1986); Perbal, A Practical Guide to Molecular Cloning (1984).
- 10 All publications, patents and patent applications cited herein, whether supra or infra, are hereby incorporated by reference in their entirety.
- 15 As used in this specification and the appended claims, the singular forms "a," "an" and "the" include plural references unless the content clearly dictates otherwise.
- 20

Polypeptide Compositions

- As used herein, the term "polypeptide" " is used in its conventional meaning, *i.e.*, as a sequence of amino acids. The polypeptides are not limited to a specific length of the product; thus, peptides, oligopeptides, and proteins are included within the definition of polypeptide, and such terms may be used interchangeably herein unless specifically indicated otherwise. This term also does not refer to or exclude post-expression modifications of the polypeptide, for example, glycosylations, acetylations, phosphorylations and the like, as well as other modifications known in the art, both naturally occurring and non-naturally occurring. A polypeptide may be an entire protein, or a subsequence thereof. Particular polypeptides of interest in the context of
- 25
- 30

this invention are amino acid subsequences comprising epitopes, i.e., antigenic determinants substantially responsible for the immunogenic properties of a polypeptide and being capable of evoking an immune response.

Particularly illustrative polypeptides of the present invention comprise
5 those encoded by a polynucleotide sequence set forth in any one of SEQ ID NO:1-35, 42-55, 58-60, 63-91 and 93-95, or a sequence that hybridizes under moderately stringent conditions, or, alternatively, under highly stringent conditions, to a polynucleotide sequence set forth in any one of SEQ ID NO:1-35, 42-55, 58-60, 63-91 and 93-95. Certain other illustrative polypeptides of the invention comprise amino acid sequences
10 as set forth in any one of SEQ ID NOs:36-41, 56, 57, 61, 62, 92 and 96.

The polypeptides of the present invention are sometimes herein referred to as lung tumor proteins or lung tumor polypeptides, as an indication that their identification has been based at least in part upon their increased levels of expression in lung tumor samples. Thus, a "lung tumor polypeptide" or "lung tumor protein," refers
15 generally to a polypeptide sequence of the present invention, or a polynucleotide sequence encoding such a polypeptide, that is expressed in a substantial proportion of lung tumor samples, for example preferably greater than about 20%, more preferably greater than about 30%, and most preferably greater than about 50% or more of lung tumor samples tested, at a level that is at least two fold, and preferably at least five fold,
20 greater than the level of expression in normal tissues, as determined using a representative assay provided herein. A lung tumor polypeptide sequence of the invention, based upon its increased level of expression in tumor cells, has particular utility both as a diagnostic marker as well as a therapeutic target, as further described below.

In certain preferred embodiments, the polypeptides of the invention are
25 immunogenic, i.e., they react detectably within an immunoassay (such as an ELISA or T-cell stimulation assay) with antisera and/or T-cells from a patient with lung cancer. Screening for immunogenic activity can be performed using techniques well known to the skilled artisan. For example, such screens can be performed using methods such as
30 those described in Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, 1988. In one illustrative example, a polypeptide may be

immobilized on a solid support and contacted with patient sera to allow binding of antibodies within the sera to the immobilized polypeptide. Unbound sera may then be removed and bound antibodies detected using, for example, ¹²⁵I-labeled Protein A.

As would be recognized by the skilled artisan, immunogenic portions of the polypeptides disclosed herein are also encompassed by the present invention. An “immunogenic portion,” as used herein, is a fragment of an immunogenic polypeptide of the invention that itself is immunologically reactive (*i.e.*, specifically binds) with the B-cells and/or T-cell surface antigen receptors that recognize the polypeptide. Immunogenic portions may generally be identified using well known techniques, such as those summarized in Paul, *Fundamental Immunology*, 3rd ed., 243-247 (Raven Press, 1993) and references cited therein. Such techniques include screening polypeptides for the ability to react with antigen-specific antibodies, antisera and/or T-cell lines or clones. As used herein, antisera and antibodies are “antigen-specific” if they specifically bind to an antigen (*i.e.*, they react with the protein in an ELISA or other immunoassay, and do not react detectably with unrelated proteins). Such antisera and antibodies may be prepared as described herein, and using well-known techniques.

In one preferred embodiment, an immunogenic portion of a polypeptide of the present invention is a portion that reacts with antisera and/or T-cells at a level that is not substantially less than the reactivity of the full-length polypeptide (*e.g.*, in an ELISA and/or T-cell reactivity assay). Preferably, the level of immunogenic activity of the immunogenic portion is at least about 50%, preferably at least about 70% and most preferably greater than about 90% of the immunogenicity for the full-length polypeptide. In some instances, preferred immunogenic portions will be identified that have a level of immunogenic activity greater than that of the corresponding full-length polypeptide, *e.g.*, having greater than about 100% or 150% or more immunogenic activity.

In certain other embodiments, illustrative immunogenic portions may include peptides in which an N-terminal leader sequence and/or transmembrane domain have been deleted. Other illustrative immunogenic portions will contain a small N- and/or C-terminal deletion (*e.g.*, 1-30 amino acids, preferably 5-15 amino acids), relative to the mature protein.

In another embodiment, a polypeptide composition of the invention may also comprise one or more polypeptides that are immunologically reactive with T cells and/or antibodies generated against a polypeptide of the invention, particularly a polypeptide having an amino acid sequence disclosed herein, or to an immunogenic
5 fragment or variant thereof.

In another embodiment of the invention, polypeptides are provided that comprise one or more polypeptides that are capable of eliciting T cells and/or antibodies that are immunologically reactive with one or more polypeptides described herein, or one or more polypeptides encoded by contiguous nucleic acid sequences contained in
10 the polynucleotide sequences disclosed herein, or immunogenic fragments or variants thereof, or to one or more nucleic acid sequences which hybridize to one or more of these sequences under conditions of moderate to high stringency.

The present invention, in another aspect, provides polypeptide fragments comprising at least about 5, 10, 15, 20, 25, 50, or 100 contiguous amino acids, or more,
15 including all intermediate lengths, of a polypeptide compositions set forth herein, such as those set forth in SEQ ID NOs:36-41, 56, 57, 61, 62, 92 and 96, or those encoded by a polynucleotide sequence set forth in a sequence of SEQ ID NOs:1-35, 42-55, 58-60, 63-91 and 93-95.

In another aspect, the present invention provides variants of the
20 polypeptide compositions described herein. Polypeptide variants generally encompassed by the present invention will typically exhibit at least about 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% or more identity (determined as described below), along its length, to a polypeptide sequences set forth herein.

In one preferred embodiment, the polypeptide fragments and variants
25 provided by the present invention are immunologically reactive with an antibody and/or T-cell that react with a full-length polypeptide specifically set forth herein.

In another preferred embodiment, the polypeptide fragments and variants provided by the present invention exhibit a level of immunogenic activity of at least
30 about 50%, preferably at least about 70%, and most preferably at least about 90% or

more of that exhibited by a full-length polypeptide sequence specifically set forth herein.

A polypeptide "variant," as the term is used herein, is a polypeptide that typically differs from a polypeptide specifically disclosed herein in one or more substitutions, deletions, additions and/or insertions. Such variants may be naturally occurring or may be synthetically generated, for example, by modifying one or more of the above polypeptide sequences of the invention and evaluating their immunogenic activity as described herein and/or using any of a number of techniques well known in the art.

For example, certain illustrative variants of the polypeptides of the invention include those in which one or more portions, such as an N-terminal leader sequence or transmembrane domain, have been removed. Other illustrative variants include variants in which a small portion (*e.g.*, 1-30 amino acids, preferably 5-15 amino acids) has been removed from the N- and/or C-terminal of the mature protein.

In many instances, a variant will contain conservative substitutions. A "conservative substitution" is one in which an amino acid is substituted for another amino acid that has similar properties, such that one skilled in the art of peptide chemistry would expect the secondary structure and hydrophobic nature of the polypeptide to be substantially unchanged. As described above, modifications may be made in the structure of the polynucleotides and polypeptides of the present invention and still obtain a functional molecule that encodes a variant or derivative polypeptide with desirable characteristics, *e.g.*, with immunogenic characteristics. When it is desired to alter the amino acid sequence of a polypeptide to create an equivalent, or even an improved, immunogenic variant or portion of a polypeptide of the invention, one skilled in the art will typically change one or more of the codons of the encoding DNA sequence according to Table 1.

For example, certain amino acids may be substituted for other amino acids in a protein structure without appreciable loss of interactive binding capacity with structures such as, for example, antigen-binding regions of antibodies or binding sites on substrate molecules. Since it is the interactive capacity and nature of a protein that defines that protein's biological functional activity, certain amino acid sequence

substitutions can be made in a protein sequence, and, of course, its underlying DNA coding sequence, and nevertheless obtain a protein with like properties. It is thus contemplated that various changes may be made in the peptide sequences of the disclosed compositions, or corresponding DNA sequences which encode said peptides without appreciable loss of their biological utility or activity.

Table 1

Amino Acids			Codons						
Alanine	Ala	A	GCA	GCC	GCG	GCU			
Cysteine	Cys	C	UGC	UGU					
Aspartic acid	Asp	D	GAC	GAU					
Glutamic acid	Glu	E	GAA	GAG					
Phenylalanine	Phe	F	UUC	UUU					
Glycine	Gly	G	GGA	GGC	GGG	GGU			
Histidine	His	H	CAC	CAU					
Isoleucine	Ile	I	AUA	AUC	AUU				
Lysine	Lys	K	AAA	AAG					
Leucine	Leu	L	UUA	UUG	CUA	CUC	CUG	CUU	
Methionine	Met	M	AUG						
Asparagine	Asn	N	AAC	AAU					
Proline	Pro	P	CCA	CCC	CCG	CCU			
Glutamine	Gln	Q	CAA	CAG					
Arginine	Arg	R	AGA	AGG	CGA	CGC	CGG	CGU	
Serine	Ser	S	AGC	AGU	UCA	UCC	UCG	UCU	
Threonine	Thr	T	ACA	ACC	ACG	ACU			
Valine	Val	V	GUA	GUC	GUG	GUU			
Tryptophan	Trp	W	UGG						
Tyrosine	Tyr	Y	UAC	UAU					

In making such changes, the hydropathic index of amino acids may be considered. The importance of the hydropathic amino acid index in conferring

interactive biologic function on a protein is generally understood in the art (Kyte and Doolittle, 1982, incorporated herein by reference). It is accepted that the relative hydrophathic character of the amino acid contributes to the secondary structure of the resultant protein, which in turn defines the interaction of the protein with other molecules, for example, enzymes, substrates, receptors, DNA, antibodies, antigens, and the like. Each amino acid has been assigned a hydrophathic index on the basis of its hydrophobicity and charge characteristics (Kyte and Doolittle, 1982). These values are: isoleucine (+4.5); valine (+4.2); leucine (+3.8); phenylalanine (+2.8); cysteine/cystine (+2.5); methionine (+1.9); alanine (+1.8); glycine (-0.4); threonine (-0.7); serine (-0.8); tryptophan (-0.9); tyrosine (-1.3); proline (-1.6); histidine (-3.2); glutamate (-3.5); glutamine (-3.5); aspartate (-3.5); asparagine (-3.5); lysine (-3.9); and arginine (-4.5).

It is known in the art that certain amino acids may be substituted by other amino acids having a similar hydrophathic index or score and still result in a protein with similar biological activity, *i.e.* still obtain a biological functionally equivalent protein. In making such changes, the substitution of amino acids whose hydrophathic indices are within ± 2 is preferred, those within ± 1 are particularly preferred, and those within ± 0.5 are even more particularly preferred. It is also understood in the art that the substitution of like amino acids can be made effectively on the basis of hydrophilicity. U. S. Patent 4,554,101 (specifically incorporated herein by reference in its entirety), states that the greatest local average hydrophilicity of a protein, as governed by the hydrophilicity of its adjacent amino acids, correlates with a biological property of the protein.

As detailed in U. S. Patent 4,554,101, the following hydrophilicity values have been assigned to amino acid residues: arginine (+3.0); lysine (+3.0); aspartate (+3.0 \pm 1); glutamate (+3.0 \pm 1); serine (+0.3); asparagine (+0.2); glutamine (+0.2); glycine (0); threonine (-0.4); proline (-0.5 \pm 1); alanine (-0.5); histidine (-0.5); cysteine (-1.0); methionine (-1.3); valine (-1.5); leucine (-1.8); isoleucine (-1.8); tyrosine (-2.3); phenylalanine (-2.5); tryptophan (-3.4). It is understood that an amino acid can be substituted for another having a similar hydrophilicity value and still obtain a biologically equivalent, and in particular, an immunologically equivalent protein. In such changes, the substitution of amino acids whose hydrophilicity values are within ± 2

is preferred, those within ± 1 are particularly preferred, and those within ± 0.5 are even more particularly preferred.

As outlined above, amino acid substitutions are generally therefore based on the relative similarity of the amino acid side-chain substituents, for example, their hydrophobicity, hydrophilicity, charge, size, and the like. Exemplary substitutions that take various of the foregoing characteristics into consideration are well known to those of skill in the art and include: arginine and lysine; glutamate and aspartate; serine and threonine; glutamine and asparagine; and valine, leucine and isoleucine.

In addition, any polynucleotide may be further modified to increase stability *in vivo*. Possible modifications include, but are not limited to, the addition of flanking sequences at the 5' and/or 3' ends; the use of phosphorothioate or 2' O-methyl rather than phosphodiesterase linkages in the backbone; and/or the inclusion of nontraditional bases such as inosine, queosine and wybutosine, as well as acetyl-methyl-, thio- and other modified forms of adenine, cytidine, guanine, thymine and uridine.

Amino acid substitutions may further be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity and/or the amphipathic nature of the residues. For example, negatively charged amino acids include aspartic acid and glutamic acid; positively charged amino acids include lysine and arginine; and amino acids with uncharged polar head groups having similar hydrophilicity values include leucine, isoleucine and valine; glycine and alanine; asparagine and glutamine; and serine, threonine, phenylalanine and tyrosine. Other groups of amino acids that may represent conservative changes include: (1) ala, pro, gly, glu, asp, gln, asn, ser, thr; (2) cys, ser, tyr, thr; (3) val, ile, leu, met, ala, phe; (4) lys, arg, his; and (5) phe, tyr, trp, his. A variant may also, or alternatively, contain nonconservative changes. In a preferred embodiment, variant polypeptides differ from a native sequence by substitution, deletion or addition of five amino acids or fewer. Variants may also (or alternatively) be modified by, for example, the deletion or addition of amino acids that have minimal influence on the immunogenicity, secondary structure and hydropathic nature of the polypeptide.

As noted above, polypeptides may comprise a signal (or leader) sequence at the N-terminal end of the protein, which co-translationally or post-translationally directs transfer of the protein. The polypeptide may also be conjugated to a linker or other sequence for ease of synthesis, purification or identification of the polypeptide (e.g., poly-His), or to enhance binding of the polypeptide to a solid support. For example, a polypeptide may be conjugated to an immunoglobulin Fc region.

When comparing polypeptide sequences, two sequences are said to be "identical" if the sequence of amino acids in the two sequences is the same when aligned for maximum correspondence, as described below. Comparisons between two sequences are typically performed by comparing the sequences over a comparison window to identify and compare local regions of sequence similarity. A "comparison window" as used herein, refers to a segment of at least about 20 contiguous positions, usually 30 to about 75, 40 to about 50, in which a sequence may be compared to a reference sequence of the same number of contiguous positions after the two sequences are optimally aligned.

Optimal alignment of sequences for comparison may be conducted using the Megalign program in the Lasergene suite of bioinformatics software (DNASTAR, Inc., Madison, WI), using default parameters. This program embodies several alignment schemes described in the following references: Dayhoff, M.O. (1978) A model of evolutionary change in proteins – Matrices for detecting distant relationships. In Dayhoff, M.O. (ed.) Atlas of Protein Sequence and Structure, National Biomedical Research Foundation, Washington DC Vol. 5, Suppl. 3, pp. 345-358; Hein J. (1990) Unified Approach to Alignment and Phylogenies pp. 626-645 *Methods in Enzymology* vol. 183, Academic Press, Inc., San Diego, CA; Higgins, D.G. and Sharp, P.M. (1989) *CABIOS* 5:151-153; Myers, E.W. and Muller W. (1988) *CABIOS* 4:11-17; Robinson, E.D. (1971) *Comb. Theor* 11:105; Saitou, N. Nei, M. (1987) *Mol. Biol. Evol.* 4:406-425; Sneath, P.H.A. and Sokal, R.R. (1973) *Numerical Taxonomy – the Principles and Practice of Numerical Taxonomy*, Freeman Press, San Francisco, CA; Wilbur, W.J. and Lipman, D.J. (1983) *Proc. Natl. Acad., Sci. USA* 80:726-730.

Alternatively, optimal alignment of sequences for comparison may be conducted by the local identity algorithm of Smith and Waterman (1981) *Add. APL*.

Math 2:482, by the identity alignment algorithm of Needleman and Wunsch (1970) *J. Mol. Biol.* 48:443, by the search for similarity methods of Pearson and Lipman (1988) *Proc. Natl. Acad. Sci. USA* 85: 2444, by computerized implementations of these algorithms (GAP, BESTFIT, BLAST, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group (GCG), 575 Science Dr., Madison, WI), or by inspection.

One preferred example of algorithms that are suitable for determining percent sequence identity and sequence similarity are the BLAST and BLAST 2.0 algorithms, which are described in Altschul et al. (1977) *Nucl. Acids Res.* 25:3389-3402 and Altschul et al. (1990) *J. Mol. Biol.* 215:403-410, respectively. BLAST and BLAST 2.0 can be used, for example with the parameters described herein, to determine percent sequence identity for the polynucleotides and polypeptides of the invention. Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information. For amino acid sequences, a scoring matrix can be used to calculate the cumulative score. Extension of the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters W, T and X determine the sensitivity and speed of the alignment.

In one preferred approach, the "percentage of sequence identity" is determined by comparing two optimally aligned sequences over a window of comparison of at least 20 positions, wherein the portion of the polypeptide sequence in the comparison window may comprise additions or deletions (*i.e.*, gaps) of 20 percent or less, usually 5 to 15 percent, or 10 to 12 percent, as compared to the reference sequences (which does not comprise additions or deletions) for optimal alignment of the two sequences. The percentage is calculated by determining the number of positions at which the identical amino acid residue occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the reference sequence (*i.e.*, the window size) and multiplying the results by 100 to yield the percentage of sequence identity.

Within other illustrative embodiments, a polypeptide may be a xenogeneic polypeptide that comprises an polypeptide having substantial sequence identity, as described above, to the human polypeptide (also termed autologous antigen) which served as a reference polypeptide, but which xenogeneic polypeptide is derived from a different, non-human species. One skilled in the art will recognize that “self” antigens are often poor stimulators of CD8+ and CD4+ T-lymphocyte responses, and therefore efficient immunotherapeutic strategies directed against tumor polypeptides require the development of methods to overcome immune tolerance to particular self tumor polypeptides. For example, humans immunized with prostate protein from a xenogeneic (non human) origin are capable of mounting an immune response against the counterpart human protein, e.g. the human prostate tumor protein present on human tumor cells. Accordingly, the present invention provides methods for purifying the xenogeneic form of the tumor proteins set forth herein, such as the polypeptides set forth in SEQ ID NO:36-41, 56, 57, 61, 62, 92 and 96, or those encoded by polynucleotide sequences set forth in SEQ ID NO:1-35, 42-55, 58-60, 63-91 and 93-95.

Therefore, one aspect of the present invention provides xenogeneic variants of the polypeptide compositions described herein. Such xenogeneic variants generally encompassed by the present invention will typically exhibit at least about 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% or more identity along their lengths, to a polypeptide sequences set forth herein.

More particularly, the invention is directed to mouse, rat, monkey, porcine and other non-human polypeptides which can be used as xenogeneic forms of human polypeptides set forth herein, to induce immune responses directed against tumor polypeptides of the invention.

Within other illustrative embodiments, a polypeptide may be a fusion polypeptide that comprises multiple polypeptides as described herein, or that comprises at least one polypeptide as described herein and an unrelated sequence, such as a known tumor protein. A fusion partner may, for example, assist in providing T helper epitopes (an immunological fusion partner), preferably T helper epitopes recognized by humans, or may assist in expressing the protein (an expression enhancer) at higher yields than the

native recombinant protein. Certain preferred fusion partners are both immunological and expression enhancing fusion partners. Other fusion partners may be selected so as to increase the solubility of the polypeptide or to enable the polypeptide to be targeted to desired intracellular compartments. Still further fusion partners include affinity tags, which facilitate purification of the polypeptide.

Fusion polypeptides may generally be prepared using standard techniques, including chemical conjugation. Preferably, a fusion polypeptide is expressed as a recombinant polypeptide, allowing the production of increased levels, relative to a non-fused polypeptide, in an expression system. Briefly, DNA sequences encoding the polypeptide components may be assembled separately, and ligated into an appropriate expression vector. The 3' end of the DNA sequence encoding one polypeptide component is ligated, with or without a peptide linker, to the 5' end of a DNA sequence encoding the second polypeptide component so that the reading frames of the sequences are in phase. This permits translation into a single fusion polypeptide that retains the biological activity of both component polypeptides.

A peptide linker sequence may be employed to separate the first and second polypeptide components by a distance sufficient to ensure that each polypeptide folds into its secondary and tertiary structures. Such a peptide linker sequence is incorporated into the fusion polypeptide using standard techniques well known in the art. Suitable peptide linker sequences may be chosen based on the following factors: (1) their ability to adopt a flexible extended conformation; (2) their inability to adopt a secondary structure that could interact with functional epitopes on the first and second polypeptides; and (3) the lack of hydrophobic or charged residues that might react with the polypeptide functional epitopes. Preferred peptide linker sequences contain Gly, Asn and Ser residues. Other near neutral amino acids, such as Thr and Ala may also be used in the linker sequence. Amino acid sequences which may be usefully employed as linkers include those disclosed in Maratea et al., *Gene* 40:39-46, 1985; Murphy et al., *Proc. Natl. Acad. Sci. USA* 83:8258-8262, 1986; U.S. Patent No. 4,935,233 and U.S. Patent No. 4,751,180. The linker sequence may generally be from 1 to about 50 amino acids in length. Linker sequences are not required when the first and second

polypeptides have non-essential N-terminal amino acid regions that can be used to separate the functional domains and prevent steric interference.

The ligated DNA sequences are operably linked to suitable transcriptional or translational regulatory elements. The regulatory elements responsible for expression of DNA are located only 5' to the DNA sequence encoding the first polypeptides. Similarly, stop codons required to end translation and transcription termination signals are only present 3' to the DNA sequence encoding the second polypeptide.

The fusion polypeptide can comprise a polypeptide as described herein together with an unrelated immunogenic protein, such as an immunogenic protein capable of eliciting a recall response. Examples of such proteins include tetanus, tuberculosis and hepatitis proteins (*see, for example, Stoute et al. New Engl. J. Med., 336:86-91, 1997*).

In one preferred embodiment, the immunological fusion partner is derived from a *Mycobacterium* sp., such as a *Mycobacterium tuberculosis*-derived Ra12 fragment. Ra12 compositions and methods for their use in enhancing the expression and/or immunogenicity of heterologous polynucleotide/polypeptide sequences is described in U.S. Patent Application 60/158,585, the disclosure of which is incorporated herein by reference in its entirety. Briefly, Ra12 refers to a polynucleotide region that is a subsequence of a *Mycobacterium tuberculosis* MTB32A nucleic acid. MTB32A is a serine protease of 32 KD molecular weight encoded by a gene in virulent and avirulent strains of *M. tuberculosis*. The nucleotide sequence and amino acid sequence of MTB32A have been described (for example, U.S. Patent Application 60/158,585; *see also, Skeiky et al., Infection and Immun. (1999) 67:3998-4007*, incorporated herein by reference). C-terminal fragments of the MTB32A coding sequence express at high levels and remain as a soluble polypeptides throughout the purification process. Moreover, Ra12 may enhance the immunogenicity of heterologous immunogenic polypeptides with which it is fused. One preferred Ra12 fusion polypeptide comprises a 14 KD C-terminal fragment corresponding to amino acid residues 192 to 323 of MTB32A. Other preferred Ra12 polynucleotides generally comprise at least about 15 consecutive nucleotides, at least about 30 nucleotides, at least

about 60 nucleotides, at least about 100 nucleotides, at least about 200 nucleotides, or at least about 300 nucleotides that encode a portion of a Ra12 polypeptide. Ra12 polynucleotides may comprise a native sequence (*i.e.*, an endogenous sequence that encodes a Ra12 polypeptide or a portion thereof) or may comprise a variant of such a sequence. Ra12 polynucleotide variants may contain one or more substitutions, additions, deletions and/or insertions such that the biological activity of the encoded fusion polypeptide is not substantially diminished, relative to a fusion polypeptide comprising a native Ra12 polypeptide. Variants preferably exhibit at least about 70% identity, more preferably at least about 80% identity and most preferably at least about 90% identity to a polynucleotide sequence that encodes a native Ra12 polypeptide or a portion thereof.

Within other preferred embodiments, an immunological fusion partner is derived from protein D, a surface protein of the gram-negative bacterium *Haemophilus influenza B* (WO 91/18926). Preferably, a protein D derivative comprises approximately the first third of the protein (*e.g.*, the first N-terminal 100-110 amino acids), and a protein D derivative may be lipidated. Within certain preferred embodiments, the first 109 residues of a Lipoprotein D fusion partner is included on the N-terminus to provide the polypeptide with additional exogenous T-cell epitopes and to increase the expression level in *E. coli* (thus functioning as an expression enhancer). The lipid tail ensures optimal presentation of the antigen to antigen presenting cells. Other fusion partners include the non-structural protein from influenzae virus, NS1 (hemagglutinin). Typically, the N-terminal 81 amino acids are used, although different fragments that include T-helper epitopes may be used.

In another embodiment, the immunological fusion partner is the protein known as LYTA, or a portion thereof (preferably a C-terminal portion). LYTA is derived from *Streptococcus pneumoniae*, which synthesizes an N-acetyl-L-alanine amidase known as amidase LYTA (encoded by the *LytA* gene; *Gene* 43:265-292, 1986). LYTA is an autolysin that specifically degrades certain bonds in the peptidoglycan backbone. The C-terminal domain of the LYTA protein is responsible for the affinity to the choline or to some choline analogues such as DEAE. This property has been exploited for the development of *E. coli* C-LYTA expressing plasmids useful for

expression of fusion proteins. Purification of hybrid proteins containing the C-LYTA fragment at the amino terminus has been described (*see Biotechnology 10:795-798, 1992*). Within a preferred embodiment, a repeat portion of LYTA may be incorporated into a fusion polypeptide. A repeat portion is found in the C-terminal region starting at residue 178. A particularly preferred repeat portion incorporates residues 188-305.

Yet another illustrative embodiment involves fusion polypeptides, and the polynucleotides encoding them, wherein the fusion partner comprises a targeting signal capable of directing a polypeptide to the endosomal/lysosomal compartment, as described in U.S. Patent No. 5,633,234. An immunogenic polypeptide of the invention, when fused with this targeting signal, will associate more efficiently with MHC class II molecules and thereby provide enhanced in vivo stimulation of CD4⁺ T-cells specific for the polypeptide.

Polypeptides of the invention are prepared using any of a variety of well known synthetic and/or recombinant techniques, the latter of which are further described below. Polypeptides, portions and other variants generally less than about 150 amino acids can be generated by synthetic means, using techniques well known to those of ordinary skill in the art. In one illustrative example, such polypeptides are synthesized using any of the commercially available solid-phase techniques, such as the Merrifield solid-phase synthesis method, where amino acids are sequentially added to a growing amino acid chain. *See Merrifield, J. Am. Chem. Soc. 85:2149-2146, 1963*. Equipment for automated synthesis of polypeptides is commercially available from suppliers such as Perkin Elmer/Applied BioSystems Division (Foster City, CA), and may be operated according to the manufacturer's instructions.

In general, polypeptide compositions (including fusion polypeptides) of the invention are isolated. An "isolated" polypeptide is one that is removed from its original environment. For example, a naturally-occurring protein or polypeptide is isolated if it is separated from some or all of the coexisting materials in the natural system. Preferably, such polypeptides are also purified, *e.g.*, are at least about 90% pure, more preferably at least about 95% pure and most preferably at least about 99% pure.

Polynucleotide Compositions

The present invention, in other aspects, provides polynucleotide compositions. The terms "DNA" and "polynucleotide" are used essentially interchangeably herein to refer to a DNA molecule that has been isolated free of total genomic DNA of a particular species. "Isolated," as used herein, means that a polynucleotide is substantially away from other coding sequences, and that the DNA molecule does not contain large portions of unrelated coding DNA, such as large chromosomal fragments or other functional genes or polypeptide coding regions. Of course, this refers to the DNA molecule as originally isolated, and does not exclude genes or coding regions later added to the segment by the hand of man.

As will be understood by those skilled in the art, the polynucleotide compositions of this invention can include genomic sequences, extra-genomic and plasmid-encoded sequences and smaller engineered gene segments that express, or may be adapted to express, proteins, polypeptides, peptides and the like. Such segments may be naturally isolated, or modified synthetically by the hand of man.

As will be also recognized by the skilled artisan, polynucleotides of the invention may be single-stranded (coding or antisense) or double-stranded, and may be DNA (genomic, cDNA or synthetic) or RNA molecules. RNA molecules may include HnRNA molecules, which contain introns and correspond to a DNA molecule in a one-to-one manner, and mRNA molecules, which do not contain introns. Additional coding or non-coding sequences may, but need not, be present within a polynucleotide of the present invention, and a polynucleotide may, but need not, be linked to other molecules and/or support materials.

Polynucleotides may comprise a native sequence (*i.e.*, an endogenous sequence that encodes a polypeptide/protein of the invention or a portion thereof) or may comprise a sequence that encodes a variant or derivative, preferably and immunogenic variant or derivative, of such a sequence.

Therefore, according to another aspect of the present invention, polynucleotide compositions are provided that comprise some or all of a polynucleotide sequence set forth in any one of SEQ ID NO:1-35, 42-55, 58-60, 63-91 and 93-95, complements of a polynucleotide sequence set forth in any one of SEQ ID NO:1-35, 42-

55, 58-60, 63-91 and 93-95, and degenerate variants of a polynucleotide sequence set forth in any one of SEQ ID NO:1-35, 42-55, 58-60, 63-91 and 93-95. In certain preferred embodiments, the polynucleotide sequences set forth herein encode immunogenic polypeptides, as described above.

5 In other related embodiments, the present invention provides polynucleotide variants having substantial identity to the sequences disclosed herein in SEQ ID NO:1-35, 42-55, 58-60, 63-91 and 93-95, for example those comprising at least 70% sequence identity, preferably at least 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% or higher, sequence identity compared to a polynucleotide sequence of this
10 invention using the methods described herein, (*e.g.*, BLAST analysis using standard parameters, as described below). One skilled in this art will recognize that these values can be appropriately adjusted to determine corresponding identity of proteins encoded by two nucleotide sequences by taking into account codon degeneracy, amino acid similarity, reading frame positioning and the like.

15 Typically, polynucleotide variants will contain one or more substitutions, additions, deletions and/or insertions, preferably such that the immunogenicity of the polypeptide encoded by the variant polynucleotide is not substantially diminished relative to a polypeptide encoded by a polynucleotide sequence specifically set forth herein). The term "variants" should also be understood to encompass homologous
20 genes of xenogenic origin.

In additional embodiments, the present invention provides polynucleotide fragments comprising or consisting of various lengths of contiguous stretches of sequence identical to or complementary to one or more of the sequences disclosed herein. For example, polynucleotides are provided by this invention that
25 comprise or consist of at least about 10, 15, 20, 30, 40, 50, 75, 100, 150, 200, 300, 400, 500 or 1000 or more contiguous nucleotides of one or more of the sequences disclosed herein as well as all intermediate lengths there between. It will be readily understood that "intermediate lengths", in this context, means any length between the quoted values, such as 16, 17, 18, 19, *etc.*; 21, 22, 23, *etc.*; 30, 31, 32, *etc.*; 50, 51, 52, 53, *etc.*;
30 100, 101, 102, 103, *etc.*; 150, 151, 152, 153, *etc.*; including all integers through 200-500; 500-1,000, and the like. A polynucleotide sequence as described here may be

extended at one or both ends by additional nucleotides not found in the native sequence. This additional sequence may consist of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 nucleotides at either end of the disclosed sequence or at both ends of the disclosed sequence.

5 In another embodiment of the invention, polynucleotide compositions are provided that are capable of hybridizing under moderate to high stringency conditions to a polynucleotide sequence provided herein, or a fragment thereof, or a complementary sequence thereof. Hybridization techniques are well known in the art of molecular biology. For purposes of illustration, suitable moderately stringent conditions for
10 testing the hybridization of a polynucleotide of this invention with other polynucleotides include prewashing in a solution of 5 X SSC, 0.5% SDS, 1.0 mM EDTA (pH 8.0); hybridizing at 50°C-60°C, 5 X SSC, overnight; followed by washing twice at 65°C for 20 minutes with each of 2X, 0.5X and 0.2X SSC containing 0.1% SDS. One skilled in the art will understand that the stringency of hybridization can be readily manipulated,
15 such as by altering the salt content of the hybridization solution and/or the temperature at which the hybridization is performed. For example, in another embodiment, suitable highly stringent hybridization conditions include those described above, with the exception that the temperature of hybridization is increased, *e.g.*, to 60-65°C or 65-70°C.

20 In certain preferred embodiments, the polynucleotides described above, *e.g.*, polynucleotide variants, fragments and hybridizing sequences, encode polypeptides that are immunologically cross-reactive with a polypeptide sequence specifically set forth herein. In other preferred embodiments, such polynucleotides encode polypeptides that have a level of immunogenic activity of at least about 50%, preferably
25 at least about 70%, and more preferably at least about 90% of that for a polypeptide sequence specifically set forth herein.

 The polynucleotides of the present invention, or fragments thereof, regardless of the length of the coding sequence itself, may be combined with other DNA sequences, such as promoters, polyadenylation signals, additional restriction enzyme
30 sites, multiple cloning sites, other coding segments, and the like, such that their overall length may vary considerably. It is therefore contemplated that a nucleic acid fragment

of almost any length may be employed, with the total length preferably being limited by the ease of preparation and use in the intended recombinant DNA protocol. For example, illustrative polynucleotide segments with total lengths of about 10,000, about 5000, about 3000, about 2,000, about 1,000, about 500, about 200, about 100, about 50
5 base pairs in length, and the like, (including all intermediate lengths) are contemplated to be useful in many implementations of this invention.

When comparing polynucleotide sequences, two sequences are said to be "identical" if the sequence of nucleotides in the two sequences is the same when aligned for maximum correspondence, as described below. Comparisons between two
10 sequences are typically performed by comparing the sequences over a comparison window to identify and compare local regions of sequence similarity. A "comparison window" as used herein, refers to a segment of at least about 20 contiguous positions, usually 30 to about 75, 40 to about 50, in which a sequence may be compared to a reference sequence of the same number of contiguous positions after the two sequences
15 are optimally aligned.

Optimal alignment of sequences for comparison may be conducted using the Megalign program in the Lasergene suite of bioinformatics software (DNASTAR, Inc., Madison, WI), using default parameters. This program embodies several alignment schemes described in the following references: Dayhoff, M.O. (1978) A
20 model of evolutionary change in proteins – Matrices for detecting distant relationships. In Dayhoff, M.O. (ed.) Atlas of Protein Sequence and Structure, National Biomedical Research Foundation, Washington DC Vol. 5, Suppl. 3, pp. 345-358; Hein J. (1990) Unified Approach to Alignment and Phylogenesis pp. 626-645 *Methods in Enzymology* vol. 183, Academic Press, Inc., San Diego, CA; Higgins, D.G. and Sharp, P.M. (1989)
25 *CABIOS* 5:151-153; Myers, E.W. and Muller W. (1988) *CABIOS* 4:11-17; Robinson, E.D. (1971) *Comb. Theor* 11:105; Santou, N. Nes, M. (1987) *Mol. Biol. Evol.* 4:406-425; Sneath, P.H.A. and Sokal, R.R. (1973) *Numerical Taxonomy – the Principles and Practice of Numerical Taxonomy*, Freeman Press, San Francisco, CA; Wilbur, W.J. and Lipman, D.J. (1983) *Proc. Natl. Acad., Sci. USA* 80:726-730.

30 Alternatively, optimal alignment of sequences for comparison may be conducted by the local identity algorithm of Smith and Waterman (1981) *Add. APL*.

Math 2:482, by the identity alignment algorithm of Needleman and Wunsch (1970) *J. Mol. Biol.* 48:443, by the search for similarity methods of Pearson and Lipman (1988) *Proc. Natl. Acad. Sci. USA* 85: 2444, by computerized implementations of these algorithms (GAP, BESTFIT, BLAST, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group (GCG), 575 Science Dr., Madison, WI), or by inspection.

One preferred example of algorithms that are suitable for determining percent sequence identity and sequence similarity are the BLAST and BLAST 2.0 algorithms, which are described in Altschul et al. (1977) *Nucl. Acids Res.* 25:3389-3402 and Altschul et al. (1990) *J. Mol. Biol.* 215:403-410, respectively. BLAST and BLAST 2.0 can be used, for example with the parameters described herein, to determine percent sequence identity for the polynucleotides of the invention. Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information. In one illustrative example, cumulative scores can be calculated using, for nucleotide sequences, the parameters M (reward score for a pair of matching residues; always >0) and N (penalty score for mismatching residues; always <0). Extension of the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters W, T and X determine the sensitivity and speed of the alignment. The BLASTN program (for nucleotide sequences) uses as defaults a wordlength (W) of 11, and expectation (E) of 10, and the BLOSUM62 scoring matrix (see Henikoff and Henikoff (1989) *Proc. Natl. Acad. Sci. USA* 89:10915) alignments, (B) of 50, expectation (E) of 10, M=5, N=-4 and a comparison of both strands.

Preferably, the "percentage of sequence identity" is determined by comparing two optimally aligned sequences over a window of comparison of at least 20 positions, wherein the portion of the polynucleotide sequence in the comparison window may comprise additions or deletions (*i.e.*, gaps) of 20 percent or less, usually 5 to 15 percent, or 10 to 12 percent, as compared to the reference sequences (which does not comprise additions or deletions) for optimal alignment of the two sequences. The

percentage is calculated by determining the number of positions at which the identical nucleic acid bases occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the reference sequence (*i.e.*, the window size) and multiplying the results by 100 to yield the percentage of sequence identity.

It will be appreciated by those of ordinary skill in the art that, as a result of the degeneracy of the genetic code, there are many nucleotide sequences that encode a polypeptide as described herein. Some of these polynucleotides bear minimal homology to the nucleotide sequence of any native gene. Nonetheless, polynucleotides that vary due to differences in codon usage are specifically contemplated by the present invention. Further, alleles of the genes comprising the polynucleotide sequences provided herein are within the scope of the present invention. Alleles are endogenous genes that are altered as a result of one or more mutations, such as deletions, additions and/or substitutions of nucleotides. The resulting mRNA and protein may, but need not, have an altered structure or function. Alleles may be identified using standard techniques (such as hybridization, amplification and/or database sequence comparison).

Therefore, in another embodiment of the invention, a mutagenesis approach, such as site-specific mutagenesis, is employed for the preparation of immunogenic variants and/or derivatives of the polypeptides described herein. By this approach, specific modifications in a polypeptide sequence can be made through mutagenesis of the underlying polynucleotides that encode them. These techniques provides a straightforward approach to prepare and test sequence variants, for example, incorporating one or more of the foregoing considerations, by introducing one or more nucleotide sequence changes into the polynucleotide.

Site-specific mutagenesis allows the production of mutants through the use of specific oligonucleotide sequences which encode the DNA sequence of the desired mutation, as well as a sufficient number of adjacent nucleotides, to provide a primer sequence of sufficient size and sequence complexity to form a stable duplex on both sides of the deletion junction being traversed. Mutations may be employed in a selected polynucleotide sequence to improve, alter, decrease, modify, or otherwise

change the properties of the polynucleotide itself, and/or alter the properties, activity, composition, stability, or primary sequence of the encoded polypeptide.

In certain embodiments of the present invention, the inventors contemplate the mutagenesis of the disclosed polynucleotide sequences to alter one or
5 more properties of the encoded polypeptide, such as the immunogenicity of a polypeptide vaccine. The techniques of site-specific mutagenesis are well-known in the art, and are widely used to create variants of both polypeptides and polynucleotides. For example, site-specific mutagenesis is often used to alter a specific portion of a DNA molecule. In such embodiments, a primer comprising typically about 14 to about 25
10 nucleotides or so in length is employed, with about 5 to about 10 residues on both sides of the junction of the sequence being altered.

As will be appreciated by those of skill in the art, site-specific mutagenesis techniques have often employed a phage vector that exists in both a single stranded and double stranded form. Typical vectors useful in site-directed mutagenesis
15 include vectors such as the M13 phage. These phage are readily commercially-available and their use is generally well-known to those skilled in the art. Double-stranded plasmids are also routinely employed in site directed mutagenesis that eliminates the step of transferring the gene of interest from a plasmid to a phage.

In general, site-directed mutagenesis in accordance herewith is
20 performed by first obtaining a single-stranded vector or melting apart of two strands of a double-stranded vector that includes within its sequence a DNA sequence that encodes the desired peptide. An oligonucleotide primer bearing the desired mutated sequence is prepared, generally synthetically. This primer is then annealed with the single-stranded vector, and subjected to DNA polymerizing enzymes such as *E. coli* polymerase I
25 Klenow fragment, in order to complete the synthesis of the mutation-bearing strand. Thus, a heteroduplex is formed wherein one strand encodes the original non-mutated sequence and the second strand bears the desired mutation. This heteroduplex vector is then used to transform appropriate cells, such as *E. coli* cells, and clones are selected which include recombinant vectors bearing the mutated sequence arrangement.

30 The preparation of sequence variants of the selected peptide-encoding DNA segments using site-directed mutagenesis provides a means of producing

potentially useful species and is not meant to be limiting as there are other ways in which sequence variants of peptides and the DNA sequences encoding them may be obtained. For example, recombinant vectors encoding the desired peptide sequence may be treated with mutagenic agents, such as hydroxylamine, to obtain sequence
5 variants. Specific details regarding these methods and protocols are found in the teachings of Maloy *et al.*, 1994; Segal, 1976; Prokop and Bajpai, 1991; Kubys, 1994; and Maniatis *et al.*, 1982, each incorporated herein by reference, for that purpose.

As used herein, the term "oligonucleotide directed mutagenesis procedure" refers to template-dependent processes and vector-mediated propagation
10 which result in an increase in the concentration of a specific nucleic acid molecule relative to its initial concentration, or in an increase in the concentration of a detectable signal, such as amplification. As used herein, the term "oligonucleotide directed mutagenesis procedure" is intended to refer to a process that involves the template-dependent extension of a primer molecule. The term template dependent
15 process refers to nucleic acid synthesis of a RNA or a DNA molecule wherein the sequence of the newly synthesized strand of nucleic acid is dictated by the well-known rules of complementary base pairing (see, for example, Watson, 1987). Typically, vector mediated methodologies involve the introduction of the nucleic acid fragment into a DNA or RNA vector, the clonal amplification of the vector, and the recovery of
20 the amplified nucleic acid fragment. Examples of such methodologies are provided by U. S. Patent No. 4,237,224, specifically incorporated herein by reference in its entirety.

In another approach for the production of polypeptide variants of the present invention, recursive sequence recombination, as described in U.S. Patent No. 5,837,458, may be employed. In this approach, iterative cycles of recombination and
25 screening or selection are performed to "evolve" individual polynucleotide variants of the invention having, for example, enhanced immunogenic activity.

In other embodiments of the present invention, the polynucleotide sequences provided herein can be advantageously used as probes or primers for nucleic acid hybridization. As such, it is contemplated that nucleic acid segments that comprise
30 or consist of a sequence region of at least about a 15 nucleotide long contiguous sequence that has the same sequence as, or is complementary to, a 15 nucleotide long

contiguous sequence disclosed herein will find particular utility. Longer contiguous identical or complementary sequences, *e.g.*, those of about 20, 30, 40, 50, 100, 200, 500, 1000 (including all intermediate lengths) and even up to full length sequences will also be of use in certain embodiments.

5 The ability of such nucleic acid probes to specifically hybridize to a sequence of interest will enable them to be of use in detecting the presence of complementary sequences in a given sample. However, other uses are also envisioned, such as the use of the sequence information for the preparation of mutant species primers, or primers for use in preparing other genetic constructions.

10 Polynucleotide molecules having sequence regions consisting of contiguous nucleotide stretches of 10-14, 15-20, 30, 50, or even of 100-200 nucleotides or so (including intermediate lengths as well), identical or complementary to a polynucleotide sequence disclosed herein, are particularly contemplated as hybridization probes for use in, *e.g.*, Southern and Northern blotting. This would allow a gene
15 product, or fragment thereof, to be analyzed, both in diverse cell types and also in various bacterial cells. The total size of fragment, as well as the size of the complementary stretch(es), will ultimately depend on the intended use or application of the particular nucleic acid segment. Smaller fragments will generally find use in hybridization embodiments, wherein the length of the contiguous complementary region
20 may be varied, such as between about 15 and about 100 nucleotides, but larger contiguous complementarity stretches may be used, according to the length complementary sequences one wishes to detect.

 The use of a hybridization probe of about 15-25 nucleotides in length allows the formation of a duplex molecule that is both stable and selective. Molecules
25 having contiguous complementary sequences over stretches greater than 15 bases in length are generally preferred, though, in order to increase stability and selectivity of the hybrid, and thereby improve the quality and degree of specific hybrid molecules obtained. One will generally prefer to design nucleic acid molecules having gene-complementary stretches of 15 to 25 contiguous nucleotides, or even longer where
30 desired.

Hybridization probes may be selected from any portion of any of the sequences disclosed herein. All that is required is to review the sequences set forth herein, or to any continuous portion of the sequences, from about 15-25 nucleotides in length up to and including the full length sequence, that one wishes to utilize as a probe or primer. The choice of probe and primer sequences may be governed by various factors. For example, one may wish to employ primers from towards the termini of the total sequence.

Small polynucleotide segments or fragments may be readily prepared by, for example, directly synthesizing the fragment by chemical means, as is commonly practiced using an automated oligonucleotide synthesizer. Also, fragments may be obtained by application of nucleic acid reproduction technology, such as the PCRTM technology of U. S. Patent 4,683,202 (incorporated herein by reference), by introducing selected sequences into recombinant vectors for recombinant production, and by other recombinant DNA techniques generally known to those of skill in the art of molecular biology.

The nucleotide sequences of the invention may be used for their ability to selectively form duplex molecules with complementary stretches of the entire gene or gene fragments of interest. Depending on the application envisioned, one will typically desire to employ varying conditions of hybridization to achieve varying degrees of selectivity of probe towards target sequence. For applications requiring high selectivity, one will typically desire to employ relatively stringent conditions to form the hybrids, *e.g.*, one will select relatively low salt and/or high temperature conditions, such as provided by a salt concentration of from about 0.02 M to about 0.15 M salt at temperatures of from about 50°C to about 70°C. Such selective conditions tolerate little, if any, mismatch between the probe and the template or target strand, and would be particularly suitable for isolating related sequences.

Of course, for some applications, for example, where one desires to prepare mutants employing a mutant primer strand hybridized to an underlying template, less stringent (reduced stringency) hybridization conditions will typically be needed in order to allow formation of the heteroduplex. In these circumstances, one may desire to employ salt conditions such as those of from about 0.15 M to about 0.9 M

salt, at temperatures ranging from about 20°C to about 55°C. Cross-hybridizing species can thereby be readily identified as positively hybridizing signals with respect to control hybridizations. In any case, it is generally appreciated that conditions can be rendered more stringent by the addition of increasing amounts of formamide, which serves to
5 destabilize the hybrid duplex in the same manner as increased temperature. Thus, hybridization conditions can be readily manipulated, and thus will generally be a method of choice depending on the desired results.

According to another embodiment of the present invention, polynucleotide compositions comprising antisense oligonucleotides are provided.
10 Antisense oligonucleotides have been demonstrated to be effective and targeted inhibitors of protein synthesis, and, consequently, provide a therapeutic approach by which a disease can be treated by inhibiting the synthesis of proteins that contribute to the disease. The efficacy of antisense oligonucleotides for inhibiting protein synthesis is well established. For example, the synthesis of polygalacturonase and the muscarine
15 type 2 acetylcholine receptor are inhibited by antisense oligonucleotides directed to their respective mRNA sequences (U. S. Patent 5,739,119 and U. S. Patent 5,759,829). Further, examples of antisense inhibition have been demonstrated with the nuclear protein cyclin, the multiple drug resistance gene (MDG1), ICAM-1, E-selectin, STK-1, striatal GABA_A receptor and human EGF (Jaskulski *et al.*, Science. 1988 Jun
20 10;240(4858):1544-6; Vasanthakumar and Ahmed, Cancer Commun. 1989;1(4):225-32; Peris *et al.*, Brain Res Mol Brain Res. 1998 Jun 15;57(2):310-20; U. S. Patent 5,801,154; U.S. Patent 5,789,573; U. S. Patent 5,718,709 and U.S. Patent 5,610,288). Antisense constructs have also been described that inhibit and can be used to treat a variety of abnormal cellular proliferations, *e.g.* cancer (U. S. Patent 5,747,470; U. S.
25 Patent 5,591,317 and U. S. Patent 5,783,683).

Therefore, in certain embodiments, the present invention provides oligonucleotide sequences that comprise all, or a portion of, any sequence that is capable of specifically binding to polynucleotide sequence described herein, or a complement thereof. In one embodiment, the antisense oligonucleotides comprise DNA
30 or derivatives thereof. In another embodiment, the oligonucleotides comprise RNA or derivatives thereof. In a third embodiment, the oligonucleotides are modified DNAs

comprising a phosphorothioated modified backbone. In a fourth embodiment, the oligonucleotide sequences comprise peptide nucleic acids or derivatives thereof. In each case, preferred compositions comprise a sequence region that is complementary, and more preferably substantially-complementary, and even more preferably, completely complementary to one or more portions of polynucleotides disclosed herein. Selection of antisense compositions specific for a given gene sequence is based upon analysis of the chosen target sequence and determination of secondary structure, T_m , binding energy, and relative stability. Antisense compositions may be selected based upon their relative inability to form dimers, hairpins, or other secondary structures that would reduce or prohibit specific binding to the target mRNA in a host cell. Highly preferred target regions of the mRNA, are those which are at or near the AUG translation initiation codon, and those sequences which are substantially complementary to 5' regions of the mRNA. These secondary structure analyses and target site selection considerations can be performed, for example, using v.4 of the OLIGO primer analysis software and/or the BLASTN 2.0.5 algorithm software (Altschul *et al.*, Nucleic Acids Res. 1997, 25(17):3389-402).

The use of an antisense delivery method employing a short peptide vector, termed MPG (27 residues), is also contemplated. The MPG peptide contains a hydrophobic domain derived from the fusion sequence of HIV gp41 and a hydrophilic domain from the nuclear localization sequence of SV40 T-antigen (Morris *et al.*, Nucleic Acids Res. 1997 Jul 15;25(14):2730-6). It has been demonstrated that several molecules of the MPG peptide coat the antisense oligonucleotides and can be delivered into cultured mammalian cells in less than 1 hour with relatively high efficiency (90%). Further, the interaction with MPG strongly increases both the stability of the oligonucleotide to nuclease and the ability to cross the plasma membrane.

According to another embodiment of the invention, the polynucleotide compositions described herein are used in the design and preparation of ribozyme molecules for inhibiting expression of the tumor polypeptides and proteins of the present invention in tumor cells. Ribozymes are RNA-protein complexes that cleave nucleic acids in a site-specific fashion. Ribozymes have specific catalytic domains that possess endonuclease activity (Kim and Cech, Proc Natl Acad Sci U S A. 1987

Dec;84(24):8788-92; Forster and Symons, Cell, 1987 Apr 24;49(2):211-20). For example, a large number of ribozymes accelerate phosphoester transfer reactions with a high degree of specificity, often cleaving only one of several phosphoesters in an oligonucleotide substrate (Cech *et al.*, Cell, 1981 Dec;27(3 Pt 2):487-96; Michel and Westhof, J Mol Biol. 1990 Dec 5;216(3):585-610; Reinhold-Hurek and Shub, Nature. 1992 May 14;357(6374):173-6). This specificity has been attributed to the requirement that the substrate bind via specific base-pairing interactions to the internal guide sequence ("IGS") of the ribozyme prior to chemical reaction.

Six basic varieties of naturally-occurring enzymatic RNAs are known presently. Each can catalyze the hydrolysis of RNA phosphodiester bonds *in trans* (and thus can cleave other RNA molecules) under physiological conditions. In general, enzymatic nucleic acids act by first binding to a target RNA. Such binding occurs through the target binding portion of a enzymatic nucleic acid which is held in close proximity to an enzymatic portion of the molecule that acts to cleave the target RNA. Thus, the enzymatic nucleic acid first recognizes and then binds a target RNA through complementary base-pairing, and once bound to the correct site, acts enzymatically to cut the target RNA. Strategic cleavage of such a target RNA will destroy its ability to direct synthesis of an encoded protein. After an enzymatic nucleic acid has bound and cleaved its RNA target, it is released from that RNA to search for another target and can repeatedly bind and cleave new targets.

The enzymatic nature of a ribozyme is advantageous over many technologies, such as antisense technology (where a nucleic acid molecule simply binds to a nucleic acid target to block its translation) since the concentration of ribozyme necessary to affect a therapeutic treatment is lower than that of an antisense oligonucleotide. This advantage reflects the ability of the ribozyme to act enzymatically. Thus, a single ribozyme molecule is able to cleave many molecules of target RNA. In addition, the ribozyme is a highly specific inhibitor, with the specificity of inhibition depending not only on the base pairing mechanism of binding to the target RNA, but also on the mechanism of target RNA cleavage. Single mismatches, or base-substitutions, near the site of cleavage can completely eliminate catalytic activity of a ribozyme. Similar mismatches in antisense molecules do not prevent their action

(Woolf *et al.*, Proc Natl Acad Sci U S A. 1992 Aug 15;89(16):7305-9). Thus, the specificity of action of a ribozyme is greater than that of an antisense oligonucleotide binding the same RNA site.

The enzymatic nucleic acid molecule may be formed in a hammerhead, hairpin, a hepatitis δ virus, group I intron or RNaseP RNA (in association with an RNA guide sequence) or Neurospora VS RNA motif. Examples of hammerhead motifs are described by Rossi *et al.* Nucleic Acids Res. 1992 Sep 11;20(17):4559-65. Examples of hairpin motifs are described by Hampel *et al.* (Eur. Pat. Appl. Publ. No. EP 0360257), Hampel and Tritz, Biochemistry 1989 Jun 13;28(12):4929-33; Hampel *et al.*, Nucleic Acids Res. 1990 Jan 25;18(2):299-304 and U. S. Patent 5,631,359. An example of the hepatitis δ virus motif is described by Perrotta and Been, Biochemistry. 1992 Dec 1;31(47):11843-52; an example of the RNaseP motif is described by Guerrier-Takada *et al.*, Cell. 1983 Dec;35(3 Pt 2):849-57; Neurospora VS RNA ribozyme motif is described by Collins (Saville and Collins, Cell. 1990 May 18;61(4):685-96; Saville and Collins, Proc Natl Acad Sci U S A. 1991 Oct 1;88(19):8826-30; Collins and Olive, Biochemistry. 1993 Mar 23;32(11):2795-9); and an example of the Group I intron is described in (U. S. Patent 4,987,071). All that is important in an enzymatic nucleic acid molecule of this invention is that it has a specific substrate binding site which is complementary to one or more of the target gene RNA regions, and that it have nucleotide sequences within or surrounding that substrate binding site which impart an RNA cleaving activity to the molecule. Thus the ribozyme constructs need not be limited to specific motifs mentioned herein.

Ribozymes may be designed as described in Int. Pat. Appl. Publ. No. WO 93/23569 and Int. Pat. Appl. Publ. No. WO 94/02595, each specifically incorporated herein by reference) and synthesized to be tested *in vitro* and *in vivo*, as described. Such ribozymes can also be optimized for delivery. While specific examples are provided, those in the art will recognize that equivalent RNA targets in other species can be utilized when necessary.

Ribozyme activity can be optimized by altering the length of the ribozyme binding arms, or chemically synthesizing ribozymes with modifications that prevent their degradation by serum ribonucleases (see *e.g.*, Int. Pat. Appl. Publ. No. WO

92/07065; Int. Pat. Appl. Publ. No. WO 93/15187; Int. Pat. Appl. Publ. No. WO 91/03162; Eur. Pat. Appl. Publ. No. 92110298.4; U. S. Patent 5,334,711; and Int. Pat. Appl. Publ. No. WO 94/13688, which describe various chemical modifications that can be made to the sugar moieties of enzymatic RNA molecules), modifications which enhance their efficacy in cells, and removal of stem II bases to shorten RNA synthesis times and reduce chemical requirements.

Sullivan *et al.* (Int. Pat. Appl. Publ. No. WO 94/02595) describes the general methods for delivery of enzymatic RNA molecules. Ribozymes may be administered to cells by a variety of methods known to those familiar to the art, including, but not restricted to, encapsulation in liposomes, by iontophoresis, or by incorporation into other vehicles, such as hydrogels, cyclodextrins, biodegradable nanocapsules, and bioadhesive microspheres. For some indications, ribozymes may be directly delivered *ex vivo* to cells or tissues with or without the aforementioned vehicles. Alternatively, the RNA/vehicle combination may be locally delivered by direct inhalation, by direct injection or by use of a catheter, infusion pump or stent. Other routes of delivery include, but are not limited to, intravascular, intramuscular, subcutaneous or joint injection, aerosol inhalation, oral (tablet or pill form), topical, systemic, ocular, intraperitoneal and/or intrathecal delivery. More detailed descriptions of ribozyme delivery and administration are provided in Int. Pat. Appl. Publ. No. WO 94/02595 and Int. Pat. Appl. Publ. No. WO 93/23569, each specifically incorporated herein by reference.

Another means of accumulating high concentrations of a ribozyme(s) within cells is to incorporate the ribozyme-encoding sequences into a DNA expression vector. Transcription of the ribozyme sequences are driven from a promoter for eukaryotic RNA polymerase I (pol I), RNA polymerase II (pol II), or RNA polymerase III (pol III). Transcripts from pol II or pol III promoters will be expressed at high levels in all cells; the levels of a given pol II promoter in a given cell type will depend on the nature of the gene regulatory sequences (enhancers, silencers, *etc.*) present nearby. Prokaryotic RNA polymerase promoters may also be used, providing that the prokaryotic RNA polymerase enzyme is expressed in the appropriate cells. Ribozymes expressed from such promoters have been shown to function in mammalian cells. Such

transcription units can be incorporated into a variety of vectors for introduction into mammalian cells, including but not restricted to, plasmid DNA vectors, viral DNA vectors (such as adenovirus or adeno-associated vectors), or viral RNA vectors (such as retroviral, semliki forest virus, sindbis virus vectors).

- 5 In another embodiment of the invention, peptide nucleic acids (PNAs) compositions are provided. PNA is a DNA mimic in which the nucleobases are attached to a pseudopeptide backbone (Good and Nielsen, *Antisense Nucleic Acid Drug*, Dev. 1997 7(4) 431-37). PNA is able to be utilized in a number of methods that traditionally have used RNA or DNA. Often PNA sequences perform better in
10 techniques than the corresponding RNA or DNA sequences and have utilities that are not inherent to RNA or DNA. A review of PNA including methods of making, characteristics of, and methods of using, is provided by Corey (*Trends Biotechnol* 1997 Jun;15(6):224-9). As such, in certain embodiments, one may prepare PNA sequences that are complementary to one or more portions of the ACE mRNA sequence, and such
15 PNA compositions may be used to regulate, alter, decrease, or reduce the translation of ACE-specific mRNA, and thereby alter the level of ACE activity in a host cell to which such PNA compositions have been administered.

- PNAs have 2-aminoethyl-glycine linkages replacing the normal phosphodiester backbone of DNA (Nielsen *et al.*, *Science* 1991 Dec 6;254(5037):1497-
20 500; Hanvey *et al.*, *Science*. 1992 Nov 27;258(5087):1481-5; Hyrup and Nielsen, *Bioorg Med Chem*. 1996 Jan;4(1):5-23). This chemistry has three important consequences: firstly, in contrast to DNA or phosphorothioate oligonucleotides, PNAs are neutral molecules; secondly, PNAs are achiral, which avoids the need to develop a stereoselective synthesis; and thirdly, PNA synthesis uses standard Boc or Fmoc
25 protocols for solid-phase peptide synthesis, although other methods, including a modified Merrifield method, have been used.

- PNA monomers or ready-made oligomers are commercially available from PerSeptive Biosystems (Framingham, MA). PNA syntheses by either Boc or Fmoc protocols are straightforward using manual or automated protocols (Norton *et al.*,
30 *Bioorg Med Chem*. 1995 Apr;3(4):437-45). The manual protocol lends itself to the

production of chemically modified PNAs or the simultaneous synthesis of families of closely related PNAs.

As with peptide synthesis, the success of a particular PNA synthesis will depend on the properties of the chosen sequence. For example, while in theory PNAs can incorporate any combination of nucleotide bases, the presence of adjacent purines can lead to deletions of one or more residues in the product. In expectation of this difficulty, it is suggested that, in producing PNAs with adjacent purines, one should repeat the coupling of residues likely to be added inefficiently. This should be followed by the purification of PNAs by reverse-phase high-pressure liquid chromatography, providing yields and purity of product similar to those observed during the synthesis of peptides.

Modifications of PNAs for a given application may be accomplished by coupling amino acids during solid-phase synthesis or by attaching compounds that contain a carboxylic acid group to the exposed N-terminal amine. Alternatively, PNAs can be modified after synthesis by coupling to an introduced lysine or cysteine. The ease with which PNAs can be modified facilitates optimization for better solubility or for specific functional requirements. Once synthesized, the identity of PNAs and their derivatives can be confirmed by mass spectrometry. Several studies have made and utilized modifications of PNAs (for example, Norton *et al.*, *Bioorg Med Chem.* 1995 Apr;3(4):437-45; Petersen *et al.*, *J Pept Sci.* 1995 May-Jun;1(3):175-83; Orum *et al.*, *Biotechniques.* 1995 Sep;19(3):472-80; Footer *et al.*, *Biochemistry.* 1996 Aug 20;35(33):10673-9; Griffith *et al.*, *Nucleic Acids Res.* 1995 Aug 11;23(15):3003-8; Pardridge *et al.*, *Proc Natl Acad Sci U S A.* 1995 Jun 6;92(12):5592-6; Boffa *et al.*, *Proc Natl Acad Sci U S A.* 1995 Mar 14;92(6):1901-5; Gambacorti-Passerini *et al.*, *Blood.* 1996 Aug 15;88(4):1411-7; Armitage *et al.*, *Proc Natl Acad Sci U S A.* 1997 Nov 11;94(23):12320-5; Seeger *et al.*, *Biotechniques.* 1997 Sep;23(3):512-7). U.S. Patent No. 5,700,922 discusses PNA-DNA-PNA chimeric molecules and their uses in diagnostics, modulating protein in organisms, and treatment of conditions susceptible to therapeutics.

Methods of characterizing the antisense binding properties of PNAs are discussed in Rose (*Anal Chem.* 1993 Dec 15;65(24):3545-9) and Jensen *et al.*

(Biochemistry. 1997 Apr 22;36(16):5072-7). Rose uses capillary gel electrophoresis to determine binding of PNAs to their complementary oligonucleotide, measuring the relative binding kinetics and stoichiometry. Similar types of measurements were made by Jensen *et al.* using BIAcore™ technology.

- 5 Other applications of PNAs that have been described and will be apparent to the skilled artisan include use in DNA strand invasion, antisense inhibition, mutational analysis, enhancers of transcription, nucleic acid purification, isolation of transcriptionally active genes, blocking of transcription factor binding, genome cleavage, biosensors, *in situ* hybridization, and the like.

10 Polynucleotide Identification, Characterization and Expression

Polynucleotides compositions of the present invention may be identified, prepared and/or manipulated using any of a variety of well established techniques (see generally, Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratories, Cold Spring Harbor, NY, 1989, and other like references). For
15 example, a polynucleotide may be identified, as described in more detail below, by screening a microarray of cDNAs for tumor-associated expression (*i.e.*, expression that is at least two fold greater in a tumor than in normal tissue, as determined using a representative assay provided herein). Such screens may be performed, for example, using the microarray technology of Affymetrix, Inc. (Santa Clara, CA) according to the
20 manufacturer's instructions (and essentially as described by Schena et al., *Proc. Natl. Acad. Sci. USA* 93:10614-10619, 1996 and Heller et al., *Proc. Natl. Acad. Sci. USA* 94:2150-2155, 1997). Alternatively, polynucleotides may be amplified from cDNA prepared from cells expressing the proteins described herein, such as tumor cells.

- Many template dependent processes are available to amplify a target
25 sequences of interest present in a sample. One of the best known amplification methods is the polymerase chain reaction (PCR™) which is described in detail in U.S. Patent Nos. 4,683,195, 4,683,202 and 4,800,159, each of which is incorporated herein by reference in its entirety. Briefly, in PCR™, two primer sequences are prepared which are complementary to regions on opposite complementary strands of the target
30 sequence. An excess of deoxynucleoside triphosphates is added to a reaction mixture

along with a DNA polymerase (*e.g.*, *Taq* polymerase). If the target sequence is present in a sample, the primers will bind to the target and the polymerase will cause the primers to be extended along the target sequence by adding on nucleotides. By raising and lowering the temperature of the reaction mixture, the extended primers will dissociate from the target to form reaction products, excess primers will bind to the target and to the reaction product and the process is repeated. Preferably reverse transcription and PCRTM amplification procedure may be performed in order to quantify the amount of mRNA amplified. Polymerase chain reaction methodologies are well known in the art.

Any of a number of other template dependent processes, many of which are variations of the PCRTM amplification technique, are readily known and available in the art. Illustratively, some such methods include the ligase chain reaction (referred to as LCR), described, for example, in Eur. Pat. Appl. Publ. No. 320,308 and U.S. Patent No. 4,883,750; Qbeta Replicase, described in PCT Intl. Pat. Appl. Publ. No. PCT/US87/00880; Strand Displacement Amplification (SDA) and Repair Chain Reaction (RCR). Still other amplification methods are described in Great Britain Pat. Appl. No. 2 202 328, and in PCT Intl. Pat. Appl. Publ. No. PCT/US89/01025. Other nucleic acid amplification procedures include transcription-based amplification systems (TAS) (PCT Intl. Pat. Appl. Publ. No. WO 88/10315), including nucleic acid sequence based amplification (NASBA) and 3SR. Eur. Pat. Appl. Publ. No. 329,822 describes a nucleic acid amplification process involving cyclically synthesizing single-stranded RNA ("ssRNA"), ssDNA, and double-stranded DNA (dsDNA). PCT Intl. Pat. Appl. Publ. No. WO 89/06700 describes a nucleic acid sequence amplification scheme based on the hybridization of a promoter/primer sequence to a target single-stranded DNA ("ssDNA") followed by transcription of many RNA copies of the sequence. Other amplification methods such as "RACE" (Frohman, 1990), and "one-sided PCR" (Ohara, 1989) are also well-known to those of skill in the art.

An amplified portion of a polynucleotide of the present invention may be used to isolate a full length gene from a suitable library (*e.g.*, a tumor cDNA library) using well known techniques. Within such techniques, a library (cDNA or genomic) is screened using one or more polynucleotide probes or primers suitable for amplification.

Preferably, a library is size-selected to include larger molecules. Random primed libraries may also be preferred for identifying 5' and upstream regions of genes. Genomic libraries are preferred for obtaining introns and extending 5' sequences.

For hybridization techniques, a partial sequence may be labeled (*e.g.*, by
5 nick-translation or end-labeling with ³²P) using well known techniques. A bacterial or bacteriophage library is then generally screened by hybridizing filters containing denatured bacterial colonies (or lawns containing phage plaques) with the labeled probe (*see* Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratories, Cold Spring Harbor, NY, 1989). Hybridizing colonies or plaques
10 are selected and expanded, and the DNA is isolated for further analysis. cDNA clones may be analyzed to determine the amount of additional sequence by, for example, PCR using a primer from the partial sequence and a primer from the vector. Restriction maps and partial sequences may be generated to identify one or more overlapping clones. The complete sequence may then be determined using standard techniques, which may
15 involve generating a series of deletion clones. The resulting overlapping sequences can then be assembled into a single contiguous sequence. A full length cDNA molecule can be generated by ligating suitable fragments, using well known techniques.

Alternatively, amplification techniques, such as those described above, can be useful for obtaining a full length coding sequence from a partial cDNA sequence.
20 One such amplification technique is inverse PCR (*see* Triglia et al., *Nucl. Acids Res.* 16:8186, 1988), which uses restriction enzymes to generate a fragment in the known region of the gene. The fragment is then circularized by intramolecular ligation and used as a template for PCR with divergent primers derived from the known region. Within an alternative approach, sequences adjacent to a partial sequence may be
25 retrieved by amplification with a primer to a linker sequence and a primer specific to a known region. The amplified sequences are typically subjected to a second round of amplification with the same linker primer and a second primer specific to the known region. A variation on this procedure, which employs two primers that initiate extension in opposite directions from the known sequence, is described in WO
30 96/38591. Another such technique is known as "rapid amplification of cDNA ends" or RACE. This technique involves the use of an internal primer and an external primer,

which hybridizes to a polyA region or vector sequence, to identify sequences that are 5' and 3' of a known sequence. Additional techniques include capture PCR (Lagerstrom et al., *PCR Methods Applic. 1*:111-19, 1991) and walking PCR (Parker et al., *Nucl. Acids. Res. 19*:3055-60, 1991). Other methods employing amplification may also be employed to obtain a full length cDNA sequence.

In certain instances, it is possible to obtain a full length cDNA sequence by analysis of sequences provided in an expressed sequence tag (EST) database, such as that available from GenBank. Searches for overlapping ESTs may generally be performed using well known programs (e.g., NCBI BLAST searches), and such ESTs may be used to generate a contiguous full length sequence. Full length DNA sequences may also be obtained by analysis of genomic fragments.

In other embodiments of the invention, polynucleotide sequences or fragments thereof which encode polypeptides of the invention, or fusion proteins or functional equivalents thereof, may be used in recombinant DNA molecules to direct expression of a polypeptide in appropriate host cells. Due to the inherent degeneracy of the genetic code, other DNA sequences that encode substantially the same or a functionally equivalent amino acid sequence may be produced and these sequences may be used to clone and express a given polypeptide.

As will be understood by those of skill in the art, it may be advantageous in some instances to produce polypeptide-encoding nucleotide sequences possessing non-naturally occurring codons. For example, codons preferred by a particular prokaryotic or eukaryotic host can be selected to increase the rate of protein expression or to produce a recombinant RNA transcript having desirable properties, such as a half-life which is longer than that of a transcript generated from the naturally occurring sequence.

Moreover, the polynucleotide sequences of the present invention can be engineered using methods generally known in the art in order to alter polypeptide encoding sequences for a variety of reasons, including but not limited to, alterations which modify the cloning, processing, and/or expression of the gene product. For example, DNA shuffling by random fragmentation and PCR reassembly of gene fragments and synthetic oligonucleotides may be used to engineer the nucleotide

sequences. In addition, site-directed mutagenesis may be used to insert new restriction sites, alter glycosylation patterns, change codon preference, produce splice variants, or introduce mutations, and so forth.

In another embodiment of the invention, natural, modified, or
5 recombinant nucleic acid sequences may be ligated to a heterologous sequence to encode a fusion protein. For example, to screen peptide libraries for inhibitors of polypeptide activity, it may be useful to encode a chimeric protein that can be recognized by a commercially available antibody. A fusion protein may also be engineered to contain a cleavage site located between the polypeptide-encoding
10 sequence and the heterologous protein sequence, so that the polypeptide may be cleaved and purified away from the heterologous moiety.

Sequences encoding a desired polypeptide may be synthesized, in whole or in part, using chemical methods well known in the art (see Caruthers, M. H. et al. (1980) *Nucl. Acids Res. Symp. Ser.* 215-223, Horn, T. et al. (1980) *Nucl. Acids Res. Symp. Ser.* 225-232). Alternatively, the protein itself may be produced using chemical
15 methods to synthesize the amino acid sequence of a polypeptide, or a portion thereof. For example, peptide synthesis can be performed using various solid-phase techniques (Roberge, J. Y. et al. (1995) *Science* 269:202-204) and automated synthesis may be achieved, for example, using the ABI 431A Peptide Synthesizer (Perkin Elmer, Palo
20 Alto, CA).

A newly synthesized peptide may be substantially purified by preparative high performance liquid chromatography (e.g., Creighton, T. (1983) *Proteins, Structures and Molecular Principles*, WH Freeman and Co., New York, N.Y.) or other comparable techniques available in the art. The composition of the synthetic peptides may be
25 confirmed by amino acid analysis or sequencing (e.g., the Edman degradation procedure). Additionally, the amino acid sequence of a polypeptide, or any part thereof, may be altered during direct synthesis and/or combined using chemical methods with sequences from other proteins, or any part thereof, to produce a variant polypeptide.

In order to express a desired polypeptide, the nucleotide sequences
30 encoding the polypeptide, or functional equivalents, may be inserted into appropriate expression vector, i.e., a vector which contains the necessary elements for the

transcription and translation of the inserted coding sequence. Methods which are well known to those skilled in the art may be used to construct expression vectors containing sequences encoding a polypeptide of interest and appropriate transcriptional and translational control elements. These methods include *in vitro* recombinant DNA techniques, synthetic techniques, and *in vivo* genetic recombination. Such techniques are described, for example, in Sambrook, J. et al. (1989) *Molecular Cloning, A Laboratory Manual*, Cold Spring Harbor Press, Plainview, N.Y., and Ausubel, F. M. et al. (1989) *Current Protocols in Molecular Biology*, John Wiley & Sons, New York, N.Y.

- 10 A variety of expression vector/host systems may be utilized to contain and express polynucleotide sequences. These include, but are not limited to, microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors; insect cell systems infected with virus expression vectors (*e.g.*, baculovirus); plant cell
15 systems transformed with virus expression vectors (*e.g.*, cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or with bacterial expression vectors (*e.g.*, Ti or pBR322 plasmids); or animal cell systems.

- The "control elements" or "regulatory sequences" present in an expression vector are those non-translated regions of the vector--enhancers, promoters,
20 5' and 3' untranslated regions--which interact with host cellular proteins to carry out transcription and translation. Such elements may vary in their strength and specificity. Depending on the vector system and host utilized, any number of suitable transcription and translation elements, including constitutive and inducible promoters, may be used. For example, when cloning in bacterial systems, inducible promoters such as the hybrid
25 lacZ promoter of the pBLUESCRIPT phagemid (Stratagene, La Jolla, Calif.) or pSPORT1 plasmid (Gibco BRL, Gaithersburg, MD) and the like may be used. In mammalian cell systems, promoters from mammalian genes or from mammalian viruses are generally preferred. If it is necessary to generate a cell line that contains multiple copies of the sequence encoding a polypeptide, vectors based on SV40 or EBV may be
30 advantageously used with an appropriate selectable marker.

In bacterial systems, any of a number of expression vectors may be selected depending upon the use intended for the expressed polypeptide. For example, when large quantities are needed, for example for the induction of antibodies, vectors which direct high level expression of fusion proteins that are readily purified may be used. Such vectors include, but are not limited to, the multifunctional *E. coli* cloning and expression vectors such as pBLUESCRIPT (Stratagene), in which the sequence encoding the polypeptide of interest may be ligated into the vector in frame with sequences for the amino-terminal Met and the subsequent 7 residues of β -galactosidase so that a hybrid protein is produced; pIN vectors (Van Heeke, G. and S. M. Schuster (1989) *J. Biol. Chem.* 264:5503-5509); and the like. pGEX Vectors (Promega, Madison, Wis.) may also be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are soluble and can easily be purified from lysed cells by adsorption to glutathione-agarose beads followed by elution in the presence of free glutathione. Proteins made in such systems may be designed to include heparin, thrombin, or factor XA protease cleavage sites so that the cloned polypeptide of interest can be released from the GST moiety at will.

In the yeast, *Saccharomyces cerevisiae*, a number of vectors containing constitutive or inducible promoters such as alpha factor, alcohol oxidase, and PGH may be used. For reviews, see Ausubel et al. (supra) and Grant et al. (1987) *Methods Enzymol.* 153:516-544.

In cases where plant expression vectors are used, the expression of sequences encoding polypeptides may be driven by any of a number of promoters. For example, viral promoters such as the 35S and 19S promoters of CaMV may be used alone or in combination with the omega leader sequence from TMV (Takamatsu, N. (1987) *EMBO J.* 6:307-311. Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock promoters may be used (Coruzzi, G. et al. (1984) *EMBO J.* 3:1671-1680; Broglie, R. et al. (1984) *Science* 224:838-843; and Winter, J. et al. (1991) *Results Probl. Cell Differ.* 17:85-105). These constructs can be introduced into plant cells by direct DNA transformation or pathogen-mediated transfection. Such techniques are described in a number of generally available reviews (see, for example, Hobbs, S. or

Murry, L. E. in McGraw Hill Yearbook of Science and Technology (1992) McGraw Hill, New York, N.Y.; pp. 191-196).

An insect system may also be used to express a polypeptide of interest. For example, in one such system, *Autographa californica* nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes in *Spodoptera frugiperda* cells or in *Trichoplusia* larvae. The sequences encoding the polypeptide may be cloned into a non-essential region of the virus, such as the polyhedrin gene, and placed under control of the polyhedrin promoter. Successful insertion of the polypeptide-encoding sequence will render the polyhedrin gene inactive and produce recombinant virus lacking coat protein. The recombinant viruses may then be used to infect, for example, *S. frugiperda* cells or *Trichoplusia* larvae in which the polypeptide of interest may be expressed (Engelhard, E. K. et al. (1994) *Proc. Natl. Acad. Sci.* 91 :3224-3227).

In mammalian host cells, a number of viral-based expression systems are generally available. For example, in cases where an adenovirus is used as an expression vector, sequences encoding a polypeptide of interest may be ligated into an adenovirus transcription/translation complex consisting of the late promoter and tripartite leader sequence. Insertion in a non-essential E1 or E3 region of the viral genome may be used to obtain a viable virus which is capable of expressing the polypeptide in infected host cells (Logan, J. and Shenk, T. (1984) *Proc. Natl. Acad. Sci.* 81:3655-3659). In addition, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, may be used to increase expression in mammalian host cells.

Specific initiation signals may also be used to achieve more efficient translation of sequences encoding a polypeptide of interest. Such signals include the ATG initiation codon and adjacent sequences. In cases where sequences encoding the polypeptide, its initiation codon, and upstream sequences are inserted into the appropriate expression vector, no additional transcriptional or translational control signals may be needed. However, in cases where only coding sequence, or a portion thereof, is inserted, exogenous translational control signals including the ATG initiation codon should be provided. Furthermore, the initiation codon should be in the correct reading frame to ensure translation of the entire insert. Exogenous translational elements and initiation codons may be of various origins, both natural and synthetic.

The efficiency of expression may be enhanced by the inclusion of enhancers which are appropriate for the particular cell system which is used, such as those described in the literature (Scharf, D. et al. (1994) *Results Probl. Cell Differ.* 20:125-162).

In addition, a host cell strain may be chosen for its ability to modulate the expression of the inserted sequences or to process the expressed protein in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing which cleaves a "prepro" form of the protein may also be used to facilitate correct insertion, folding and/or function. Different host cells such as CHO, COS, HeLa, MDCK, HEK293, and WI38, which have specific cellular machinery and characteristic mechanisms for such post-translational activities, may be chosen to ensure the correct modification and processing of the foreign protein.

For long-term, high-yield production of recombinant proteins, stable expression is generally preferred. For example, cell lines which stably express a polynucleotide of interest may be transformed using expression vectors which may contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector. Following the introduction of the vector, cells may be allowed to grow for 1-2 days in an enriched media before they are switched to selective media. The purpose of the selectable marker is to confer resistance to selection, and its presence allows growth and recovery of cells which successfully express the introduced sequences. Resistant clones of stably transformed cells may be proliferated using tissue culture techniques appropriate to the cell type.

Any number of selection systems may be used to recover transformed cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase (Wigler, M. et al. (1977) *Cell* 11:223-32) and adenine phosphoribosyltransferase (Lowy, I. et al. (1990) *Cell* 22:817-23) genes which can be employed in tk.sup.- or aprt.sup.- cells, respectively. Also, antimetabolite, antibiotic or herbicide resistance can be used as the basis for selection; for example, dhfr which confers resistance to methotrexate (Wigler, M. et al. (1980) *Proc. Natl. Acad. Sci.* 77:3567-70); npt, which confers resistance to the aminoglycosides, neomycin and G-418 (Colbere-Garapin, F. et al (1981) *J. Mol. Biol.* 150:1-14); and als or pat, which confer resistance to

chlorsulfuron and phosphinotricin acetyltransferase, respectively (Murry, *supra*). Additional selectable genes have been described, for example, *trpB*, which allows cells to utilize indole in place of tryptophan, or *hisD*, which allows cells to utilize histinol in place of histidine (Hartman, S. C. and R. C. Mulligan (1988) *Proc. Natl. Acad. Sci.* 85:8047-51). The use of visible markers has gained popularity with such markers as anthocyanins, beta-glucuronidase and its substrate GUS, and luciferase and its substrate luciferin, being widely used not only to identify transformants, but also to quantify the amount of transient or stable protein expression attributable to a specific vector system (Rhodes, C. A. et al. (1995) *Methods Mol. Biol.* 55:121-131).

Although the presence/absence of marker gene expression suggests that the gene of interest is also present, its presence and expression may need to be confirmed. For example, if the sequence encoding a polypeptide is inserted within a marker gene sequence, recombinant cells containing sequences can be identified by the absence of marker gene function. Alternatively, a marker gene can be placed in tandem with a polypeptide-encoding sequence under the control of a single promoter. Expression of the marker gene in response to induction or selection usually indicates expression of the tandem gene as well.

Alternatively, host cells that contain and express a desired polynucleotide sequence may be identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations and protein bioassay or immunoassay techniques which include, for example, membrane, solution, or chip based technologies for the detection and/or quantification of nucleic acid or protein.

A variety of protocols for detecting and measuring the expression of polynucleotide-encoded products, using either polyclonal or monoclonal antibodies specific for the product are known in the art. Examples include enzyme-linked immunosorbent assay (ELISA), radioimmunoassay (RIA), and fluorescence activated cell sorting (FACS). A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering epitopes on a given polypeptide may be preferred for some applications, but a competitive binding assay may also be employed. These and other assays are described, among other places, in Hampton, R. et al. (1990);

Serological Methods, a Laboratory Manual, APS Press, St Paul, Minn.) and Maddox, D. E. et al. (1983; *J. Exp. Med.* 158:1211-1216).

A wide variety of labels and conjugation techniques are known by those skilled in the art and may be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to polynucleotides include oligolabeling, nick translation, end-labeling or PCR amplification using a labeled nucleotide. Alternatively, the sequences, or any portions thereof may be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes in vitro by addition of an appropriate RNA polymerase such as T7, T3, or SP6 and labeled nucleotides. These procedures may be conducted using a variety of commercially available kits. Suitable reporter molecules or labels, which may be used include radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

Host cells transformed with a polynucleotide sequence of interest may be cultured under conditions suitable for the expression and recovery of the protein from cell culture. The protein produced by a recombinant cell may be secreted or contained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides of the invention may be designed to contain signal sequences which direct secretion of the encoded polypeptide through a prokaryotic or eukaryotic cell membrane. Other recombinant constructions may be used to join sequences encoding a polypeptide of interest to nucleotide sequence encoding a polypeptide domain which will facilitate purification of soluble proteins. Such purification facilitating domains include, but are not limited to, metal chelating peptides such as histidine-tryptophan modules that allow purification on immobilized metals, protein A domains that allow purification on immobilized immunoglobulin, and the domain utilized in the FLAGS extension/affinity purification system (Immunex Corp., Seattle, Wash.). The inclusion of cleavable linker sequences such as those specific for Factor XA or enterokinase (Invitrogen, San Diego, Calif.) between the purification domain and the encoded polypeptide may be used to facilitate purification. One such expression vector provides for expression of a fusion

protein containing a polypeptide of interest and a nucleic acid encoding 6 histidine residues preceding a thioredoxin or an enterokinase cleavage site. The histidine residues facilitate purification on IMIAC (immobilized metal ion affinity chromatography) as described in Porath, J. et al. (1992, *Prot. Exp. Purif.* 3:263-281) while the enterokinase cleavage site provides a means for purifying the desired polypeptide from the fusion protein. A discussion of vectors which contain fusion proteins is provided in Kroll, D. J. et al. (1993; *DNA Cell Biol.* 12:441-453).

In addition to recombinant production methods, polypeptides of the invention, and fragments thereof, may be produced by direct peptide synthesis using solid-phase techniques (Merrifield J. (1963) *J. Am. Chem. Soc.* 85:2149-2154). Protein synthesis may be performed using manual techniques or by automation. Automated synthesis may be achieved, for example, using Applied Biosystems 431A Peptide Synthesizer (Perkin Elmer). Alternatively, various fragments may be chemically synthesized separately and combined using chemical methods to produce the full length molecule.

Antibody Compositions, Fragments Thereof and Other Binding Agents

According to another aspect, the present invention further provides binding agents, such as antibodies and antigen-binding fragments thereof, that exhibit immunological binding to a tumor polypeptide disclosed herein, or to a portion, variant or derivative thereof. An antibody, or antigen-binding fragment thereof, is said to "specifically bind," "immunologically bind," and/or is "immunologically reactive" to a polypeptide of the invention if it reacts at a detectable level (within, for example, an ELISA assay) with the polypeptide, and does not react detectably with unrelated polypeptides under similar conditions.

Immunological binding, as used in this context, generally refers to the non-covalent interactions of the type which occur between an immunoglobulin molecule and an antigen for which the immunoglobulin is specific. The strength, or affinity of immunological binding interactions can be expressed in terms of the dissociation constant (K_d) of the interaction, wherein a smaller K_d represents a greater affinity. Immunological binding properties of selected polypeptides can be quantified

using methods well known in the art. One such method entails measuring the rates of antigen-binding site/antigen complex formation and dissociation, wherein those rates depend on the concentrations of the complex partners, the affinity of the interaction, and on geometric parameters that equally influence the rate in both directions. Thus, both the "on rate constant" (K_{on}) and the "off rate constant" (K_{off}) can be determined by calculation of the concentrations and the actual rates of association and dissociation. The ratio of K_{off}/K_{on} enables cancellation of all parameters not related to affinity, and is thus equal to the dissociation constant K_d . See, generally, Davies et al. (1990) Annual Rev. Biochem. 59:439-473.

10 An "antigen-binding site," or "binding portion" of an antibody refers to the part of the immunoglobulin molecule that participates in antigen binding. The antigen binding site is formed by amino acid residues of the N-terminal variable ("V") regions of the heavy ("H") and light ("L") chains. Three highly divergent stretches within the V regions of the heavy and light chains are referred to as "hypervariable regions" which are interposed between more conserved flanking stretches known as "framework regions," or "FRs". Thus the term "FR" refers to amino acid sequences which are naturally found between and adjacent to hypervariable regions in immunoglobulins. In an antibody molecule, the three hypervariable regions of a light chain and the three hypervariable regions of a heavy chain are disposed relative to each other in three dimensional space to form an antigen-binding surface. The antigen-binding surface is complementary to the three-dimensional surface of a bound antigen, and the three hypervariable regions of each of the heavy and light chains are referred to as "complementarity-determining regions," or "CDRs."

25 Binding agents may be further capable of differentiating between patients with and without a cancer, such as lung cancer, using the representative assays provided herein. For example, antibodies or other binding agents that bind to a tumor protein will preferably generate a signal indicating the presence of a cancer in at least about 20% of patients with the disease, more preferably at least about 30% of patients. Alternatively, or in addition, the antibody will generate a negative signal indicating the absence of the disease in at least about 90% of individuals without the cancer. To determine whether a binding agent satisfies this requirement, biological samples (e.g.,

blood, sera, sputum, urine and/or tumor biopsies) from patients with and without a cancer (as determined using standard clinical tests) may be assayed as described herein for the presence of polypeptides that bind to the binding agent. Preferably, a statistically significant number of samples with and without the disease will be assayed. Each

5 binding agent should satisfy the above criteria; however, those of ordinary skill in the art will recognize that binding agents may be used in combination to improve sensitivity.

Any agent that satisfies the above requirements may be a binding agent. For example, a binding agent may be a ribosome, with or without a peptide component,

10 an RNA molecule or a polypeptide. In a preferred embodiment, a binding agent is an antibody or an antigen-binding fragment thereof. Antibodies may be prepared by any of a variety of techniques known to those of ordinary skill in the art. See, e.g., Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, 1988. In general, antibodies can be produced by cell culture techniques, including the generation

15 of monoclonal antibodies as described herein, or via transfection of antibody genes into suitable bacterial or mammalian cell hosts, in order to allow for the production of recombinant antibodies. In one technique, an immunogen comprising the polypeptide is initially injected into any of a wide variety of mammals (e.g., mice, rats, rabbits, sheep or goats). In this step, the polypeptides of this invention may serve as the immunogen

20 without modification. Alternatively, particularly for relatively short polypeptides, a superior immune response may be elicited if the polypeptide is joined to a carrier protein, such as bovine serum albumin or keyhole limpet hemocyanin. The immunogen is injected into the animal host, preferably according to a predetermined schedule incorporating one or more booster immunizations, and the animals are bled periodically.

25 Polyclonal antibodies specific for the polypeptide may then be purified from such antisera by, for example, affinity chromatography using the polypeptide coupled to a suitable solid support.

Monoclonal antibodies specific for an antigenic polypeptide of interest may be prepared, for example, using the technique of Kohler and Milstein, *Eur. J.*

30 *Immunol.* 6:511-519, 1976, and improvements thereto. Briefly, these methods involve the preparation of immortal cell lines capable of producing antibodies having the

desired specificity (*i.e.*, reactivity with the polypeptide of interest). Such cell lines may be produced, for example, from spleen cells obtained from an animal immunized as described above. The spleen cells are then immortalized by, for example, fusion with a myeloma cell fusion partner, preferably one that is syngeneic with the immunized animal. A variety of fusion techniques may be employed. For example, the spleen cells and myeloma cells may be combined with a nonionic detergent for a few minutes and then plated at low density on a selective medium that supports the growth of hybrid cells, but not myeloma cells. A preferred selection technique uses HAT (hypoxanthine, aminopterin, thymidine) selection. After a sufficient time, usually about 1 to 2 weeks, colonies of hybrids are observed. Single colonies are selected and their culture supernatants tested for binding activity against the polypeptide. Hybridomas having high reactivity and specificity are preferred.

Monoclonal antibodies may be isolated from the supernatants of growing hybridoma colonies. In addition, various techniques may be employed to enhance the yield, such as injection of the hybridoma cell line into the peritoneal cavity of a suitable vertebrate host, such as a mouse. Monoclonal antibodies may then be harvested from the ascites fluid or the blood. Contaminants may be removed from the antibodies by conventional techniques, such as chromatography, gel filtration, precipitation, and extraction. The polypeptides of this invention may be used in the purification process in, for example, an affinity chromatography step.

A number of therapeutically useful molecules are known in the art which comprise antigen-binding sites that are capable of exhibiting immunological binding properties of an antibody molecule. The proteolytic enzyme papain preferentially cleaves IgG molecules to yield several fragments, two of which (the "F(ab)" fragments) each comprise a covalent heterodimer that includes an intact antigen-binding site. The enzyme pepsin is able to cleave IgG molecules to provide several fragments, including the "F(ab')₂" fragment which comprises both antigen-binding sites. An "Fv" fragment can be produced by preferential proteolytic cleavage of an IgM, and on rare occasions IgG or IgA immunoglobulin molecule. Fv fragments are, however, more commonly derived using recombinant techniques known in the art. The Fv fragment includes a non-covalent V_H::V_L heterodimer including an antigen-binding site which retains much

of the antigen recognition and binding capabilities of the native antibody molecule. Inbar et al. (1972) Proc. Nat. Acad. Sci. USA 69:2659-2662; Hochman et al. (1976) Biochem 15:2706-2710; and Ehrlich et al. (1980) Biochem 19:4091-4096.

A single chain Fv ("sFv") polypeptide is a covalently linked $V_H::V_L$ heterodimer which is expressed from a gene fusion including V_H - and V_L -encoding genes linked by a peptide-encoding linker. Huston et al. (1988) Proc. Nat. Acad. Sci. USA 85(16):5879-5883. A number of methods have been described to discern chemical structures for converting the naturally aggregated--but chemically separated--light and heavy polypeptide chains from an antibody V region into an sFv molecule which will fold into a three dimensional structure substantially similar to the structure of an antigen-binding site. See, e.g., U.S. Pat. Nos. 5,091,513 and 5,132,405, to Huston et al.; and U.S. Pat. No. 4,946,778, to Ladner et al.

Each of the above-described molecules includes a heavy chain and a light chain CDR set, respectively interposed between a heavy chain and a light chain FR set which provide support to the CDRs and define the spatial relationship of the CDRs relative to each other. As used herein, the term "CDR set" refers to the three hypervariable regions of a heavy or light chain V region. Proceeding from the N-terminus of a heavy or light chain, these regions are denoted as "CDR1," "CDR2," and "CDR3" respectively. An antigen-binding site, therefore, includes six CDRs, comprising the CDR set from each of a heavy and a light chain V region. A polypeptide comprising a single CDR, (e.g., a CDR1, CDR2 or CDR3) is referred to herein as a "molecular recognition unit." Crystallographic analysis of a number of antigen-antibody complexes has demonstrated that the amino acid residues of CDRs form extensive contact with bound antigen, wherein the most extensive antigen contact is with the heavy chain CDR3. Thus, the molecular recognition units are primarily responsible for the specificity of an antigen-binding site.

As used herein, the term "FR set" refers to the four flanking amino acid sequences which frame the CDRs of a CDR set of a heavy or light chain V region. Some FR residues may contact bound antigen; however, FRs are primarily responsible for folding the V region into the antigen-binding site, particularly the FR residues directly adjacent to the CDRs. Within FRs, certain amino residues and certain structural

features are very highly conserved. In this regard, all V region sequences contain an internal disulfide loop of around 90 amino acid residues. When the V regions fold into a binding-site, the CDRs are displayed as projecting loop motifs which form an antigen-binding surface. It is generally recognized that there are conserved structural regions of FRs which influence the folded shape of the CDR loops into certain "canonical" structures--regardless of the precise CDR amino acid sequence. Further, certain FR residues are known to participate in non-covalent interdomain contacts which stabilize the interaction of the antibody heavy and light chains.

A number of "humanized" antibody molecules comprising an antigen-binding site derived from a non-human immunoglobulin have been described, including chimeric antibodies having rodent V regions and their associated CDRs fused to human constant domains (Winter et al. (1991) *Nature* 349:293-299; Lobuglio et al. (1989) *Proc. Nat. Acad. Sci. USA* 86:4220-4224; Shaw et al. (1987) *J Immunol.* 138:4534-4538; and Brown et al. (1987) *Cancer Res.* 47:3577-3583), rodent CDRs grafted into a human supporting FR prior to fusion with an appropriate human antibody constant domain (Riechmann et al. (1988) *Nature* 332:323-327; Verhoeyen et al. (1988) *Science* 239:1534-1536; and Jones et al. (1986) *Nature* 321:522-525), and rodent CDRs supported by recombinantly veneered rodent FRs (European Patent Publication No. 519,596, published Dec. 23, 1992). These "humanized" molecules are designed to minimize unwanted immunological response toward rodent antihuman antibody molecules which limits the duration and effectiveness of therapeutic applications of those moieties in human recipients.

As used herein, the terms "veneered FRs" and "recombinantly veneered FRs" refer to the selective replacement of FR residues from, e.g., a rodent heavy or light chain V region, with human FR residues in order to provide a xenogeneic molecule comprising an antigen-binding site which retains substantially all of the native FR polypeptide folding structure. Veneering techniques are based on the understanding that the ligand binding characteristics of an antigen-binding site are determined primarily by the structure and relative disposition of the heavy and light chain CDR sets within the antigen-binding surface. Davies et al. (1990) *Ann. Rev. Biochem.* 59:439-473. Thus, antigen binding specificity can be preserved in a humanized antibody only wherein the

CDR structures, their interaction with each other, and their interaction with the rest of the V region domains are carefully maintained. By using veneering techniques, exterior (e.g., solvent-accessible) FR residues which are readily encountered by the immune system are selectively replaced with human residues to provide a hybrid molecule that comprises either a weakly immunogenic, or substantially non-immunogenic veneered surface.

The process of veneering makes use of the available sequence data for human antibody variable domains compiled by Kabat et al., in Sequences of Proteins of Immunological Interest, 4th ed., (U.S. Dept. of Health and Human Services, U.S. Government Printing Office, 1987), updates to the Kabat database, and other accessible U.S. and foreign databases (both nucleic acid and protein). Solvent accessibilities of V region amino acids can be deduced from the known three-dimensional structure for human and murine antibody fragments. There are two general steps in veneering a murine antigen-binding site. Initially, the FRs of the variable domains of an antibody molecule of interest are compared with corresponding FR sequences of human variable domains obtained from the above-identified sources. The most homologous human V regions are then compared residue by residue to corresponding murine amino acids. The residues in the murine FR which differ from the human counterpart are replaced by the residues present in the human moiety using recombinant techniques well known in the art. Residue switching is only carried out with moieties which are at least partially exposed (solvent accessible), and care is exercised in the replacement of amino acid residues which may have a significant effect on the tertiary structure of V region domains, such as proline, glycine and charged amino acids.

In this manner, the resultant "veneered" murine antigen-binding sites are thus designed to retain the murine CDR residues, the residues substantially adjacent to the CDRs, the residues identified as buried or mostly buried (solvent inaccessible), the residues believed to participate in non-covalent (e.g., electrostatic and hydrophobic) contacts between heavy and light chain domains, and the residues from conserved structural regions of the FRs which are believed to influence the "canonical" tertiary structures of the CDR loops. These design criteria are then used to prepare recombinant nucleotide sequences which combine the CDRs of both the heavy and light chain of a

murine antigen-binding site into human-appearing FRs that can be used to transfect mammalian cells for the expression of recombinant human antibodies which exhibit the antigen specificity of the murine antibody molecule.

In another embodiment of the invention, monoclonal antibodies of the present invention may be coupled to one or more therapeutic agents. Suitable agents in this regard include radionuclides, differentiation inducers, drugs, toxins, and derivatives thereof. Preferred radionuclides include ^{90}Y , ^{123}I , ^{125}I , ^{131}I , ^{186}Re , ^{188}Re , ^{211}At , and ^{212}Bi . Preferred drugs include methotrexate, and pyrimidine and purine analogs. Preferred differentiation inducers include phorbol esters and butyric acid. Preferred toxins include ricin, abrin, diphtheria toxin, cholera toxin, gelonin, *Pseudomonas* exotoxin, *Shigella* toxin, and pokeweed antiviral protein.

A therapeutic agent may be coupled (*e.g.*, covalently bonded) to a suitable monoclonal antibody either directly or indirectly (*e.g.*, via a linker group). A direct reaction between an agent and an antibody is possible when each possesses a substituent capable of reacting with the other. For example, a nucleophilic group, such as an amino or sulfhydryl group, on one may be capable of reacting with a carbonyl-containing group, such as an anhydride or an acid halide, or with an alkyl group containing a good leaving group (*e.g.*, a halide) on the other.

Alternatively, it may be desirable to couple a therapeutic agent and an antibody via a linker group. A linker group can function as a spacer to distance an antibody from an agent in order to avoid interference with binding capabilities. A linker group can also serve to increase the chemical reactivity of a substituent on an agent or an antibody, and thus increase the coupling efficiency. An increase in chemical reactivity may also facilitate the use of agents, or functional groups on agents, which otherwise would not be possible.

It will be evident to those skilled in the art that a variety of bifunctional or polyfunctional reagents, both homo- and hetero-functional (such as those described in the catalog of the Pierce Chemical Co., Rockford, IL), may be employed as the linker group. Coupling may be effected, for example, through amino groups, carboxyl groups, sulfhydryl groups or oxidized carbohydrate residues. There are numerous references describing such methodology, *e.g.*, U.S. Patent No. 4,671,958, to Rodwell et al.

Where a therapeutic agent is more potent when free from the antibody portion of the immunoconjugates of the present invention, it may be desirable to use a linker group which is cleavable during or upon internalization into a cell. A number of different cleavable linker groups have been described. The mechanisms for the intracellular release of an agent from these linker groups include cleavage by reduction of a disulfide bond (*e.g.*, U.S. Patent No. 4,489,710, to Spittler), by irradiation of a photolabile bond (*e.g.*, U.S. Patent No. 4,625,014, to Senter et al.), by hydrolysis of derivatized amino acid side chains (*e.g.*, U.S. Patent No. 4,638,045, to Kohn et al.), by serum complement-mediated hydrolysis (*e.g.*, U.S. Patent No. 4,671,958, to Rodwell et al.), and acid-catalyzed hydrolysis (*e.g.*, U.S. Patent No. 4,569,789, to Blattler et al.).

It may be desirable to couple more than one agent to an antibody. In one embodiment, multiple molecules of an agent are coupled to one antibody molecule. In another embodiment, more than one type of agent may be coupled to one antibody. Regardless of the particular embodiment, immunoconjugates with more than one agent may be prepared in a variety of ways. For example, more than one agent may be coupled directly to an antibody molecule, or linkers that provide multiple sites for attachment can be used. Alternatively, a carrier can be used.

A carrier may bear the agents in a variety of ways, including covalent bonding either directly or via a linker group. Suitable carriers include proteins such as albumins (*e.g.*, U.S. Patent No. 4,507,234, to Kato et al.), peptides and polysaccharides such as aminodextran (*e.g.*, U.S. Patent No. 4,699,784, to Shih et al.). A carrier may also bear an agent by noncovalent bonding or by encapsulation, such as within a liposome vesicle (*e.g.*, U.S. Patent Nos. 4,429,008 and 4,873,088). Carriers specific for radionuclide agents include radiohalogenated small molecules and chelating compounds. For example, U.S. Patent No. 4,735,792 discloses representative radiohalogenated small molecules and their synthesis. A radionuclide chelate may be formed from chelating compounds that include those containing nitrogen and sulfur atoms as the donor atoms for binding the metal, or metal oxide, radionuclide. For example, U.S. Patent No. 4,673,562, to Davison et al. discloses representative chelating compounds and their synthesis.

T Cell Compositions

The present invention, in another aspect, provides T cells specific for a tumor polypeptide disclosed herein, or for a variant or derivative thereof. Such cells may generally be prepared *in vitro* or *ex vivo*, using standard procedures. For example,

5 T cells may be isolated from bone marrow, peripheral blood, or a fraction of bone marrow or peripheral blood of a patient, using a commercially available cell separation system, such as the Isolex™ System, available from Nexell Therapeutics, Inc. (Irvine, CA; see also U.S. Patent No. 5,240,856; U.S. Patent No. 5,215,926; WO 89/06280; WO 91/16116 and WO 92/07243). Alternatively, T cells may be derived from related or

10 unrelated humans, non-human mammals, cell lines or cultures.

T cells may be stimulated with a polypeptide, polynucleotide encoding a polypeptide and/or an antigen presenting cell (APC) that expresses such a polypeptide. Such stimulation is performed under conditions and for a time sufficient to permit the generation of T cells that are specific for the polypeptide of interest. Preferably, a tumor

15 polypeptide or polynucleotide of the invention is present within a delivery vehicle, such as a microsphere, to facilitate the generation of specific T cells.

T cells are considered to be specific for a polypeptide of the present invention if the T cells specifically proliferate, secrete cytokines or kill target cells coated with the polypeptide or expressing a gene encoding the polypeptide. T cell

20 specificity may be evaluated using any of a variety of standard techniques. For example, within a chromium release assay or proliferation assay, a stimulation index of more than two fold increase in lysis and/or proliferation, compared to negative controls, indicates T cell specificity. Such assays may be performed, for example, as described in Chen et al., *Cancer Res.* 54:1065-1070, 1994. Alternatively, detection of the

25 proliferation of T cells may be accomplished by a variety of known techniques. For example, T cell proliferation can be detected by measuring an increased rate of DNA synthesis (*e.g.*, by pulse-labeling cultures of T cells with tritiated thymidine and measuring the amount of tritiated thymidine incorporated into DNA). Contact with a tumor polypeptide (100 ng/ml - 100 µg/ml, preferably 200 ng/ml - 25 µg/ml) for 3 - 7

30 days will typically result in at least a two fold increase in proliferation of the T cells. Contact as described above for 2-3 hours should result in activation of the T cells, as

measured using standard cytokine assays in which a two fold increase in the level of cytokine release (*e.g.*, TNF or IFN- γ) is indicative of T cell activation (*see* Coligan et al., Current Protocols in Immunology, vol. 1, Wiley Interscience (Greene 1998)). T cells that have been activated in response to a tumor polypeptide, polynucleotide or polypeptide-expressing APC may be CD4⁺ and/or CD8⁺. Tumor polypeptide-specific T cells may be expanded using standard techniques. Within preferred embodiments, the T cells are derived from a patient, a related donor or an unrelated donor, and are administered to the patient following stimulation and expansion.

For therapeutic purposes, CD4⁺ or CD8⁺ T cells that proliferate in response to a tumor polypeptide, polynucleotide or APC can be expanded in number either *in vitro* or *in vivo*. Proliferation of such T cells *in vitro* may be accomplished in a variety of ways. For example, the T cells can be re-exposed to a tumor polypeptide, or a short peptide corresponding to an immunogenic portion of such a polypeptide, with or without the addition of T cell growth factors, such as interleukin-2, and/or stimulator cells that synthesize a tumor polypeptide. Alternatively, one or more T cells that proliferate in the presence of the tumor polypeptide can be expanded in number by cloning. Methods for cloning cells are well known in the art, and include limiting dilution.

T Cell Receptor Compositions

The T cell receptor (TCR) consists of 2 different, highly variable polypeptide chains, termed the T-cell receptor α and β chains, that are linked by a disulfide bond (Janeway, Travers, Walport. *Immunobiology*. Fourth Ed., 148-159. Elsevier Science Ltd/Garland Publishing, 1999). The α/β heterodimer complexes with the invariant CD3 chains at the cell membrane. This complex recognizes specific antigenic peptides bound to MHC molecules. The enormous diversity of TCR specificities is generated much like immunoglobulin diversity, through somatic gene rearrangement. The β chain genes contain over 50 variable (V), 2 diversity (D), over 10 joining (J) segments, and 2 constant region segments (C). The α chain genes contain over 70 V segments, and over 60 J segments but no D segments, as well as one C segment. During T cell development in the thymus, the D to J gene rearrangement of

the β chain occurs, followed by the V gene segment rearrangement to the DJ. This functional VDJ β exon is transcribed and spliced to join to a C β . For the α chain, a V α gene segment rearranges to a J α gene segment to create the functional exon that is then transcribed and spliced to the C α . Diversity is further increased during the recombination process by the random addition of P and N-nucleotides between the V, D, and J segments of the β chain and between the V and J segments in the α chain (Janeway, Travers, Walport. *Immunobiology*. Fourth Ed., 98 and 150. Elsevier Science Ltd/Garland Publishing. 1999).

The present invention, in another aspect, provides TCRs specific for a polypeptide disclosed herein, or for a variant or derivative thereof. In accordance with the present invention, polynucleotide and amino acid sequences are provided for the V-J or V-D-J junctional regions or parts thereof for the alpha and beta chains of the T-cell receptor which recognize tumor polypeptides described herein. In general, this aspect of the invention relates to T-cell receptors which recognize or bind tumor polypeptides presented in the context of MHC. In a preferred embodiment the tumor antigens recognized by the T-cell receptors comprise a polypeptide of the present invention. For example, cDNA encoding a TCR specific for a _tumor peptide can be isolated from T cells specific for a tumor polypeptide using standard molecular biological and recombinant DNA techniques.

This invention further includes the T-cell receptors or analogs thereof having substantially the same function or activity as the T-cell receptors of this invention which recognize or bind tumor polypeptides. Such receptors include, but are not limited to, a fragment of the receptor, or a substitution, addition or deletion mutant of a T-cell receptor provided herein. This invention also encompasses polypeptides or peptides that are substantially homologous to the T-cell receptors provided herein or that retain substantially the same activity. The term "analog" includes any protein or polypeptide having an amino acid residue sequence substantially identical to the T-cell receptors provided herein in which one or more residues, preferably no more than 5 residues, more preferably no more than 25 residues have been conservatively substituted with a functionally similar residue and which displays the functional aspects of the T-cell receptor as described herein.

The present invention further provides for suitable mammalian host cells, for example, non-specific T cells, that are transfected with a polynucleotide encoding TCRs specific for a polypeptide described herein, thereby rendering the host cell specific for the polypeptide. The α and β chains of the TCR may be contained on
5 separate expression vectors or alternatively, on a single expression vector that also contains an internal ribosome entry site (IRES) for cap-independent translation of the gene downstream of the IRES. Said host cells expressing TCRs specific for the polypeptide may be used, for example, for adoptive immunotherapy of lung cancer as discussed further below.

10 In further aspects of the present invention, cloned TCRs specific for a polypeptide recited herein may be used in a kit for the diagnosis of lung cancer. For example, the nucleic acid sequence or portions thereof, of tumor-specific TCRs can be used as probes or primers for the detection of expression of the rearranged genes encoding the specific TCR in a biological sample. Therefore, the present invention
15 further provides for an assay for detecting messenger RNA or DNA encoding the TCR specific for a polypeptide.

Pharmaceutical Compositions

In additional embodiments, the present invention concerns formulation of one or more of the polynucleotide, polypeptide, T-cell, TCR, and/or antibody
20 compositions disclosed herein in pharmaceutically-acceptable carriers for administration to a cell or an animal, either alone, or in combination with one or more other modalities of therapy.

It will be understood that, if desired, a composition as disclosed herein may be administered in combination with other agents as well, such as, *e.g.*, other
25 proteins or polypeptides or various pharmaceutically-active agents. In fact, there is virtually no limit to other components that may also be included, given that the additional agents do not cause a significant adverse effect upon contact with the target cells or host tissues. The compositions may thus be delivered along with various other agents as required in the particular instance. Such compositions may be purified from

host cells or other biological sources, or alternatively may be chemically synthesized as described herein. Likewise, such compositions may further comprise substituted or derivatized RNA or DNA compositions.

Therefore, in another aspect of the present invention, pharmaceutical
5 compositions are provided comprising one or more of the polynucleotide, polypeptide, antibody, TCR, and/or T-cell compositions described herein in combination with a physiologically acceptable carrier. In certain preferred embodiments, the pharmaceutical compositions of the invention comprise immunogenic polynucleotide and/or polypeptide compositions of the invention for use in prophylactic and therapeutic
10 vaccine applications. Vaccine preparation is generally described in, for example, M.F. Powell and M.J. Newman, eds., "Vaccine Design (the subunit and adjuvant approach)," Plenum Press (NY, 1995). Generally, such compositions will comprise one or more polynucleotide and/or polypeptide compositions of the present invention in combination with one or more immunostimulants.

15 It will be apparent that any of the pharmaceutical compositions described herein can contain pharmaceutically acceptable salts of the polynucleotides and polypeptides of the invention. Such salts can be prepared, for example, from pharmaceutically acceptable non-toxic bases, including organic bases (*e.g.*, salts of primary, secondary and tertiary amines and basic amino acids) and inorganic bases (*e.g.*,
20 sodium, potassium, lithium, ammonium, calcium and magnesium salts).

In another embodiment, illustrative immunogenic compositions, *e.g.*, vaccine compositions, of the present invention comprise DNA encoding one or more of the polypeptides as described above, such that the polypeptide is generated *in situ*. As noted above, the polynucleotide may be administered within any of a variety of delivery
25 systems known to those of ordinary skill in the art. Indeed, numerous gene delivery techniques are well known in the art, such as those described by Rolland, *Crit. Rev. Therap. Drug Carrier Systems* 15:143-198, 1998, and references cited therein. Appropriate polynucleotide expression systems will, of course, contain the necessary regulatory DNA regulatory sequences for expression in a patient (such as a suitable
30 promoter and terminating signal). Alternatively, bacterial delivery systems may involve

the administration of a bacterium (such as *Bacillus-Calmette-Guerrin*) that expresses an immunogenic portion of the polypeptide on its cell surface or secretes such an epitope.

Therefore, in certain embodiments, polynucleotides encoding immunogenic polypeptides described herein are introduced into suitable mammalian host cells for expression using any of a number of known viral-based systems. In one illustrative embodiment, retroviruses provide a convenient and effective platform for gene delivery systems. A selected nucleotide sequence encoding a polypeptide of the present invention can be inserted into a vector and packaged in retroviral particles using techniques known in the art. The recombinant virus can then be isolated and delivered to a subject. A number of illustrative retroviral systems have been described (e.g., U.S. Pat. No. 5,219,740; Miller and Rosman (1989) *BioTechniques* 7:980-990; Miller, A. D. (1990) *Human Gene Therapy* 1:5-14; Scarpa et al. (1991) *Virology* 180:849-852; Burns et al. (1993) *Proc. Natl. Acad. Sci. USA* 90:8033-8037; and Boris-Lawrie and Temin (1993) *Cur. Opin. Genet. Develop.* 3:102-109.

In addition, a number of illustrative adenovirus-based systems have also been described. Unlike retroviruses which integrate into the host genome, adenoviruses persist extrachromosomally thus minimizing the risks associated with insertional mutagenesis (Haj-Ahmad and Graham (1986) *J. Virol.* 57:267-274; Bett et al. (1993) *J. Virol.* 67:5911-5921; Mittereder et al. (1994) *Human Gene Therapy* 5:717-729; Seth et al. (1994) *J. Virol.* 68:933-940; Barr et al. (1994) *Gene Therapy* 1:51-58; Berkner, K. L. (1988) *BioTechniques* 6:616-629; and Rich et al. (1993) *Human Gene Therapy* 4:461-476).

Various adeno-associated virus (AAV) vector systems have also been developed for polynucleotide delivery. AAV vectors can be readily constructed using techniques well known in the art. See, e.g., U.S. Pat. Nos. 5,173,414 and 5,139,941; International Publication Nos. WO 92/01070 and WO 93/03769; Lebkowski et al. (1988) *Molec. Cell. Biol.* 8:3988-3996; Vincent et al. (1990) *Vaccines* 90 (Cold Spring Harbor Laboratory Press); Carter, B. J. (1992) *Current Opinion in Biotechnology* 3:533-539; Muzyczka, N. (1992) *Current Topics in Microbiol. and Immunol.* 158:97-129; Kotin, R. M. (1994) *Human Gene Therapy* 5:793-801; Shelling and Smith (1994) *Gene Therapy* 1:165-169; and Zhou et al. (1994) *J. Exp. Med.* 179:1867-1875.

Additional viral vectors useful for delivering the polynucleotides encoding polypeptides of the present invention by gene transfer include those derived from the pox family of viruses, such as vaccinia virus and avian poxvirus. By way of example, vaccinia virus recombinants expressing the novel molecules can be constructed as follows. The DNA encoding a polypeptide is first inserted into an appropriate vector so that it is adjacent to a vaccinia promoter and flanking vaccinia DNA sequences, such as the sequence encoding thymidine kinase (TK). This vector is then used to transfect cells which are simultaneously infected with vaccinia. Homologous recombination serves to insert the vaccinia promoter plus the gene encoding the polypeptide of interest into the viral genome. The resulting TK.sup.(-) recombinant can be selected by culturing the cells in the presence of 5-bromodeoxyuridine and picking viral plaques resistant thereto.

A vaccinia-based infection/transfection system can be conveniently used to provide for inducible, transient expression or coexpression of one or more polypeptides described herein in host cells of an organism. In this particular system, cells are first infected in vitro with a vaccinia virus recombinant that encodes the bacteriophage T7 RNA polymerase. This polymerase displays exquisite specificity in that it only transcribes templates bearing T7 promoters. Following infection, cells are transfected with the polynucleotide or polynucleotides of interest, driven by a T7 promoter. The polymerase expressed in the cytoplasm from the vaccinia virus recombinant transcribes the transfected DNA into RNA which is then translated into polypeptide by the host translational machinery. The method provides for high level, transient, cytoplasmic production of large quantities of RNA and its translation products. See, *e.g.*, Elroy-Stein and Moss, Proc. Natl. Acad. Sci. USA (1990) 87:6743-6747; Fuerst et al. Proc. Natl. Acad. Sci. USA (1986) 83:8122-8126.

Alternatively, avipoxviruses, such as the fowlpox and canarypox viruses, can also be used to deliver the coding sequences of interest. Recombinant avipox viruses, expressing immunogens from mammalian pathogens, are known to confer protective immunity when administered to non-avian species. The use of an Avipox vector is particularly desirable in human and other mammalian species since members of the Avipox genus can only productively replicate in susceptible avian species and

therefore are not infective in mammalian cells. Methods for producing recombinant Avipoxviruses are known in the art and employ genetic recombination, as described above with respect to the production of vaccinia viruses. See, e.g., WO 91/12882; WO 89/03429; and WO 92/03545.

5 Any of a number of alphavirus vectors can also be used for delivery of polynucleotide compositions of the present invention, such as those vectors described in U.S. Patent Nos. 5,843,723; 6,015,686; 6,008,035 and 6,015,694. Certain vectors based on Venezuelan Equine Encephalitis (VEE) can also be used, illustrative examples of which can be found in U.S. Patent Nos. 5,505,947 and 5,643,576.

10 Moreover, molecular conjugate vectors, such as the adenovirus chimeric vectors described in Michael et al. *J. Biol. Chem.* (1993) 268:6866-6869 and Wagner et al. *Proc. Natl. Acad. Sci. USA* (1992) 89:6099-6103, can also be used for gene delivery under the invention.

Additional illustrative information on these and other known viral-based
15 delivery systems can be found, for example, in Fisher-Hoch et al., *Proc. Natl. Acad. Sci. USA* 86:317-321, 1989; Flexner et al., *Ann. N.Y. Acad. Sci.* 569:86-103, 1989; Flexner et al., *Vaccine* 8:17-21, 1990; U.S. Patent Nos. 4,603,112, 4,769,330, and 5,017,487; WO 89/01973; U.S. Patent No. 4,777,127; GB 2,200,651; EP 0,345,242; WO 91/02805; Berkner, *Biotechniques* 6:616-627, 1988; Rosenfeld et al., *Science*
20 252:431-434, 1991; Kolls et al., *Proc. Natl. Acad. Sci. USA* 91:215-219, 1994; Kass-Eisler et al., *Proc. Natl. Acad. Sci. USA* 90:11498-11502, 1993; Guzman et al., *Circulation* 88:2838-2848, 1993; and Guzman et al., *Cir. Res.* 73:1202-1207, 1993.

In certain embodiments, a polynucleotide may be integrated into the genome of a target cell. This integration may be in the specific location and orientation
25 via homologous recombination (gene replacement) or it may be integrated in a random, non-specific location (gene augmentation). In yet further embodiments, the polynucleotide may be stably maintained in the cell as a separate, episomal segment of DNA. Such polynucleotide segments or "episomes" encode sequences sufficient to permit maintenance and replication independent of or in synchronization with the host
30 cell cycle. The manner in which the expression construct is delivered to a cell and

where in the cell the polynucleotide remains is dependent on the type of expression construct employed.

In another embodiment of the invention, a polynucleotide is administered/delivered as "naked" DNA, for example as described in Ulmer et al.,
5 *Science* 259:1745-1749, 1993 and reviewed by Cohen, *Science* 259:1691-1692, 1993. The uptake of naked DNA may be increased by coating the DNA onto biodegradable beads, which are efficiently transported into the cells.

In still another embodiment, a composition of the present invention can be delivered via a particle bombardment approach, many of which have been described.
10 In one illustrative example, gas-driven particle acceleration can be achieved with devices such as those manufactured by Powderject Pharmaceuticals PLC (Oxford, UK) and Powderject Vaccines Inc. (Madison, WI), some examples of which are described in U.S. Patent Nos. 5,846,796; 6,010,478; 5,865,796; 5,584,807; and EP Patent No. 0500 799. This approach offers a needle-free delivery approach wherein a dry powder
15 formulation of microscopic particles, such as polynucleotide or polypeptide particles, are accelerated to high speed within a helium gas jet generated by a hand held device, propelling the particles into a target tissue of interest.

In a related embodiment, other devices and methods that may be useful for gas-driven needle-less injection of compositions of the present invention include
20 those provided by Bioject, Inc. (Portland, OR), some examples of which are described in U.S. Patent Nos. 4,790,824; 5,064,413; 5,312,335; 5,383,851; 5,399,163; 5,520,639 and 5,993,412.

According to another embodiment, the pharmaceutical compositions described herein will comprise one or more immunostimulants in addition to the
25 immunogenic polynucleotide, polypeptide, antibody, T-cell, TCR, and/or APC compositions of this invention. An immunostimulant refers to essentially any substance that enhances or potentiates an immune response (antibody and/or cell-mediated) to an exogenous antigen. One preferred type of immunostimulant comprises an adjuvant. Many adjuvants contain a substance designed to protect the antigen from rapid
30 catabolism, such as aluminum hydroxide or mineral oil, and a stimulator of immune responses, such as lipid A, *Bordetella pertussis* or *Mycobacterium tuberculosis* derived

proteins. Certain adjuvants are commercially available as, for example, Freund's Incomplete Adjuvant and Complete Adjuvant (Difco Laboratories, Detroit, MI); Merck Adjuvant 65 (Merck and Company, Inc., Rahway, NJ); AS-2 (SmithKline Beecham, Philadelphia, PA); aluminum salts such as aluminum hydroxide gel (alum) or aluminum phosphate; salts of calcium, iron or zinc; an insoluble suspension of acylated tyrosine; acylated sugars; cationically or anionically derivatized polysaccharides; polyphosphazenes; biodegradable microspheres; monophosphoryl lipid A and quil A. Cytokines, such as GM-CSF, interleukin-2, -7, -12, and other like growth factors, may also be used as adjuvants.

10 Within certain embodiments of the invention, the adjuvant composition is preferably one that induces an immune response predominantly of the Th1 type. High levels of Th1-type cytokines (*e.g.*, IFN- γ , TNF α , IL-2 and IL-12) tend to favor the induction of cell mediated immune responses to an administered antigen. In contrast, high levels of Th2-type cytokines (*e.g.*, IL-4, IL-5, IL-6 and IL-10) tend to favor the
15 induction of humoral immune responses. Following application of a vaccine as provided herein, a patient will support an immune response that includes Th1- and Th2-type responses. Within a preferred embodiment, in which a response is predominantly Th1-type, the level of Th1-type cytokines will increase to a greater extent than the level of Th2-type cytokines. The levels of these cytokines may be readily assessed using
20 standard assays. For a review of the families of cytokines, see Mosmann and Coffman, *Ann. Rev. Immunol.* 7:145-173, 1989.

Certain preferred adjuvants for eliciting a predominantly Th1-type response include, for example, a combination of monophosphoryl lipid A, preferably 3-de-O-acylated monophosphoryl lipid A, together with an aluminum salt. MPL[®]
25 adjuvants are available from Corixa Corporation (Seattle, WA; *see*, for example, US Patent Nos. 4,436,727; 4,877,611; 4,866,034 and 4,912,094). CpG-containing oligonucleotides (in which the CpG dinucleotide is unmethylated) also induce a predominantly Th1 response. Such oligonucleotides are well known and are described, for example, in WO 96/02555, WO 99/33488 and U.S. Patent Nos. 6,008,200 and
30 5,856,462. Immunostimulatory DNA sequences are also described, for example, by Sato et al., *Science* 273:352, 1996. Another preferred adjuvant comprises a saponin,

such as Quil A, or derivatives thereof, including QS21 and QS7 (Aquila Biopharmaceuticals Inc., Framingham, MA); Escin; Digitonin; or *Gypsophila* or *Chenopodium quinoa* saponins. Other preferred formulations include more than one saponin in the adjuvant combinations of the present invention, for example combinations of at least two of the following group comprising QS21, QS7, Quil A, β -escin, or digitonin.


Alternatively the saponin formulations may be combined with vaccine vehicles composed of chitosan or other polycationic polymers, polylactide and polylactide-co-glycolide particles, poly-N-acetyl glucosamine-based polymer matrix, particles composed of polysaccharides or chemically modified polysaccharides, liposomes and lipid-based particles, particles composed of glycerol monoesters, etc. The saponins may also be formulated in the presence of cholesterol to form particulate structures such as liposomes or ISCOMs. Furthermore, the saponins may be formulated together with a polyoxyethylene ether or ester, in either a non-particulate solution or suspension, or in a particulate structure such as a paucilamellar liposome or ISCOM. The saponins may also be formulated with excipients such as Carbopol^R to increase viscosity, or may be formulated in a dry powder form with a powder excipient such as lactose.

In one preferred embodiment, the adjuvant system includes the combination of a monophosphoryl lipid A and a saponin derivative, such as the combination of QS21 and 3D-MPL[®] adjuvant, as described in WO 94/00153, or a less reactogenic composition where the QS21 is quenched with cholesterol, as described in WO 96/33739. Other preferred formulations comprise an oil-in-water emulsion and tocopherol. Another particularly preferred adjuvant formulation employing QS21, 3D-MPL[®] adjuvant and tocopherol in an oil-in-water emulsion is described in WO 95/17210.

Another enhanced adjuvant system involves the combination of a CpG-containing oligonucleotide and a saponin derivative particularly the combination of CpG and QS21 is disclosed in WO 00/09159. Preferably the formulation additionally comprises an oil in water emulsion and tocopherol.

Additional illustrative adjuvants for use in the pharmaceutical compositions of the invention include Montanide ISA 720 (Seppic, France), SAF (Chiron, California, United States), ISCOMS (CSL), MF-59 (Chiron), the SBAS series of adjuvants (*e.g.*, SBAS-2 or SBAS-4, available from SmithKline Beecham, Rixensart, Belgium), Detox (Enhazyn[®]) (Corixa, Hamilton, MT), RC-529 (Corixa, Hamilton, MT) and other aminoalkyl glucosaminide 4-phosphates (AGPs), such as those described in pending U.S. Patent Application Serial Nos. 08/853,826 and 09/074,720, the disclosures of which are incorporated herein by reference in their entireties, and polyoxyethylene ether adjuvants such as those described in WO 99/52549A1.

Other preferred adjuvants include adjuvant molecules of the general formula



wherein, n is 1-50, A is a bond or $-\text{C}(\text{O})-$, R is C_{1-50} alkyl or Phenyl C_{1-50} alkyl.

One embodiment of the present invention consists of a vaccine formulation comprising a polyoxyethylene ether of general formula (I), wherein n is between 1 and 50, preferably 4-24, most preferably 9; the R component is C_{1-50} , preferably $\text{C}_4\text{-C}_{20}$ alkyl and most preferably C_{12} alkyl, and A is a bond. The concentration of the polyoxyethylene ethers should be in the range 0.1-20%, preferably from 0.1-10%, and most preferably in the range 0.1-1%. Preferred polyoxyethylene ethers are selected from the following group: polyoxyethylene-9-lauryl ether, polyoxyethylene-9-stearyl ether, polyoxyethylene-8-stearyl ether, polyoxyethylene-4-lauryl ether, polyoxyethylene-35-lauryl ether, and polyoxyethylene-23-lauryl ether. Polyoxyethylene ethers such as polyoxyethylene lauryl ether are described in the Merck index (12th edition: entry 7717). These adjuvant molecules are described in WO 99/52549.

The polyoxyethylene ether according to the general formula (I) above may, if desired, be combined with another adjuvant. For example, a preferred adjuvant combination is preferably with CpG as described in the pending UK patent application GB 9820956.2.

According to another embodiment of this invention, an immunogenic composition described herein is delivered to a host via antigen presenting cells (APCs),

such as dendritic cells, macrophages, B cells, monocytes and other cells that may be engineered to be efficient APCs. Such cells may, but need not, be genetically modified to increase the capacity for presenting the antigen, to improve activation and/or maintenance of the T cell response, to have anti-tumor effects *per se* and/or to be immunologically compatible with the receiver (*i.e.*, matched HLA haplotype). APCs may generally be isolated from any of a variety of biological fluids and organs, including tumor and peritumoral tissues, and may be autologous, allogeneic, syngeneic or xenogeneic cells.

Certain preferred embodiments of the present invention use dendritic cells or progenitors thereof as antigen-presenting cells. Dendritic cells are highly potent APCs (Banchereau and Steinman, *Nature* 392:245-251, 1998) and have been shown to be effective as a physiological adjuvant for eliciting prophylactic or therapeutic antitumor immunity (*see* Timmerman and Levy, *Ann. Rev. Med.* 50:507-529, 1999). In general, dendritic cells may be identified based on their typical shape (stellate *in situ*, with marked cytoplasmic processes (dendrites) visible *in vitro*), their ability to take up, process and present antigens with high efficiency and their ability to activate naïve T cell responses. Dendritic cells may, of course, be engineered to express specific cell-surface receptors or ligands that are not commonly found on dendritic cells *in vivo* or *ex vivo*, and such modified dendritic cells are contemplated by the present invention. As an alternative to dendritic cells, secreted vesicles antigen-loaded dendritic cells (called exosomes) may be used within a vaccine (*see* Zitvogel et al., *Nature Med.* 4:594-600, 1998).

Dendritic cells and progenitors may be obtained from peripheral blood, bone marrow, tumor-infiltrating cells, peritumoral tissues-infiltrating cells, lymph nodes, spleen, skin, umbilical cord blood or any other suitable tissue or fluid. For example, dendritic cells may be differentiated *ex vivo* by adding a combination of cytokines such as GM-CSF, IL-4, IL-13 and/or TNF α to cultures of monocytes harvested from peripheral blood. Alternatively, CD34 positive cells harvested from peripheral blood, umbilical cord blood or bone marrow may be differentiated into dendritic cells by adding to the culture medium combinations of GM-CSF, IL-3, TNF α ,

CD40 ligand, LPS, flt3 ligand and/or other compound(s) that induce differentiation, maturation and proliferation of dendritic cells.

Dendritic cells are conveniently categorized as "immature" and "mature" cells, which allows a simple way to discriminate between two well characterized phenotypes. However, this nomenclature should not be construed to exclude all possible intermediate stages of differentiation. Immature dendritic cells are characterized as APC with a high capacity for antigen uptake and processing, which correlates with the high expression of Fcγ receptor and mannose receptor. The mature phenotype is typically characterized by a lower expression of these markers, but a high expression of cell surface molecules responsible for T cell activation such as class I and class II MHC, adhesion molecules (e.g., CD54 and CD11) and costimulatory molecules (e.g., CD40, CD80, CD86 and 4-1BB).

APCs may generally be transfected with a polynucleotide of the invention (or portion or other variant thereof) such that the encoded polypeptide, or an immunogenic portion thereof, is expressed on the cell surface. Such transfection may take place *ex vivo*, and a pharmaceutical composition comprising such transfected cells may then be used for therapeutic purposes, as described herein. Alternatively, a gene delivery vehicle that targets a dendritic or other antigen presenting cell may be administered to a patient, resulting in transfection that occurs *in vivo*. *In vivo* and *ex vivo* transfection of dendritic cells, for example, may generally be performed using any methods known in the art, such as those described in WO 97/24447, or the gene gun approach described by Mahvi et al., *Immunology and cell Biology* 75:456-460, 1997. Antigen loading of dendritic cells may be achieved by incubating dendritic cells or progenitor cells with the tumor polypeptide, DNA (naked or within a plasmid vector) or RNA; or with antigen-expressing recombinant bacterium or viruses (e.g., vaccinia, fowlpox, adenovirus or lentivirus vectors). Prior to loading, the polypeptide may be covalently conjugated to an immunological partner that provides T cell help (e.g., a carrier molecule). Alternatively, a dendritic cell may be pulsed with a non-conjugated immunological partner, separately or in the presence of the polypeptide.

While any suitable carrier known to those of ordinary skill in the art may be employed in the pharmaceutical compositions of this invention, the type of carrier

will typically vary depending on the mode of administration. Compositions of the present invention may be formulated for any appropriate manner of administration, including for example, topical, oral, nasal, mucosal, intravenous, intracranial, intraperitoneal, subcutaneous and intramuscular administration.

5 Carriers for use within such pharmaceutical compositions are biocompatible, and may also be biodegradable. In certain embodiments, the formulation preferably provides a relatively constant level of active component release. In other embodiments, however, a more rapid rate of release immediately upon administration may be desired. The formulation of such compositions is well within the
10 level of ordinary skill in the art using known techniques. Illustrative carriers useful in this regard include microparticles of poly(lactide-co-glycolide), polyacrylate, latex, starch, cellulose, dextran and the like. Other illustrative delayed-release carriers include supramolecular biovectors, which comprise a non-liquid hydrophilic core (e.g., a cross-linked polysaccharide or oligosaccharide) and, optionally, an external layer
15 comprising an amphiphilic compound, such as a phospholipid (*see e.g.*, U.S. Patent No. 5,151,254 and PCT applications WO 94/20078, WO/94/23701 and WO 96/06638). The amount of active compound contained within a sustained release formulation depends upon the site of implantation, the rate and expected duration of release and the nature of the condition to be treated or prevented.

20 In another illustrative embodiment, biodegradable microspheres (e.g., polylactate polyglycolate) are employed as carriers for the compositions of this invention. Suitable biodegradable microspheres are disclosed, for example, in U.S. Patent Nos. 4,897,268; 5,075,109; 5,928,647; 5,811,128; 5,820,883; 5,853,763; 5,814,344, 5,407,609 and 5,942,252. Modified hepatitis B core protein carrier systems.
25 such as described in WO/99 40934, and references cited therein, will also be useful for many applications. Another illustrative carrier/delivery system employs a carrier comprising particulate-protein complexes, such as those described in U.S. Patent No. 5,928,647, which are capable of inducing a class I-restricted cytotoxic T lymphocyte responses in a host.

30 In another illustrative embodiment, calcium phosphate core particles are employed as carriers, vaccine adjuvants, or as controlled release matrices for the

compositions of this invention. Exemplary calcium phosphate particles are disclosed, for example, in published patent application No. WO/0046147.

The pharmaceutical compositions of the invention will often further comprise one or more buffers (*e.g.*, neutral buffered saline or phosphate buffered saline), carbohydrates (*e.g.*, glucose, mannose, sucrose or dextrans), mannitol, proteins, polypeptides or amino acids such as glycine, antioxidants, bacteriostats, chelating agents such as EDTA or glutathione, adjuvants (*e.g.*, aluminum hydroxide), solutes that render the formulation isotonic, hypotonic or weakly hypertonic with the blood of a recipient, suspending agents, thickening agents and/or preservatives. Alternatively, compositions of the present invention may be formulated as a lyophilizate.

The pharmaceutical compositions described herein may be presented in unit-dose or multi-dose containers, such as sealed ampoules or vials. Such containers are typically sealed in such a way to preserve the sterility and stability of the formulation until use. In general, formulations may be stored as suspensions, solutions or emulsions in oily or aqueous vehicles. Alternatively, a pharmaceutical composition may be stored in a freeze-dried condition requiring only the addition of a sterile liquid carrier immediately prior to use.

The development of suitable dosing and treatment regimens for using the particular compositions described herein in a variety of treatment regimens, including *e.g.*, oral, parenteral, intravenous, intranasal, and intramuscular administration and formulation, is well known in the art, some of which are briefly discussed below for general purposes of illustration.

In certain applications, the pharmaceutical compositions disclosed herein may be delivered *via* oral administration to an animal. As such, these compositions may be formulated with an inert diluent or with an assimilable edible carrier, or they may be enclosed in hard- or soft-shell gelatin capsule, or they may be compressed into tablets, or they may be incorporated directly with the food of the diet.

The active compounds may even be incorporated with excipients and used in the form of ingestible tablets, buccal tables, troches, capsules, elixirs, suspensions, syrups, wafers, and the like (see, for example, Mathiowitz *et al.*, Nature 1997 Mar 27;386(6623):410-4; Hwang *et al.*, Crit Rev Ther Drug Carrier Syst

1998;15(3):243-84; U. S. Patent 5,641,515; U. S. Patent 5,580,579 and U. S. Patent 5,792,451). Tablets, troches, pills, capsules and the like may also contain any of a variety of additional components, for example, a binder, such as gum tragacanth, acacia, cornstarch, or gelatin; excipients, such as dicalcium phosphate; a disintegrating agent, such as corn starch, potato starch, alginic acid and the like; a lubricant, such as magnesium stearate; and a sweetening agent, such as sucrose, lactose or saccharin may be added or a flavoring agent, such as peppermint, oil of wintergreen, or cherry flavoring. When the dosage unit form is a capsule, it may contain, in addition to materials of the above type, a liquid carrier. Various other materials may be present as coatings or to otherwise modify the physical form of the dosage unit. For instance, tablets, pills, or capsules may be coated with shellac, sugar, or both. Of course, any material used in preparing any dosage unit form should be pharmaceutically pure and substantially non-toxic in the amounts employed. In addition, the active compounds may be incorporated into sustained-release preparation and formulations.

Typically, these formulations will contain at least about 0.1% of the active compound or more, although the percentage of the active ingredient(s) may, of course, be varied and may conveniently be between about 1 or 2% and about 60% or 70% or more of the weight or volume of the total formulation. Naturally, the amount of active compound(s) in each therapeutically useful composition may be prepared in such a way that a suitable dosage will be obtained in any given unit dose of the compound. Factors such as solubility, bioavailability, biological half-life, route of administration, product shelf life, as well as other pharmacological considerations will be contemplated by one skilled in the art of preparing such pharmaceutical formulations, and as such, a variety of dosages and treatment regimens may be desirable.

For oral administration the compositions of the present invention may alternatively be incorporated with one or more excipients in the form of a mouthwash, dentifrice, buccal tablet, oral spray, or sublingual orally-administered formulation. Alternatively, the active ingredient may be incorporated into an oral solution such as one containing sodium borate, glycerin and potassium bicarbonate, or dispersed in a dentifrice, or added in a therapeutically-effective amount to a composition that may include water, binders, abrasives, flavoring agents, foaming agents, and humectants.

Alternatively the compositions may be fashioned into a tablet or solution form that may be placed under the tongue or otherwise dissolved in the mouth.

In certain circumstances it will be desirable to deliver the pharmaceutical compositions disclosed herein parenterally, intravenously, intramuscularly, or even intraperitoneally. Such approaches are well known to the skilled artisan, some of which are further described, for example, in U. S. Patent 5,543,158; U. S. Patent 5,641,515 and U. S. Patent 5,399,363. In certain embodiments, solutions of the active compounds as free base or pharmacologically acceptable salts may be prepared in water suitably mixed with a surfactant, such as hydroxypropylcellulose. Dispersions may also be prepared in glycerol, liquid polyethylene glycols, and mixtures thereof and in oils. Under ordinary conditions of storage and use, these preparations generally will contain a preservative to prevent the growth of microorganisms.

Illustrative pharmaceutical forms suitable for injectable use include sterile aqueous solutions or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions (for example, see U. S. Patent 5,466,468). In all cases the form must be sterile and must be fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms, such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (*e.g.*, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), suitable mixtures thereof, and/or vegetable oils. Proper fluidity may be maintained, for example, by the use of a coating, such as lecithin, by the maintenance of the required particle size in the case of dispersion and/or by the use of surfactants. The prevention of the action of microorganisms can be facilitated by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars or sodium chloride. Prolonged absorption of the injectable compositions can be brought about by the use in the compositions of agents delaying absorption, for example, aluminum monostearate and gelatin.

In one embodiment, for parenteral administration in an aqueous solution, the solution should be suitably buffered if necessary and the liquid diluent first rendered isotonic with sufficient saline or glucose. These particular aqueous solutions are especially suitable for intravenous, intramuscular, subcutaneous and intraperitoneal administration. In this connection, a sterile aqueous medium that can be employed will be known to those of skill in the art in light of the present disclosure. For example, one dosage may be dissolved in 1 ml of isotonic NaCl solution and either added to 1000 ml of hypodermoclysis fluid or injected at the proposed site of infusion, (see for example, "Remington's Pharmaceutical Sciences" 15th Edition, pages 1035-1038 and 1570-1580). Some variation in dosage will necessarily occur depending on the condition of the subject being treated. Moreover, for human administration, preparations will of course preferably meet sterility, pyrogenicity, and the general safety and purity standards as required by FDA Office of Biologics standards.

In another embodiment of the invention, the compositions disclosed herein may be formulated in a neutral or salt form. Illustrative pharmaceutically-acceptable salts include the acid addition salts (formed with the free amino groups of the protein) and which are formed with inorganic acids such as, for example, hydrochloric or phosphoric acids, or such organic acids as acetic, oxalic, tartaric, mandelic, and the like. Salts formed with the free carboxyl groups can also be derived from inorganic bases such as, for example, sodium, potassium, ammonium, calcium, or ferric hydroxides, and such organic bases as isopropylamine, trimethylamine, histidine, procaine and the like. Upon formulation, solutions will be administered in a manner compatible with the dosage formulation and in such amount as is therapeutically effective.

The carriers can further comprise any and all solvents, dispersion media, vehicles, coatings, diluents, antibacterial and antifungal agents, isotonic and absorption delaying agents, buffers, carrier solutions, suspensions, colloids, and the like. The use of such media and agents for pharmaceutical active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active ingredient, its use in the therapeutic compositions is contemplated. Supplementary active ingredients can also be incorporated into the compositions. The phrase

"pharmaceutically-acceptable" refers to molecular entities and compositions that do not produce an allergic or similar untoward reaction when administered to a human.

In certain embodiments, the pharmaceutical compositions may be delivered by intranasal sprays, inhalation, and/or other aerosol delivery vehicles.

5 Methods for delivering genes, nucleic acids, and peptide compositions directly to the lungs *via* nasal aerosol sprays has been described, *e.g.*, in U. S. Patent 5,756,353 and U. S. Patent 5,804,212. Likewise, the delivery of drugs using intranasal microparticle resins (Takenaga *et al.*, J Controlled Release 1998 Mar 2;52(1-2):81-7) and lysophosphatidyl-glycerol compounds (U. S. Patent 5,725,871) are also well-known in

10 the pharmaceutical arts. Likewise, illustrative transmucosal drug delivery in the form of a polytetrafluoroethylene support matrix is described in U. S. Patent 5,780,045.

In certain embodiments, liposomes, nanocapsules, microparticles, lipid particles, vesicles, and the like, are used for the introduction of the compositions of the present invention into suitable host cells/organisms. In particular, the compositions of

15 the present invention may be formulated for delivery either encapsulated in a lipid particle, a liposome, a vesicle, a nanosphere, or a nanoparticle or the like. Alternatively, compositions of the present invention can be bound, either covalently or non-covalently, to the surface of such carrier vehicles.

The formation and use of liposome and liposome-like preparations as

20 potential drug carriers is generally known to those of skill in the art (see for example, Lasic, Trends Biotechnol 1998 Jul;16(7):307-21; Takakura, Nippon Rinsho 1998 Mar;56(3):691-5; Chandran *et al.*, Indian J Exp Biol. 1997 Aug;35(8):801-9; Margalit, Crit Rev Ther Drug Carrier Syst. 1995;12(2-3):233-61; U.S. Patent 5,567,434; U.S. Patent 5,552,157; U.S. Patent 5,565,213; U.S. Patent 5,738,868 and U.S. Patent

25 5,795,587, each specifically incorporated herein by reference in its entirety).

Liposomes have been used successfully with a number of cell types that are normally difficult to transfect by other procedures, including T cell suspensions, primary hepatocyte cultures and PC 12 cells (Renneisen *et al.*, J Biol Chem. 1990 Sep 25;265(27):16337-42; Muller *et al.*, DNA Cell Biol. 1990 Apr;9(3):221-9). In addition,

30 liposomes are free of the DNA length constraints that are typical of viral-based delivery systems. Liposomes have been used effectively to introduce genes, various drugs,

radiotherapeutic agents, enzymes, viruses, transcription factors, allosteric effectors and the like, into a variety of cultured cell lines and animals. Furthermore, the use of liposomes does not appear to be associated with autoimmune responses or unacceptable toxicity after systemic delivery.

- 5 In certain embodiments, liposomes are formed from phospholipids that are dispersed in an aqueous medium and spontaneously form multilamellar concentric bilayer vesicles (also termed multilamellar vesicles (MLVs)).

- Alternatively, in other embodiments, the invention provides for pharmaceutically-acceptable nanocapsule formulations of the compositions of the present invention. Nanocapsules can generally entrap compounds in a stable and reproducible way (see, for example, Quintanar-Guerrero *et al.*, Drug Dev Ind Pharm. 1998 Dec;24(12):1113-28). To avoid side effects due to intracellular polymeric overloading, such ultrafine particles (sized around 0.1 μm) may be designed using polymers able to be degraded *in vivo*. Such particles can be made as described, for example, by Couvreur *et al.*, Crit Rev Ther Drug Carrier Syst. 1988;5(1):1-20; zur Muhlen *et al.*, Eur J Pharm Biopharm. 1998 Mar;45(2):149-55; Zambaux *et al.* J Controlled Release. 1998 Jan 2;50(1-3):31-40; and U. S. Patent 5,145,684.

Cancer Therapeutic Methods

- Immunologic approaches to cancer therapy are based on the recognition that cancer cells can often evade the body's defenses against aberrant or foreign cells and molecules, and that these defenses might be therapeutically stimulated to regain the lost ground, *e.g.* pgs. 623-648 in Klein, Immunology (Wiley-Interscience, New York, 1982). Numerous recent observations that various immune effectors can directly or indirectly inhibit growth of tumors has led to renewed interest in this approach to cancer therapy, *e.g.* Jager, *et al.*, Oncology 2001;60(1):1-7; Renner, *et al.*, Ann Hematol 2000 Dec;79(12):651-9.

- Four basic cell types whose function has been associated with antitumor cell immunity and the elimination of tumor cells from the body are: i) B-lymphocytes which secrete immunoglobulins into the blood plasma for identifying and labeling the nonself invader cells; ii) monocytes which secrete the complement proteins that are

responsible for lysing and processing the immunoglobulin-coated target invader cells; iii) natural killer lymphocytes having two mechanisms for the destruction of tumor cells, antibody-dependent cellular cytotoxicity and natural killing; and iv) T-lymphocytes possessing antigen-specific receptors and having the capacity to recognize a tumor cell carrying complementary marker molecules (Schreiber, H., 1989, in Fundamental Immunology (ed) W. E. Paul, pp. 923-955).

Cancer immunotherapy generally focuses on inducing humoral immune responses, cellular immune responses, or both. Moreover, it is well established that induction of CD4⁺ T helper cells is necessary in order to secondarily induce either antibodies or cytotoxic CD8⁺ T cells. Polypeptide antigens that are selective or ideally specific for cancer cells, particularly lung cancer cells, offer a powerful approach for inducing immune responses against lung cancer, and are an important aspect of the present invention.

Therefore, in further aspects of the present invention, the pharmaceutical compositions described herein may be used to stimulate an immune response against cancer, particularly for the immunotherapy of lung cancer. Within such methods, the pharmaceutical compositions described herein are administered to a patient, typically a warm-blooded animal, preferably a human. A patient may or may not be afflicted with cancer. Pharmaceutical compositions and vaccines may be administered either prior to or following surgical removal of primary tumors and/or treatment such as administration of radiotherapy or conventional chemotherapeutic drugs. As discussed above, administration of the pharmaceutical compositions may be by any suitable method, including administration by intravenous, intraperitoneal, intramuscular, subcutaneous, intranasal, intradermal, anal, vaginal, topical and oral routes.

Within certain embodiments, immunotherapy may be active immunotherapy, in which treatment relies on the *in vivo* stimulation of the endogenous host immune system to react against tumors with the administration of immune response-modifying agents (such as polypeptides and polynucleotides as provided herein).

Within other embodiments, immunotherapy may be passive immunotherapy, in which treatment involves the delivery of agents with established

tumor-immune reactivity (such as effector cells or antibodies) that can directly or indirectly mediate antitumor effects and does not necessarily depend on an intact host immune system. Examples of effector cells include T cells as discussed above, T lymphocytes (such as CD8⁺ cytotoxic T lymphocytes and CD4⁺ T-helper tumor-infiltrating lymphocytes), killer cells (such as Natural Killer cells and lymphokine-activated killer cells), B cells and antigen-presenting cells (such as dendritic cells and macrophages) expressing a polypeptide provided herein. T cell receptors and antibody receptors specific for the polypeptides recited herein may be cloned, expressed and transferred into other vectors or effector cells for adoptive immunotherapy. The polypeptides provided herein may also be used to generate antibodies or anti-idiotypic antibodies (as described above and in U.S. Patent No. 4,918,164) for passive immunotherapy.

Monoclonal antibodies may be labeled with any of a variety of labels for desired selective usages in detection, diagnostic assays or therapeutic applications (as described in U.S. Patent Nos. 6,090,365; 6,015,542; 5,843,398; 5,595,721; and 4,708,930, hereby incorporated by reference in their entirety as if each was incorporated individually). In each case, the binding of the labelled monoclonal antibody to the determinant site of the antigen will signal detection or delivery of a particular therapeutic agent to the antigenic determinant on the non-normal cell. A further object of this invention is to provide the specific monoclonal antibody suitably labelled for achieving such desired selective usages thereof.

Effector cells may generally be obtained in sufficient quantities for adoptive immunotherapy by growth *in vitro*, as described herein. Culture conditions for expanding single antigen-specific effector cells to several billion in number with retention of antigen recognition *in vivo* are well known in the art. Such *in vitro* culture conditions typically use intermittent stimulation with antigen, often in the presence of cytokines (such as IL-2) and non-dividing feeder cells. As noted above, immunoreactive polypeptides as provided herein may be used to rapidly expand antigen-specific T cell cultures in order to generate a sufficient number of cells for immunotherapy. In particular, antigen-presenting cells, such as dendritic, macrophage, monocyte, fibroblast and/or B cells, may be pulsed with immunoreactive polypeptides

or transfected with one or more polynucleotides using standard techniques well known in the art. For example, antigen-presenting cells can be transfected with a polynucleotide having a promoter appropriate for increasing expression in a recombinant virus or other expression system. Cultured effector cells for use in therapy must be able to grow and distribute widely, and to survive long term *in vivo*. Studies have shown that cultured effector cells can be induced to grow *in vivo* and to survive long term in substantial numbers by repeated stimulation with antigen supplemented with IL-2 (*see, for example, Cheever et al., Immunological Reviews 157:177, 1997.*

Alternatively, a vector expressing a polypeptide recited herein may be introduced into antigen presenting cells taken from a patient and clonally propagated *ex vivo* for transplant back into the same patient. Transfected cells may be reintroduced into the patient using any means known in the art, preferably in sterile form by intravenous, intracavitary, intraperitoneal or intratumor administration.

Routes and frequency of administration of the therapeutic compositions described herein, as well as dosage, will vary from individual to individual, and may be readily established using standard techniques. In general, the pharmaceutical compositions and vaccines may be administered by injection (*e.g.,* intracutaneous, intramuscular, intravenous or subcutaneous), intranasally (*e.g.,* by aspiration) or orally. Preferably, between 1 and 10 doses may be administered over a 52 week period. Preferably, 6 doses are administered, at intervals of 1 month, and booster vaccinations may be given periodically thereafter. Alternate protocols may be appropriate for individual patients. A suitable dose is an amount of a compound that, when administered as described above, is capable of promoting an anti-tumor immune response, and is at least 10-50% above the basal (*i.e.,* untreated) level. Such response can be monitored by measuring the anti-tumor antibodies in a patient or by vaccine-dependent generation of cytolytic effector cells capable of killing the patient's tumor cells *in vitro*. Such vaccines should also be capable of causing an immune response that leads to an improved clinical outcome (*e.g.,* more frequent remissions, complete or partial or longer disease-free survival) in vaccinated patients as compared to non-vaccinated patients. In general, for pharmaceutical compositions and vaccines comprising one or more polypeptides, the amount of each polypeptide present in a dose

ranges from about 25 μg to 5 mg per kg of host. Suitable dose sizes will vary with the size of the patient, but will typically range from about 0.1 mL to about 5 mL.

In general, an appropriate dosage and treatment regimen provides the active compound(s) in an amount sufficient to provide therapeutic and/or prophylactic benefit. Such a response can be monitored by establishing an improved clinical outcome (e.g., more frequent remissions, complete or partial, or longer disease-free survival) in treated patients as compared to non-treated patients. Increases in preexisting immune responses to a tumor protein generally correlate with an improved clinical outcome. Such immune responses may generally be evaluated using standard proliferation, cytotoxicity or cytokine assays, which may be performed using samples obtained from a patient before and after treatment.

Cancer Detection and Diagnostic Compositions, Methods and Kits

In general, a cancer may be detected in a patient based on the presence of one or more lung tumor proteins and/or polynucleotides encoding such proteins in a biological sample (for example, blood, sera, sputum urine and/or tumor biopsies) obtained from the patient. In other words, such proteins may be used as markers to indicate the presence or absence of a cancer such as lung cancer. In addition, such proteins may be useful for the detection of other cancers. The binding agents provided herein generally permit detection of the level of antigen that binds to the agent in the biological sample.

Polynucleotide primers and probes may be used to detect the level of mRNA encoding a tumor protein, which is also indicative of the presence or absence of a cancer. In general, a tumor sequence should be present at a level that is at least two-fold, preferably three-fold, and more preferably five-fold or higher in tumor tissue than in normal tissue of the same type from which the tumor arose. Expression levels of a particular tumor sequence in tissue types different from that in which the tumor arose are irrelevant in certain diagnostic embodiments since the presence of tumor cells can be confirmed by observation of predetermined differential expression levels, e.g., 2-fold, 5-fold, etc, in tumor tissue to expression levels in normal tissue of the same type.

Other differential expression patterns can be utilized advantageously for diagnostic purposes. For example, in one aspect of the invention, overexpression of a tumor sequence in tumor tissue and normal tissue of the same type, but not in other normal tissue types, e.g. PBMCs, can be exploited diagnostically. In this case, the presence of metastatic tumor cells, for example in a sample taken from the circulation or some other tissue site different from that in which the tumor arose, can be identified and/or confirmed by detecting expression of the tumor sequence in the sample, for example using RT-PCR analysis. In many instances, it will be desired to enrich for tumor cells in the sample of interest, e.g., PBMCs, using cell capture or other like techniques.

There are a variety of assay formats known to those of ordinary skill in the art for using a binding agent to detect polypeptide markers in a sample. See, e.g., Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, 1988. In general, the presence or absence of a cancer in a patient may be determined by (a) contacting a biological sample obtained from a patient with a binding agent; (b) detecting in the sample a level of polypeptide that binds to the binding agent; and (c) comparing the level of polypeptide with a predetermined cut-off value.

In a preferred embodiment, the assay involves the use of binding agent immobilized on a solid support to bind to and remove the polypeptide from the remainder of the sample. The bound polypeptide may then be detected using a detection reagent that contains a reporter group and specifically binds to the binding agent/polypeptide complex. Such detection reagents may comprise, for example, a binding agent that specifically binds to the polypeptide or an antibody or other agent that specifically binds to the binding agent, such as an anti-immunoglobulin, protein G, protein A or a lectin. Alternatively, a competitive assay may be utilized, in which a polypeptide is labeled with a reporter group and allowed to bind to the immobilized binding agent after incubation of the binding agent with the sample. The extent to which components of the sample inhibit the binding of the labeled polypeptide to the binding agent is indicative of the reactivity of the sample with the immobilized binding agent. Suitable polypeptides for use within such assays include full length lung tumor

proteins and polypeptide portions thereof to which the binding agent binds, as described above.

The solid support may be any material known to those of ordinary skill in the art to which the tumor protein may be attached. For example, the solid support may be a test well in a microtiter plate or a nitrocellulose or other suitable membrane. Alternatively, the support may be a bead or disc, such as glass, fiberglass, latex or a plastic material such as polystyrene or polyvinylchloride. The support may also be a magnetic particle or a fiber optic sensor, such as those disclosed, for example, in U.S. Patent No. 5,359,681. The binding agent may be immobilized on the solid support using a variety of techniques known to those of skill in the art, which are amply described in the patent and scientific literature. In the context of the present invention, the term "immobilization" refers to both noncovalent association, such as adsorption, and covalent attachment (which may be a direct linkage between the agent and functional groups on the support or may be a linkage by way of a cross-linking agent). Immobilization by adsorption to a well in a microtiter plate or to a membrane is preferred. In such cases, adsorption may be achieved by contacting the binding agent, in a suitable buffer, with the solid support for a suitable amount of time. The contact time varies with temperature, but is typically between about 1 hour and about 1 day. In general, contacting a well of a plastic microtiter plate (such as polystyrene or polyvinylchloride) with an amount of binding agent ranging from about 10 ng to about 10 μ g, and preferably about 100 ng to about 1 μ g, is sufficient to immobilize an adequate amount of binding agent.

Covalent attachment of binding agent to a solid support may generally be achieved by first reacting the support with a bifunctional reagent that will react with both the support and a functional group, such as a hydroxyl or amino group, on the binding agent. For example, the binding agent may be covalently attached to supports having an appropriate polymer coating using benzoquinone or by condensation of an aldehyde group on the support with an amine and an active hydrogen on the binding partner (*see, e.g.,* Pierce Immunotechnology Catalog and Handbook, 1991, at A12-A13).

In certain embodiments, the assay is a two-antibody sandwich assay.

This assay may be performed by first contacting an antibody that has been immobilized on a solid support, commonly the well of a microtiter plate, with the sample, such that polypeptides within the sample are allowed to bind to the immobilized antibody.

- 5 Unbound sample is then removed from the immobilized polypeptide-antibody complexes and a detection reagent (preferably a second antibody capable of binding to a different site on the polypeptide) containing a reporter group is added. The amount of detection reagent that remains bound to the solid support is then determined using a method appropriate for the specific reporter group.

- 10 More specifically, once the antibody is immobilized on the support as described above, the remaining protein binding sites on the support are typically blocked. Any suitable blocking agent known to those of ordinary skill in the art, such as bovine serum albumin or Tween 20™ (Sigma Chemical Co., St. Louis, MO). The immobilized antibody is then incubated with the sample, and polypeptide is allowed to
15 bind to the antibody. The sample may be diluted with a suitable diluent, such as phosphate-buffered saline (PBS) prior to incubation. In general, an appropriate contact time (*i.e.*, incubation time) is a period of time that is sufficient to detect the presence of polypeptide within a sample obtained from an individual with lung least about 95% of that achieved at equilibrium between bound and unbound polypeptide. Those of
20 ordinary skill in the art will recognize that the time necessary to achieve equilibrium may be readily determined by assaying the level of binding that occurs over a period of time. At room temperature, an incubation time of about 30 minutes is generally sufficient.

- Unbound sample may then be removed by washing the solid support
25 with an appropriate buffer, such as PBS containing 0.1% Tween 20™. The second antibody, which contains a reporter group, may then be added to the solid support. Preferred reporter groups include those groups recited above.

- The detection reagent is then incubated with the immobilized antibody-polypeptide complex for an amount of time sufficient to detect the bound polypeptide.
30 An appropriate amount of time may generally be determined by assaying the level of binding that occurs over a period of time. Unbound detection reagent is then removed

and bound detection reagent is detected using the reporter group. The method employed for detecting the reporter group depends upon the nature of the reporter group. For radioactive groups, scintillation counting or autoradiographic methods are generally appropriate. Spectroscopic methods may be used to detect dyes, luminescent groups and fluorescent groups. Biotin may be detected using avidin, coupled to a different reporter group (commonly a radioactive or fluorescent group or an enzyme). Enzyme reporter groups may generally be detected by the addition of substrate (generally for a specific period of time), followed by spectroscopic or other analysis of the reaction products.

To determine the presence or absence of a cancer, such as lung cancer, the signal detected from the reporter group that remains bound to the solid support is generally compared to a signal that corresponds to a predetermined cut-off value. In one preferred embodiment, the cut-off value for the detection of a cancer is the average mean signal obtained when the immobilized antibody is incubated with samples from patients without the cancer. In general, a sample generating a signal that is three standard deviations above the predetermined cut-off value is considered positive for the cancer. In an alternate preferred embodiment, the cut-off value is determined using a Receiver Operator Curve, according to the method of Sackett et al., *Clinical Epidemiology: A Basic Science for Clinical Medicine*, Little Brown and Co., 1985, p. 106-7. Briefly, in this embodiment, the cut-off value may be determined from a plot of pairs of true positive rates (*i.e.*, sensitivity) and false positive rates (100%-specificity) that correspond to each possible cut-off value for the diagnostic test result. The cut-off value on the plot that is the closest to the upper left-hand corner (*i.e.*, the value that encloses the largest area) is the most accurate cut-off value, and a sample generating a signal that is higher than the cut-off value determined by this method may be considered positive. Alternatively, the cut-off value may be shifted to the left along the plot, to minimize the false positive rate, or to the right, to minimize the false negative rate. In general, a sample generating a signal that is higher than the cut-off value determined by this method is considered positive for a cancer.

In a related embodiment, the assay is performed in a flow-through or strip test format, wherein the binding agent is immobilized on a membrane, such as

nitrocellulose. In the flow-through test, polypeptides within the sample bind to the immobilized binding agent as the sample passes through the membrane. A second, labeled binding agent then binds to the binding agent-polypeptide complex as a solution containing the second binding agent flows through the membrane. The detection of bound second binding agent may then be performed as described above. In the strip test format, one end of the membrane to which binding agent is bound is immersed in a solution containing the sample. The sample migrates along the membrane through a region containing second binding agent and to the area of immobilized binding agent. Concentration of second binding agent at the area of immobilized antibody indicates the presence of a cancer. Typically, the concentration of second binding agent at that site generates a pattern, such as a line, that can be read visually. The absence of such a pattern indicates a negative result. In general, the amount of binding agent immobilized on the membrane is selected to generate a visually discernible pattern when the biological sample contains a level of polypeptide that would be sufficient to generate a positive signal in the two-antibody sandwich assay, in the format discussed above. Preferred binding agents for use in such assays are antibodies and antigen-binding fragments thereof. Preferably, the amount of antibody immobilized on the membrane ranges from about 25 ng to about 1 μ g, and more preferably from about 50 ng to about 500 ng. Such tests can typically be performed with a very small amount of biological sample.

Of course, numerous other assay protocols exist that are suitable for use with the tumor proteins or binding agents of the present invention. The above descriptions are intended to be exemplary only. For example, it will be apparent to those of ordinary skill in the art that the above protocols may be readily modified to use tumor polypeptides to detect antibodies that bind to such polypeptides in a biological sample. The detection of such tumor protein specific antibodies may correlate with the presence of a cancer.

A cancer may also, or alternatively, be detected based on the presence of T cells that specifically react with a tumor protein in a biological sample. Within certain methods, a biological sample comprising CD4⁺ and/or CD8⁺ T cells isolated from a patient is incubated with a tumor polypeptide, a polynucleotide encoding such a

polypeptide and/or an APC that expresses at least an immunogenic portion of such a polypeptide, and the presence or absence of specific activation of the T cells is detected. Suitable biological samples include, but are not limited to, isolated T cells. For example, T cells may be isolated from a patient by routine techniques (such as by Ficol/Hypaque density gradient centrifugation of peripheral blood lymphocytes). T cells may be incubated *in vitro* for 2-9 days (typically 4 days) at 37°C with polypeptide (e.g., 5 - 25 µg/ml). It may be desirable to incubate another aliquot of a T cell sample in the absence of tumor polypeptide to serve as a control. For CD4⁺ T cells, activation is preferably detected by evaluating proliferation of the T cells. For CD8⁺ T cells, activation is preferably detected by evaluating cytolytic activity. A level of proliferation that is at least two fold greater and/or a level of cytolytic activity that is at least 20% greater than in disease-free patients indicates the presence of a cancer in the patient.

As noted above, a cancer may also, or alternatively, be detected based on the level of mRNA encoding a tumor protein in a biological sample. For example, at least two oligonucleotide primers may be employed in a polymerase chain reaction (PCR) based assay to amplify a portion of a tumor cDNA derived from a biological sample, wherein at least one of the oligonucleotide primers is specific for (*i.e.*, hybridizes to) a polynucleotide encoding the tumor protein. The amplified cDNA is then separated and detected using techniques well known in the art, such as gel electrophoresis.

Similarly, oligonucleotide probes that specifically hybridize to a polynucleotide encoding a tumor protein may be used in a hybridization assay to detect the presence of polynucleotide encoding the tumor protein in a biological sample.

To permit hybridization under assay conditions, oligonucleotide primers and probes should comprise an oligonucleotide sequence that has at least about 60%, preferably at least about 75% and more preferably at least about 90%, identity to a portion of a polynucleotide encoding a tumor protein of the invention that is at least 10 nucleotides, and preferably at least 20 nucleotides, in length. Preferably, oligonucleotide primers and/or probes hybridize to a polynucleotide encoding a polypeptide described herein under moderately stringent conditions, as defined above. Oligonucleotide primers and/or probes which may be usefully employed in the

diagnostic methods described herein preferably are at least 10-40 nucleotides in length. In a preferred embodiment, the oligonucleotide primers comprise at least 10 contiguous nucleotides, more preferably at least 15 contiguous nucleotides, of a DNA molecule having a sequence as disclosed herein. Techniques for both PCR based assays and hybridization assays are well known in the art (*see*, for example, Mullis et al., *Cold Spring Harbor Symp. Quant. Biol.*, 51:263, 1987; Erlich ed., *PCR Technology*, Stockton Press, NY, 1989).

One preferred assay employs RT-PCR, in which PCR is applied in conjunction with reverse transcription. Typically, RNA is extracted from a biological sample, such as biopsy tissue, and is reverse transcribed to produce cDNA molecules. PCR amplification using at least one specific primer generates a cDNA molecule, which may be separated and visualized using, for example, gel electrophoresis. Amplification may be performed on biological samples taken from a test patient and from an individual who is not afflicted with a cancer. The amplification reaction may be performed on several dilutions of cDNA spanning two orders of magnitude. A two-fold or greater increase in expression in several dilutions of the test patient sample as compared to the same dilutions of the non-cancerous sample is typically considered positive.

In another aspect of the present invention, cell capture technologies may be used in conjunction, with, for example, real-time PCR to provide a more sensitive tool for detection of metastatic cells expressing lung tumor antigens. Detection of lung cancer cells in biological samples, e.g., bone marrow samples, peripheral blood, and small needle aspiration samples is desirable for diagnosis and prognosis in lung cancer patients.

Immunomagnetic beads coated with specific monoclonal antibodies to surface cell markers, or tetrameric antibody complexes, may be used to first enrich or positively select cancer cells in a sample. Various commercially available kits may be used, including Dynabeads® Epithelial Enrich (DynaL Biotech, Oslo, Norway), StemSep™ (StemCell Technologies, Inc., Vancouver, BC), and RosetteSep (StemCell Technologies). A skilled artisan will recognize that other methodologies and kits may also be used to enrich or positively select desired cell populations. Dynabeads®

Epithelial Enrich contains magnetic beads coated with mAbs specific for two glycoprotein membrane antigens expressed on normal and neoplastic epithelial tissues. The coated beads may be added to a sample and the sample then applied to a magnet, thereby capturing the cells bound to the beads. The unwanted cells are washed away and the magnetically isolated cells eluted from the beads and used in further analyses.

RosetteSep can be used to enrich cells directly from a blood sample and consists of a cocktail of tetrameric antibodies that targets a variety of unwanted cells and crosslinks them to glycophorin A on red blood cells (RBC) present in the sample, forming rosettes. When centrifuged over Ficoll, targeted cells pellet along with the free RBC. The combination of antibodies in the depletion cocktail determines which cells will be removed and consequently which cells will be recovered. Antibodies that are available include, but are not limited to: CD2, CD3, CD4, CD5, CD8, CD10, CD11b, CD14, CD15, CD16, CD19, CD20, CD24, CD25, CD29, CD33, CD34, CD36, CD38, CD41, CD45, CD45RA, CD45RO, CD56, CD66B, CD66e, HLA-DR, IgE, and TCR $\alpha\beta$.

Additionally, it is contemplated in the present invention that mAbs specific for lung tumor antigens can be generated and used in a similar manner. For example, mAbs that bind to tumor-specific cell surface antigens may be conjugated to magnetic beads, or formulated in a tetrameric antibody complex, and used to enrich or positively select metastatic lung tumor cells from a sample. Once a sample is enriched or positively selected, cells may be lysed and RNA isolated. RNA may then be subjected to RT-PCR analysis using lung tumor-specific primers in a real-time PCR assay as described herein. One skilled in the art will recognize that enriched or selected populations of cells may be analyzed by other methods (*e.g. in situ* hybridization or flow cytometry).

In another embodiment, the compositions described herein may be used as markers for the progression of cancer. In this embodiment, assays as described above for the diagnosis of a cancer may be performed over time, and the change in the level of reactive polypeptide(s) or polynucleotide(s) evaluated. For example, the assays may be performed every 24-72 hours for a period of 6 months to 1 year, and thereafter performed as needed. In general, a cancer is progressing in those patients in whom the level of polypeptide or polynucleotide detected increases over time. In contrast, the

cancer is not progressing when the level of reactive polypeptide or polynucleotide either remains constant or decreases with time.

Certain *in vivo* diagnostic assays may be performed directly on a tumor. One such assay involves contacting tumor cells with a binding agent. The bound
5 binding agent may then be detected directly or indirectly via a reporter group. Such binding agents may also be used in histological applications. Alternatively, polynucleotide probes may be used within such applications.

As noted above, to improve sensitivity, multiple tumor protein markers may be assayed within a given sample. It will be apparent that binding agents specific
10 for different proteins provided herein may be combined within a single assay. Further, multiple primers or probes may be used concurrently. The selection of tumor protein markers may be based on routine experiments to determine combinations that results in optimal sensitivity. In addition, or alternatively, assays for tumor proteins provided herein may be combined with assays for other known tumor antigens.

15 The present invention further provides kits for use within any of the above diagnostic methods. Such kits typically comprise two or more components necessary for performing a diagnostic assay. Components may be compounds, reagents, containers and/or equipment. For example, one container within a kit may contain a monoclonal antibody or fragment thereof that specifically binds to a tumor protein.
20 Such antibodies or fragments may be provided attached to a support material, as described above. One or more additional containers may enclose elements, such as reagents or buffers, to be used in the assay. Such kits may also, or alternatively, contain a detection reagent as described above that contains a reporter group suitable for direct or indirect detection of antibody binding.

25 Alternatively, a kit may be designed to detect the level of mRNA encoding a tumor protein in a biological sample. Such kits generally comprise at least one oligonucleotide probe or primer, as described above, that hybridizes to a polynucleotide encoding a tumor protein. Such an oligonucleotide may be used, for example, within a PCR or hybridization assay. Additional components that may be
30 present within such kits include a second oligonucleotide and/or a diagnostic reagent or container to facilitate the detection of a polynucleotide encoding a tumor protein.

The following Examples are offered by way of illustration and not by way of limitation.

EXAMPLES

5

EXAMPLE 1

IDENTIFICATION OF LUNG TUMOR PROTEIN CDNAs

Lung-specific genes were identified by electronic subtraction. The method used was similar to that described by Vasmatizis et al., *Proc. Natl. Acad. Sci. USA* 95:300-304, 1998, but there were several key differences. Sequences of EST clones (1,453,679) were downloaded from the GenBank public human EST database. Human cDNA libraries were downloaded to create a database of these cDNA libraries and the EST sequences derived from them. The cDNA libraries were grouped into three groups: Plus, Minus and Other/Neutral. The Plus group included 30 libraries constructed from lung tumor and fetal lung tissues (and therefore including those containing lung tumor-specific ESTs); the Minus group consisted of 206 libraries derived from all adult normal tissues; the Other/Neutral group contained libraries from tissues where expression is considered irrelevant (e.g., non-lung-fetal tissue, non-lung tumors, cell lines other than lung tumor cell lines). A total of 93,526 ESTs were derived from the 30 lung tumor and fetal lung libraries. These ESTs were preprocessed to remove common sequence repeats and cloning adapters, resulting in a final Plus group of 90,365 (a decrease of 3%).

Each Plus group (lung tumor or fetal lung) EST sequence was used as a query "seed" sequence in a BLASTN (version 2.0.9; May 7, 1999) search against the total human EST database. Standard measures of similarity are insufficient in this sort of analysis, as EST relationships often include short stretches and poor sequence data. Criteria employed in this study required a matching segment to be at least 75 nucleotides in length, and the density of exact matches within this segment to be at least 80%. This was considered conservative criteria designed to avoid short spurious matches while allowing for polymorphisms and errors in sequencing. Each BLAST search generated a cluster of related sequences based on direct overlap with the query

“seed” sequence. A second level of clustering was performed to merge closely related clusters and to eliminate redundancy resulting from the fact that similar clusters are generated if the clusters contain more than one seed (*i.e.*, sequences from the Plus EST group). The resulting “super clusters” were discarded if they grew in size to 200 or more ESTs, since these probably represented repetitive elements that were not removed by the initial preprocessing of the seeds, or highly expressed genes such as those for ribosomal proteins. Superclusters were merged if they shared at least one third of their sequences.

The BLAST searches gave rise to a total of 49,154 clusters. In the first super clustering stage, 18,665 clusters grew beyond the limit of 200 clones. The remainder was reduced to a total of 30,489 super clusters. This number was reduced to 29,501 after adjacent clusters were merged. Resulting super clusters were analyzed to determine the tissue source of each EST clone contained within it and this expression profile was used to classify the superclusters into four groups: Type 1 – this supercluster contains EST clones found in the Plus group only, with no expression in the Minus or Other/Neutral group libraries; Type 2 – EST clones in the supercluster are found in the Plus and Other/Neutral group libraries, with no expression in the Minus group; Type 3 – super cluster EST clones found in all groups, but the number of ESTs in the Plus group is higher than in either of the Minus or Other/Neutral groups; Type 4 – super cluster EST clones found in all groups, but the number in the Plus group is higher than in the Minus group with expression in the Other/Neutral group non relevant. Sequences derived from the Plus library group that were placed in Types 1, 2 and 3 superclusters resulted in 20,487 polynucleotide sequences. The electronic subtraction procedures identified these sequences as having significant differential expression in lung tissue.

EXAMPLE 2

ANALYSIS OF CDNA EXPRESSION USING MICROARRAY TECHNOLOGY

2208 of the clones identified from the lung electronic subtraction procedure were evaluated for overexpression in specific tumor tissues by microarray analysis. Using this approach, cDNA sequences are PCR amplified and their mRNA expression profiles in tumor and normal tissues are examined using cDNA microarray

technology essentially as described (Shena, M. *et al.*, 1995 Science 270:467-70). In brief, the 2208 clones were arrayed onto glass slides as multiple replicas, with each location corresponding to a unique cDNA clone (as many as 5500 clones can be arrayed on a single slide or chip). Each chip was hybridized with a pair of cDNA probes that were fluorescence-labeled with Cy3 and Cy5, respectively. Typically, 1µg of polyA⁺ RNA was used to generate each cDNA probe. Since one cDNA probe is generated from tumor tissue RNA and the other is generated from normal tissue RNA, sequences that are differentially overexpressed in tumor tissue will generate a stronger signal from the tumor specific probe than the normal tissue probe, thus allowing the identification of those sequences that exhibit elevated expression in tumor versus normal tissue.

After hybridization, the chips were scanned and the fluorescence intensity recorded for both Cy3 and Cy5 channels. There were multiple built-in quality control steps. First, the probe quality was monitored using a panel of 18 ubiquitously expressed genes. Secondly, the control plate also had yeast DNA fragments of which complementary RNA was spiked into the probe synthesis for measuring the quality of the probe and the sensitivity of the analysis. Currently, the technology offers a sensitivity of 1 in 100,000 copies of mRNA. Finally, the reproducibility of this technology was ensured by including duplicated control cDNA elements at different locations. Further validation of the process was indicated in that several differentially expressed genes were identified multiple times in the study, and the expression profiles for these genes are very comparable. The clones were arrayed on Lung Chip 6.

Of those analyzed by microarray, 781 sequences met the criteria of having at least 2-fold overexpression in lung tumor tissue compared to normal tissues. Of these 781 clones, 459 were found to meet the additional criteria of having a mean normal tissue expression value less than or equal to 0.2. These 459 clones were then analyzed visually and certain clones with favorable expression profiles (e.g., high expression in tumors with little or no expression in normal tissues) were sequenced and searched against public sequence databases to facilitate identification of extended sequence for the clones.

SEQ ID NO:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 32 and 34 represent a subset of those 459 clones that met the above criteria of being at least 2-

fold overexpressed in tumor versus normal tissues and having a mean normal tissue expression of less than or equal to 0.2. Additional information about these sequences is provided in Table 2 below.

Table 2

SEQ ID NO:	SEQ ID NO: from 60/207,485	Clone Name:	Clone ID #	MICROARRAY ANALYSIS (Lung Chip #)	MICROARRAY RATIO (Lung Tumor:Normal Tissue)
9	4538	L1027C	55571	6	2.94
5	4978	L1037C	58267	6	2.61
7	1796	L1038C	58245	6	3.5
3	7264	L1039C	58269	6	2.81
1	2337	L1040C	55964	6	5.07
15	1548/4619	L1041C	58346	6	2.33
25	15127	n/a	56016	6	> 2
27	3816	n/a	55987	6	> 2
29	2046	n/a	55956	6	> 2
31	1912	n/a	55952	6	> 2
32	2064	n/a	55957	6	> 2
34	1502/3852	n/a	55559	6	> 2
11	2814	n/a	55978	6	> 2
13	3478	n/a	55980	6	> 2
17	553	n/a	55561	6	> 2
19	3275	n/a	55984	6	> 2
21	2809	n/a	58261	6	> 2
23	1677	n/a	58348	6	> 2

Each of the sequences was then used as a query to search the public databases in order to facilitate identification of extended sequences for these clones. Extended sequence information for the above sequences, obtained by searching public sequence databases, is set forth in SEQ ID NO:2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 33, and 35, respectively.

EXAMPLE 3

QUANTITATIVE REAL-TIME RT-PCR ANALYSIS

Briefly, quantitation of PCR product relies on the few cycles where the amount of DNA amplifies logarithmically from barely above the background to the plateau. Using continuous fluorescence monitoring, the threshold cycle number where DNA amplifies logarithmically is easily determined in each PCR reaction. There are two fluorescence detecting systems. One is based upon a double-strand DNA specific binding dye SYBR Green I dye. The other uses TaqMan probe containing a Reporter dye at the 5' end (FAM) and a Quencher dye at the 3' end (TAMRA) (Perkin Elmer/Applied Biosystems Division, Foster City, CA). Target-specific PCR amplification results in cleavage and release of the Reporter dye from the Quencher-containing probe by the nuclease activity of AmpliTaq Gold™ (Perkin Elmer/Applied Biosystems Division, Foster City, CA). Thus, fluorescence signal generated from released reporter dye is proportional to the amount of PCR product. Both detection methods have been found to generate comparable results. To compare the relative level of gene expression in multiple tissue samples, a panel of cDNAs is constructed using RNA from tissues and/or cell lines, and Real-Time PCR is performed using gene specific primers to quantify the copy number in each cDNA sample. Each cDNA sample is generally performed in duplicate and each reaction repeated in duplicated plates. The final Real-time PCR result is typically reported as an average of copy number of a gene of interest normalized against internal actin number in each cDNA sample. Real-time PCR reactions may be performed on a GeneAmp 5700 Detector using SYBR Green I dye or an ABI PRISM 7700 Detector using the TaqMan probe (Perkin Elmer/Applied Biosystems Division, Foster City, CA).

Using this approach, Real Time PCR[®] profiles were generated for L1027, L1037, L1038, L1039, L1040 and L1041, and are provided in Table 3.

Table 3

SEQ ID NO:	CLONE NAME	REAL TIME PROFILE
9	L1027C	Real Time PCR shows over-expression in small cell lung carcinoma as well as in bone marrow. Expression is also observed for multiple normal tissue.
5	L1037C	Real Time PCR shows over-expression in small cell lung carcinoma as well as in bone marrow and lymph node. Expression is also observed for multiple normal tissue.
7	L1038C	Real Time PCR shows over-expression in small cell lung carcinoma as well as in brain, pituitary gland and adrenal gland. Expression is also observed for multiple normal tissue.
3	L1039C	Real Time PCR shows over-expression in small cell lung carcinoma as well as in lymph node. Expression is also observed for multiple normal tissue.
1	L1040C	Real Time PCR shows over-expression in small cell lung carcinoma as well as in brain, pituitary gland and adrenal gland. Expression is also observed for multiple normal tissue.
15	L0141C	Real Time PCR shows over-expression in small cell lung carcinoma as well as in adrenal gland, bone marrow and thymus. Expression is also observed for multiple normal tissue.

5

EXAMPLE 4

CLONING OF FULL-LENGTH CDNA SEQUENCES AND ORF FOR L1027C

cDNA sequences encoding the full-length sequence for L1027C were isolated by screening a small cell primary tumor full length cloning library with a radioactively labeled probe of the original isolate sequence (SEQ ID NO:9). In order to determine the transcript size of the gene, a multiple tissue Northern blot was probed with the radioactively labelled original isolate sequence, SEQ ID NO:9. The Northern blot included 1µg of small cell primary tumor polyA⁺ RNA. Visual analysis of the

exposed film revealed a single transcript of approximately 2.5 kb. Approximately 500,000 clones from the full-length cloning library were screened and four clones were obtained from this library. The inserts were sequenced and yielded DNA nucleotide molecules of about 2.32 and 2.37 kb. These sequences are provided in SEQ ID NO:93 and 94, respectively. Both of these sequences contain the same single ORF of 450 bp (SEQ ID NO:95), and encode a deduced amino acid sequence of 150 amino acid residues (SEQ ID NO:96). These sequences were searched against the Genbank nonredundant and GeneSeq DNA databases and showed no hits.

10

EXAMPLE 5

ANALYSIS OF CDNA EXPRESSION USING MICROARRAY TECHNOLOGY

An additional 5054 of the resulting clones obtained from the lung electronic subtraction of Example 1 were probed by microarray chip technology to further characterize the expression of these clones. The microarray analysis was carried out as provided in Example 2. The clones were arrayed on Lung Chip 7. CorixArray analysis was performed on the microarray results to compare expression in lung tumors and in normal tissues. Clones were selected based on two criteria: 2-fold overexpression in lung tumors when compared to non-lung tissue and a mean expression level of less than 0.2 in these same non-lung tissues. Of those analyzed, 2372 clones met the criteria.

20

Microarray analysis for five of these clones is presented in Table 4:

Table 4

SEQ ID NO:	SEQ ID NO: from 60/207,485	Clone Name:	Clone ID #	MICROARRAY ANALYSIS (Lung Chip #)	MICROARRAY RATIO (Lung Tumor:Normal Tissue)
42	18618	L1053C	63575	7	13.5
43	14788	L1054C	63582	7	5.29
44	7744	L1055C	63598	7	15.25
45	4257	L1056C	64963	7	9.31
46	20087	L1058C	64988	7	5.66

5

EXAMPLE 6

QUANTITATIVE REAL-TIME PCR ANALYSIS

170 of the 2372 clones of Example 4 were further analyzed by visual analysis based on high expression in tumors and little or no expression in normal tissues. Seven clones were selected for Real-time PCR analysis. The Real-time PCR was carried out as disclosed in Example 3. The Real-time PCR profiles of these seven clones are presented in Table 5. The sequences of these seven clones are provided in SEQ ID NO:42-48, respectively.

Table 5

SEQ ID NO:	CLONE NAME	CLONE ID #	REAL TIME PROFILE
42	L1053C	63575	Real Time PCR shows over-expression in small cell lung carcinoma as well as in pituitary. Expression is also observed for multiple normal tissues.
43	L1054C	63582	Real Time PCR shows over-expression in small cell lung carcinoma as well as in pituitary, brain and spinal cord. Expression is also observed for adrenal and pancreas.
44	L1055C	63598	Real Time PCR shows over-expression in small

			cell lung carcinoma as well as in pituitary and brain. Expression is also observed for multiple normal tissues.
45	L1056C	64963	Real Time PCR shows over-expression in one small cell lung carcinoma sample. No expression is otherwise observed.
46	L1058C	64988	Real Time PCR shows over-expression in small cell lung carcinoma. Low level expression is also observed for adrenal gland, pancreas, and bone marrow.
47	n/a	63485	Real Time PCR shows over-expression in metastatic tumor as well as low level expression in multiple normal tissues.
48	n/a	65010	Real Time PCR shows low expression in one lung sample. No expression is otherwise observed.

Each of the sequences was then used as a query to search the public databases in order to facilitate identification of extended sequences for these clones. SEQ ID NO:42, 43 and 45 matched to known genes in Genbank, and these results are presented in Table 6. The full-length cDNA sequences of the known genes are disclosed in SEQ ID NO:49, 50 and 52, respectively. The deduced amino acid sequences encoded by SEQ ID NO:49 and 50 are also provided as SEQ ID NO:56 and 57, respectively. SEQ ID NO:44 and 46-48 were found to be novel with respect to known genes, but matched to public EST sequences. The sequences of SEQ ID NO:44 and 46-48 were aligned with the matching EST sequences in order to obtain extended sequence data. These extended sequences are provided in SEQ ID NO:51 and 53-55, respectively.

Table 6

SEQ ID NO:	CLONE NAME	GENBANK DESCRIPTION
42	L1053C	Insulinoma-associated 1
43	L1054C	KIAA0535
45	L1056C	Human DAZ mRNA 3' UTR

EXAMPLE 7

CLONING OF cDNA ENCODING FULL-LENGTH L1058C

The cDNA sequence encoding full-length L1058C was isolated by screening a small cell primary tumor full length cloning library with a radioactively labeled probe of the original isolate sequence (SEQ ID NO:46). In order to determine the transcript size of the gene, a multiple tissue Northern blot was probed with the radioactively labelled original isolate sequence, SEQ ID NO:46. The Northern blot included 1µg of small cell primary tumor, carcinoid metastasis and small cell (tumor) cell line polyA+ RNA. Visual analysis of the exposed film revealed a single transcript of approximately 2.5 kb. Approximately 500,000 clones from the full-length cloning library were screened and one clone was obtained from this library. The insert was sequenced and yields a 2165 bp DNA nucleotide molecule. The full-length sequence is provided in SEQ ID NO:58. The full-length sequence is predicted to have two ORFs. A first ORF (SEQ ID NO:59) is predicted to encode a polypeptide having 392 amino acid residues (SEQ ID NO:61), and the second ORF (SEQ ID NO:60) is predicted to encode a polypeptide of 363 amino acid residues (SEQ ID NO:62) but does not show the starting methionine. This 2165 bp DNA was searched against the Genbank nonredundant and GeneSeq DNA databases and showed no hits.

EXAMPLE 8

ANALYSIS OF CDNA EXPRESSION USING MICROARRAY TECHNOLOGY

An additional 3453 of the resulting clones obtained from the lung electronic subtraction of Example 1 were probed by microarray chip technology to further characterize the expression of these clones. The microarray analysis was carried out as provided in Example 2. The clones were arrayed on Lung Chip 8. CorixArray analysis was performed on the microarray results to compare expression in lung tumors and in normal tissues. Clones were selected based on two criteria: 2-fold overexpression in lung tumors when compared to non-lung tissue and a mean expression level of less than 0.2 in these same non-lung tissues. Of those analyzed, 557 clones met the criteria.

300 of the 557 clones were visually analyzed for overexpression in tumor versus normal tissue. Twenty-eight clones showing overexpression in tumor versus normal tissue were then sequenced. These DNA sequences are provided in SEQ ID NO:63-92, respectively. The microarray analysis for these 28 clones is presented in

5 Table 7.

Table 7

SEQ ID NO:	CLONE ID #	RATIO	MEDIAN SIGNAL 1	MEDIAN SIGNAL 2
63	72761	2.22	0.154	0.07
64	72762	2.33	0.105	0.045
65	72763	2.41	0.233	0.097
66	72764	2.72	0.199	0.073
67	72765	2.62	0.158	0.06
68	72766	2.84	0.149	0.053
69	72772	2.25	0.109	0.049
70	72775	2.36	0.103	0.044
71	72776	2.34	0.146	0.062
72	72779	2.25	0.22	0.098
73	72781	2.51	0.149	0.059
74	72784	2.35	0.212	0.09
75	72788	2.85	0.152	0.053
76	72789	2.69	0.196	0.073
77	72790	2.46	0.181	0.074
78	72791	2.39	0.143	0.06
79	72792	2.43	0.197	0.081
80	72794	3.04	0.258	0.085
81	72795	2.37	0.143	0.06
82	72797	2.96	0.233	0.079
83	72798	2.82	0.218	0.077
84	72804	2.33	0.14	0.06
85	72805	2.33	0.102	0.043
86	72806	2.32	0.121	0.052
87	72807	3.02	0.117	0.039
88	72808	2.74	0.109	0.04
89	72809	2.26	0.126	0.056
90	72811	2.92	0.151	0.052
91	72813 (L1080C)	2.66	0.138	0.052

Each of the sequences was then used as a query to search the public sequence databases to identify novel and known genes. Results of this search are provided in Table 8.

5

Table 8

SEQ ID NO:	GEN BANK ACC #	GENESEQ	DESCRIPTION
63	AC004590		Chromosome 17
64	Z78409	T62661	transcription factor E2F5
65	S45828	Z86797; A09328	cDNA DKFZp564L2416; nekl=serine/threonine-and tyrosine-specific protein kinase [mice, erythroleukemia cells]
66			Novel
67	AL136169		Chromosome Xq26.1-27.1
68	AC011742 AK021426		Chromosome 2, Homo sapiens cDNA FLJ11364 fis. clone HEMBA 1000264.
69	NM 005414	Q03742	SKI-like (SKIL)
70	NM 002335	V85551	low density lipoprotein receptor- related protein 5
71	XM_004587 AB000520		Homo sapiens adaptor protein with pleckstrin homology and src homology 2 domains (APS), mRNA. Homo sapiens mRNA for APS, complete cds.
72	AK024119		cDNA FLJ14057 fis. clone HEMBB1000337.
73	U86338		Mus musculus zinc finger protein Png-1 (Png-1)
74			Novel
75			Novel
76	NM_002271	C03734	Homo sapiens karyopherin (importin) beta 3 (KPNB3) mRNA
77	NM_001401	T48669;	Homo sapiens endothelial

		T44104	differentiation, lysophosphatidic acid G-protein-coupled receptor, 2(EDG2), mRNA.
78	U40583		Human alpha / neuronal nicotinic acetylcholine receptor mRNA, complete cds.
79		Z15509	Novel
80	Z59860	V34162	H. sapiens CpG island DNA genomic MseI fragment, clone 178c7, reverse read cpg178c7.rta.
81			Novel
82	Z59860	HNGIT2 2	DNA genomic MseI fragment, clone 178c7
83	XM-004477	Q72451	Homo sapiens glutamate-cysteine ligase, catalytic subunit (GCLC), mRNA.
84		Z16421	Novel
85			Novel
86	AC022013	V52850	Chromosome 3
87			Novel
88	AL354993	Z91766	Chromosome 20q13.2-13. Continas a peptidylprolyl isomerase A (cyclophilin A) pseudogene, the gene for OVC10-2, ESTs, STSs and GSSs, complete sequence
89	AC005021		Chromosome 7q21-q22, complete sequence.
90	AK023904		cDNA FLJ13842 fis, clone THYRO1000793.

EXAMPLE 9

QUANTITATIVE REAL-TIME PCR ANALYSIS

- 5 One of the clones of Example 7, clone L1080C, was further selected for Real-time PCR analysis. The Real-time PCR was carried out as disclosed in Example 3. The Real-time PCR shows over-expression in small cell lung carcinoma as well as in

brain and pituitary. Expression was also observed in thyroid, adrenal and salivary glands.

EXAMPLE 10

5 IDENTIFYING FULL-LENGTH cDNA SEQUENCE ENCODING L1080C

The cDNA sequence encoding full-length L1080C was predicted by using a partial sequence as a query to search the public sequence databases to obtain extended sequence. The query resulted in the identification of a full-length cDNA sequence for L1080C (SEQ ID NO:91). The deduced amino acid sequence encoded by 10 the full-length cDNA sequence is provided in SEQ ID NO:92.

EXAMPLE 11

PEPTIDE PRIMING OF T-HELPER LINES

Generation of CD4⁺ T helper lines and identification of peptide epitopes 15 derived from tumor-specific antigens that are capable of being recognized by CD4⁺ T cells in the context of HLA class II molecules, is carried out as follows:

Fifteen-mer peptides overlapping by 10 amino acids, derived from a tumor-specific antigen, are generated using standard procedures. Dendritic cells (DC) are derived from PBMC of a normal donor using GM-CSF and IL-4 by standard 20 protocols. CD4⁺ T cells are generated from the same donor as the DC using MACS beads (Miltenyi Biotec, Auburn, CA) and negative selection. DC are pulsed overnight with pools of the 15-mer peptides, with each peptide at a final concentration of 0.25 µg/ml. Pulsed DC are washed and plated at 1×10^4 cells/well of 96-well V-bottom plates and purified CD4⁺ T cells are added at 1×10^5 /well. Cultures are supplemented 25 with 60 ng/ml IL-6 and 10 ng/ml IL-12 and incubated at 37°C. Cultures are restimulated as above on a weekly basis using DC generated and pulsed as above as antigen presenting cells, supplemented with 5 ng/ml IL-7 and 10 U/ml IL-2. Following 4 *in vitro* stimulation cycles, resulting CD4⁺ T cell lines (each line corresponding to one well) are tested for specific proliferation and cytokine production in response to the 30 stimulating pools of peptide with an irrelevant pool of peptides used as a control.

EXAMPLE 12

GENERATION OF TUMOR-SPECIFIC CTL LINES USING IN VITRO WHOLE-GENE PRIMING

Using *in vitro* whole-gene priming with tumor antigen-vaccinia infected

5 DC (see, for example, Yee et al, *The Journal of Immunology*, 157(9):4079-86, 1996), human CTL lines are derived that specifically recognize autologous fibroblasts transduced with a specific tumor antigen, as determined by interferon- γ ELISPOT analysis. Specifically, dendritic cells (DC) are differentiated from monocyte cultures derived from PBMC of normal human donors by growing for five days in RPMI

10 medium containing 10% human serum, 50 ng/ml human GM-CSF and 30 ng/ml human IL-4. Following culture, DC are infected overnight with tumor antigen-recombinant vaccinia virus at a multiplicity of infection (M.O.I) of five, and matured overnight by the addition of 3 μ g/ml CD40 ligand. Virus is then inactivated by UV irradiation. CD8+ T cells are isolated using a magnetic bead system, and priming cultures are

15 initiated using standard culture techniques. Cultures are restimulated every 7-10 days using autologous primary fibroblasts retrovirally transduced with previously identified tumor antigens. Following four stimulation cycles, CD8+ T cell lines are identified that specifically produce interferon- γ when stimulated with tumor antigen-transduced autologous fibroblasts. Using a panel of HLA-mismatched B-LCL lines transduced

20 with a vector expressing a tumor antigen, and measuring interferon- γ production by the CTL lines in an ELISPOT assay, the HLA restriction of the CTL lines is determined.

EXAMPLE 13

GENERATION AND CHARACTERIZATION OF ANTI-TUMOR ANTIGEN MONOCLONAL

25

ANTIBODIES

Mouse monoclonal antibodies are raised against *E. coli* derived tumor antigen proteins as follows: Mice are immunized with Complete Freund's Adjuvant (CFA) containing 50 μ g recombinant tumor protein, followed by a subsequent intraperitoneal boost with Incomplete Freund's Adjuvant (IFA) containing 10 μ g

30 recombinant protein. Three days prior to removal of the spleens, the mice are immunized intravenously with approximately 50 μ g of soluble recombinant protein. The

spleen of a mouse with a positive titer to the tumor antigen is removed, and a single-cell suspension made and used for fusion to SP2/O myeloma cells to generate B cell hybridomas. The supernatants from the hybrid clones are tested by ELISA for specificity to recombinant tumor protein, and epitope mapped using peptides that span the entire tumor protein sequence. The mAbs are also tested by flow cytometry for their ability to detect tumor protein on the surface of cells stably transfected with the cDNA encoding the tumor protein.

EXAMPLE 14

SYNTHESIS OF POLYPEPTIDES

Polypeptides are synthesized on a Perkin Elmer/Applied Biosystems Division 430A peptide synthesizer using Fmoc chemistry with HPTU (O-Benzotriazole-N,N,N',N'-tetramethyluronium hexafluorophosphate) activation. A Gly-Cys-Gly sequence is attached to the amino terminus of the peptide to provide a method of conjugation, binding to an immobilized surface, or labeling of the peptide. Cleavage of the peptides from the solid support is carried out using the following cleavage mixture: trifluoroacetic acid:ethanedithiol:thioanisole:water:phenol (40:1:2:2:3). After cleaving for 2 hours, the peptides are precipitated in cold methyl-t-butyl-ether. The peptide pellets are then dissolved in water containing 0.1% trifluoroacetic acid (TFA) and lyophilized prior to purification by C18 reverse phase HPLC. A gradient of 0%-60% acetonitrile (containing 0.1% TFA) in water (containing 0.1% TFA) is used to elute the peptides. Following lyophilization of the pure fractions, the peptides are characterized using electrospray or other types of mass spectrometry and by amino acid analysis.

From the foregoing it will be appreciated that, although specific embodiments of the invention have been described herein for purposes of illustration, various modifications may be made without deviating from the spirit and scope of the invention. Accordingly, the invention is not limited except as by the appended claims.

CLAIMS

What is Claimed:

1. An isolated polynucleotide comprising a sequence selected from the group consisting of:

(a) sequences provided in SEQ ID NO:1-3, 5, 7, 9, 11-19, 25-35, 44, 46, 47, 48, 53-55, 58-60, 66, 74, 75, 79, 81, 84, 85, 87, 93, 94 and 95;

(b) complements of the sequences provided in SEQ ID NO:1-3, 5, 7, 9, 11-19, 25-35, 44, 46, 47, 48, 53-55, 58-60, 66, 74, 75, 79, 81, 84, 85, 87, 93, 94 and 95;

(c) sequences consisting of at least 20 contiguous residues of a sequence provided in SEQ ID NO:1-3, 5, 7, 9, 11-19, 25-35, 44, 46, 47, 48, 53-55, 58-60, 66, 74, 75, 79, 81, 84, 85, 87, 93, 94 and 95;

(d) sequences that hybridize to a sequence provided in SEQ ID NO:1-3, 5, 7, 9, 11-19, 25-35, 44, 46, 47, 48, 53-55, 58-60, 66, 74, 75, 79, 81, 84, 85, 87, 93, 94 and 95, under highly stringent conditions;

(e) sequences having at least 75% identity to a sequence of SEQ ID NO:1-3, 5, 7, 9, 11-19, 25-35, 44, 46, 47, 48, 53-55, 58-60, 66, 74, 75, 79, 81, 84, 85, 87, 93, 94 and 95;

(f) sequences having at least 90% identity to a sequence of SEQ ID NO:1-3, 5, 7, 9, 11-19, 25-35, 44, 46, 47, 48, 53-55, 58-60, 66, 74, 75, 79, 81, 84, 85, 87, 93, 94 and 95; and

(g) degenerate variants of a sequence provided in SEQ ID NO:1-3, 5, 7, 9, 11-19, 25-35, 44, 46, 47, 48, 53-55, 58-60, 66, 74, 75, 79, 81, 84, 85, 87, 93, 94 and 95.

2. An isolated polypeptide comprising an amino acid sequence selected from the group consisting of:

(a) sequences having an amino acid sequence of any one of SEQ ID NO:61, 62 and 96;

- (b) sequences encoded by a polynucleotide of claim 1;
 - (c) sequences having at least 70% identity to a sequence encoded by a polynucleotide of claim 1; and
 - (d) sequences having at least 90% identity to a sequence encoded by a polynucleotide of claim 1.
3. An expression vector comprising a polynucleotide of claim 1 operably linked to an expression control sequence.
4. A host cell transformed or transfected with an expression vector according to claim 3.
5. An isolated antibody, or antigen-binding fragment thereof, that specifically binds to a polypeptide of claim 2.
6. A method for detecting the presence of a cancer in a patient, comprising the steps of:
- (a) obtaining a biological sample from the patient;
 - (b) contacting the biological sample with a binding agent that binds to a polypeptide of claim 2;
 - (c) detecting in the sample an amount of polypeptide that binds to the binding agent; and
 - (d) comparing the amount of polypeptide to a predetermined cut-off value and therefrom determining the presence of a cancer in the patient.
7. A fusion protein comprising at least one polypeptide according to claim 2.
8. An oligonucleotide that hybridizes to a sequence recited in SEQ ID NO:1-3, 5, 7, 9, 11-19, 25-35, 44, 46, 47, 48, 53-55, 58-60, 66, 74, 75, 79, 81, 84, 85, 87, 93, 94 and 95 under highly stringent conditions.

9. A method for stimulating and/or expanding T cells specific for a tumor protein, comprising contacting T cells with at least one component selected from the group consisting of:

(a) polypeptides according to claim 2;
(b) polynucleotides according to claim 1; and
(c) polynucleotides having a nucleotide sequence of any one of SEQ ID NO:4, 6, 8, 10, 20-24, 42, 43, 45, 49-52, 63-65, 67-73, 76-78, 80, 82, 83, 86 and 88-91;

(d) antigen-presenting cells that express a polynucleotide according to claim 1,

under conditions and for a time sufficient to permit the stimulation and/or expansion of T cells.

10. An isolated T cell population, comprising T cells prepared according to the method of claim 9.

11. A composition comprising a first component selected from the group consisting of physiologically acceptable carriers and immunostimulants, and a second component selected from the group consisting of:

(a) polypeptides according to claim 2;
(b) polynucleotides according to claim 1;
(c) polynucleotides having a nucleotide sequence of any one of SEQ ID NO:4, 6, 8, 10, 20-24, 42, 43, 45, 49-52, 63-65, 67-73, 76-78, 80, 82, 83, 86 and 88-91;

(d) antibodies according to claim 5;
(e) fusion proteins according to claim 7;
(f) T cell populations according to claim 10; and
(g) antigen presenting cells that express a polypeptide according to claim 2.

12. A method for stimulating an immune response in a patient, comprising administering to the patient a composition of claim 11.

13. A method for the treatment of a lung cancer in a patient, comprising administering to the patient a composition of claim 11.

14. A method for determining the presence of a cancer in a patient, comprising the steps of:

- (a) obtaining a biological sample from the patient;
- (b) contacting the biological sample with an oligonucleotide according to claim 8;
- (c) detecting in the sample an amount of a polynucleotide that hybridizes to the oligonucleotide; and
- (d) compare the amount of polynucleotide that hybridizes to the oligonucleotide to a predetermined cut-off value, and therefrom determining the presence of the cancer in the patient.

15. A diagnostic kit comprising at least one oligonucleotide according to claim 8.

16. A diagnostic kit comprising at least one antibody according to claim 5 and a detection reagent, wherein the detection reagent comprises a reporter group.

17. A method for the treatment of lung cancer in a patient, comprising the steps of:

- (a) incubating CD4+ and/or CD8+ T cells isolated from a patient with at least one component selected from the group consisting of: (i) polypeptides according to claim 2; (ii) polynucleotides according to claim 1; and (iii) antigen presenting cells that express a polypeptide of claim 2, such that T cell proliferate;

(b) administering to the patient an effective amount of the proliferated T cells,
and thereby inhibiting the development of a cancer in the patient.

SEQUENCE LISTING

<110> Corixa Corporation
 Harlocker, Susan L.
 Wang, Tongtong
 Bangur, Chaitanya S.
 Klee, Jennifer
 Switzer, Anne

<120> COMPOSITIONS AND METHODS FOR THE THERAPY
 AND DIAGNOSIS OF LUNG CANCER.

<130> 210121.502PC

<140> PCT

<141> 2001-05-25

<160> 96

<210> 1

<211> 644

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<222> (1)...(644)

<223> n = A, T, C or G

<400> 1

```

ttactcctct agagggaaag catgacacog aacactaagc acacagcttt ttgttgtttt 60
ggttttttct cccgcaaatc ttaaagtgat tcccatgacc ttggccaagg acactttetta 120
aagattaatg actggcactg acattgcccc aggcggggca ctctccacac tggctctcag 180
ttccagccca tgccctgggc tcagtcactt ctattccacc ctctgagact ccattgggtg 240
cacacaaggt gtctctcttg ctttgatttt gagaatcccc tatitttctact tccagatctg 300
tcagctgccca tggaggaata atagaaaaac agaaatcgct gttagaggag atttctaaaa 360
cttcccttgt gtgcccatag ttgtagtttt gggttctggc aggtggaaca ccttgaaacc 420
tggaatcatt ctatgagaat acagttcaga ctttgagacc tccagcccat actaactgtc 480
atgaagcttg actctctgtc ataatgcagc catctggag gaaattggca ttctgtctta 540
gatgntggc agggtcgcgc tcagctttgc tttctacact aaattacata gcattaatto 600
aagnattggt ttccaatttc ccatccctga tttccagctt tctt
644

```

<210> 2

<211> 1115

<212> DNA

<213> Homo sapiens

<400> 2

```

gtaggaaggt acagtaaatg gtagttcatt cttacttaca cacatagcta atcttttttt 60
tttcaactgg aattatgttg aatgtttcat ttgacaaaa aagtagacta gaaggtatgt 120
yctttaagtt gtcttgcctc cattatataa gaagaagaca ggtgagagga agagcagaaa 180
ctctgagactg gctgatgttc agagcactta ctctctaga gggaaagcat gacaccgaac 240
actaagcaca cagctttttg ttgttttggt tttttctccc gcaaatctta aagtgtatcc 300
ctgacccttg gccaaagaca cttcttaag attaatgact ggcactgaca ttgcccagg 360
cgggccactc ctcacactgg ctctcagttc ccagccatgc ctggggctca gtcacttcta 420

```

ttccaccctc	tgagactcca	ttggtgtcac	acaaggtgtc	ttcttggctt	tgattttgag	480
aatccctcat	tttcaactcc	agatctgtca	gctgccatgg	aggaataata	gaaaaccaga	540
aatgcgtgta	gaggggagatt	tctaaaaactt	cccttgtgtc	gcccatagatt	gtagttttgg	600
gttctggcag	gtggaaacacc	ctgaaacctg	gaatcaattct	atgagaataac	agttcagact	660
ttgcagactc	cagcccatatc	taactgtcat	gaagcttgac	ttcttgtcat	aatgcagcca	720
tcttggagga	aattggcccat	ttctgcttag	atggttggca	gggtcgcgct	cagctttgtg	780
ttctacaacta	attacatagc	attattcaag	tattgttttc	catttccocat	cctgtatttc	840
cagcttctta	aagctgactg	ttcttgcagg	ggccaacttgc	ttctctcaga	gtacaaaagt	900
aaggcgcttc	cttactaact	gcagggtctc	tctattacac	ctcaacatc	acaotttgcg	960
gctactgttt	gtactgtcta	cagtagaatt	tccttatctt	gctcctggta	gtgcaattaca	1020
ggcaagcatg	aaatgtaaa	tatttattta	aataaaaaga	aaacctctaa	attggtaatt	1080
gaawwammwm	mmwrwammw	tatagtttgt	gacat			1115

<210> 3

<211> 540

<212> DNA

<213> Homo sapiens

<400> 3

gggcagaaat	tcggccgagg	cctgcaaacg	agaaggctgt	ggatttgatt	attgtacgaa	60
gtgtctctgt	aattatcata	ctactaaga	ctgttcagat	ggcaagctcc	tcaaaagccag	120
ttgtaaaata	ggctccctgc	ctgtgtacaa	gaaaagcaaa	aagaattttac	gaagatttgg	180
atctcttatt	aaatcaattg	ttactgtaca	tgaattgtta	ttagaataatg	taggttttta	240
acttaaaaaa	aattgtattg	tgattttcaa	ttttatgttg	aaatcggtgt	tatctcctga	300
ggtttttttc	ccccagaaag	ataaagagga	tagacaacct	ottaaaatat	ttttacaatt	360
taattgagaaa	aagtttaaaa	ttctcaatc	aaatcaaaaca	atttaaatat	tttaagaaaa	420
aaggaaaaagt	agatagtgat	actgagggtg	aaaaaaaatt	gattcaattt	tatggttaag	480
gaaacccatg	caattttacc	tagacagtct	taaatatgtc	tggtttttcca	tctgttagca	540

<210> 4

<211> 2076

<212> DNA

<213> Homo sapiens

<400> 4

agggtgtctca	gctgcccccg	gagcgggttc	ttcaactgtg	gcagacacca	cctcggttgg	60
cagtgcgcgg	cgccccctgca	gctgcgcctc	acggcocaacc	cgtcgtctct	cgctgcgcag	120
ccccagcgcca	gtgacagccg	ccggcgccgc	tcgacccctg	gatagttgta	aagaagaaag	180
ttctaaccctt	tctgtcaaaa	tgaagtgtga	ttttaattgt	aaacatgttc	attcccgact	240
taaacttgta	aaactgtgatg	acattggaag	actagtttcc	tacacccctg	catatctgtga	300
aggttctcgt	aaagactgca	ttaaagacta	tgaaggctgt	tcagtgtattg	ggtcaccgat	360
tgtgagccct	aggatgttac	aacttgaaac	tgaagcaag	cgtctgcata	acaagaaaaa	420
tcacacatgt	caacagacac	ttaatgtatc	aaatgaaata	gaagcactag	agacccagtag	480
actttatgaa	gacagtggct	attcctcatt	ttctotacaa	agtggcctca	gtgaacatga	540
agaaggttagc	ctcctggagg	agaatttcgg	tgacagtcta	caatcctgcc	tgctacaaat	600
acaaagcccca	gaccaatata	ccaacaaaaa	cttgcctcca	gttcttcatt	ttgaaaaagt	660
ggtttgttca	acattaaaaa	agaatgcaaa	acgaaatctc	aaagttagatc	gggagatgct	720
gaaggaaatt	atagccagag	gaaatttttag	actgcagaat	ataattggca	gaaaaatggg	780
octagaatgt	gtagatatct	tcagcgaaat	ctttcgaaag	ggactcaagc	atgtcttagc	840
aactattttta	gcacaactca	gtgacatgga	cttaatcaat	gtgtctaagg	tgagcacaaac	900
tttgaagaag	atgataaagg	ggcattccag	ttgtacagat	aagcaataca	agcaataca	960
aagagttacc	gaaaacaaca	ataaattttc	acctcatgct	tcacacagag	aataattgtat	1020
ctgtcagaacc	ccactggtctt	ctgttcagaa	atcagcagcc	cagactttctc	tcaaaaaaga	1080
tgctcaaac	aagttatcca	atcaaggtga	tcagaaaggt	tctacttata	tgtagacacaa	1140
tgaattctct	gaggtgtcca	agacattgaa	aaagaacgaa	agcctcaag	cctgtattctg	1200
ctgtaattca	cctgcacaaat	atgattgtcta	tttacaacgg	gcaacctgca	aacgagaagg	1260
ctgtggattt	gattattgta	cgaagtgtct	ctgtaattat	catactacta	aagactgttc	1320
agatggcgaag	ctcctcaaaag	ccagttgttaa	aataggtccc	ctgcctgtga	caaagaaaaa	1380

caaaaagaat	ttacgaagat	tgtgatctct	tattaaatca	attgttactg	atcatgaatg	1440
ttagttagaa	aatgttaggt	tttaacttaa	aaaaaattgt	attgtgattt	tcaattttat	1500
gttgaatctg	gtgtagtatc	ctgaggtttt	tttcccccca	gaagataaag	aggatagaca	1560
acctctttaa	atatttttaac	aattttaatga	gaaaaagtgt	aaaattctca	atacaaatca	1620
aacaatttaa	atatttttaag	aaaaaaggaa	aagtagatag	tgatactgag	ggtaaaaaaa	1680
aaattgtatt	aatttttatgg	taaagggaac	ccatgcaatt	ttacctagac	agtcttaaat	1740
atgtctggtt	ttccatctgt	tagcatttca	gacattttat	gttctcttta	ctcaattgat	1800
accaacagaa	atatacaactt	ctggagtcta	ttaaatgtgt	tgctacottt	ctaaagcttt	1860
ttttcatgtg	gtgtatttcc	caagaagata	tcctttgtaa	aaacttgott	gttttccotta	1920
ttttcgaaat	ctgttttaaat	atttttgtat	acatgtaaat	atttctgtat	tttttatatg	1980
tcaagaata	tgtctcttgt	atgtacatat	aaaaataaat	tttgcctaat	aaaattgtaa	2040
gcttaaaaaa	aaaaaaaaaa	aactcgagac	tagtgc			2076

<210> 5

<211> 634

<212> DNA

<213> Homo sapiens

<400> 5

gggcagaatt	cggaacgagga	cttttctcca	gtgttgacct	taggggtgcag	ctggatgttt	60
ttaccctcag	cggcttttcgg	actgtacaga	tcctggaaag	acaaaagatc	ctggctaaact	120
gttctctctc	ctaccaggta	gacctgtttg	gtatagcaga	tttagcacat	ttactattgt	180
tcaaggaaaca	ctacacaggct	ttctgggatg	ggctctctctg	gaaacttagc	caaaatattt	240
ctgagctaaa	agatgtgtgaa	ttgtggaata	aattctttgt	gcgactctcg	aattgccaatg	300
atgagggcac	agtgctctgt	ctgggggagc	ttgcagcaga	aatgaatggg	gtttttgaca	360
ctacatttcca	aagtccacctg	aaacaagcct	tatggaaagt	agggaaagta	actagctctg	420
gggcttttgt	ctttcagtga	gctaggcaat	caagtctcac	agattgtctg	ctcagagcaa	480
tggtgttatt	gtggaacact	gaaactgtat	gtgtgttaat	ttaattttag	acacatttag	540
atgcactacc	attgtctgtc	tacttttttg	tacagggtata	ttttgacgtc	actgatattt	600
tttatacagt	gatatactta	ctcatggcct	tgct			634

<210> 6

<211> 3725

<212> DNA

<213> Homo sapiens

<400> 6

accgttaaat	ttgaaacttg	gcgggtaggg	gtgtgggctt	gagggtggcg	gtttgttagg	60
gagtcgtgtg	cgtgccttgg	tcgctcttgt	agctccggag	gcaggtttgg	gaagaaagcc	120
caggcggtct	gtggcccaga	ggaaggcctt	gcagcaggac	gaggacctga	gccaggaaatg	180
caggatggcg	gcggtgaaga	aggaaggggg	tgctctgagt	gaagccatgt	ccctggaggg	240
agatgaatgg	gaactgagta	aagaaaatgt	acaaccttta	aggcaaggcg	ggatcatgtc	300
cacgcttcag	ggagcactgg	cacaagaatc	tgctgttaac	aatactcttc	agcagcagaa	360
acgggcatctt	gaatatgaaa	ttcgatttta	cactggaaat	gcacctctgg	atgtttggga	420
taggtatatt	agctggacag	agcagaacta	tcctcaaggt	gggaaggaga	gtaatatgtc	480
aaacgttatta	gaaagagctg	tagaagcact	acaaggagaa	aaacgatatt	atagtgatcc	540
tcgatttctc	aatctctggc	ttaaatttag	gcgtttatgc	aatgagcctt	tggtatgtga	600
cagttactgt	cacaacCaag	ggattgggtg	ttcacttgct	gcgttctata	tctcatgggc	660
agaagaatat	gaagctagag	aaaactttag	gaaagcagct	gcgatatttc	aggaagggat	720
tcaacagag	gctgaaccac	tagaaagact	acagtcocag	caccgacaat	tccaagctcg	780
agtgctctgg	caaactctgt	tggcacttga	gaaagaagaa	gaggaggaaag	tttttgagtc	840
ttctgtacca	caacgagcaa	cactagctga	actaaagagc	aaagggaataa	agacagcaag	900
agctccaatc	atccgtgtag	gaggtgctct	caaggctcca	agccagaaaca	gaggactcca	960
aaatccattt	ctctaacaga	tgcaaaataa	tagtagaatt	actgtttttg	atgaaatagt	1020
tgtataggct	tcctagacag	agttgtctaa	gcctacagtc	cagccatgga	tagcaccctcc	1080
catgccacag	gccaaagaga	atgagctgca	agcaggccct	tggaacacag	gcaggtctctt	1140
ggaacacagg	cctcgtggca	atacagcttc	actgatagct	taccctcgag	tgctccccag	1200
tttcaactcca	tatgtgggaag	agactgcaca	acagccagtt	atgacaccat	gtaaaaatga	1260

```

acctagtata aaccacatcc taagcaccag aaagcctgga aaggaagaag gagatccctc 1320
acaaaggggtt cagagccatc agcaagcatc tgaggagaag aaagagaaga tgatgtattg 1380
taaggagaag atttatgcag gagtagggga attctccttt gaagaaattc ggggtgaagt 1440
tttccgggaag aaattaaaaag agcaaaaggga agccgagcta ttgaccattg cagagaagag 1500
agcagaatgt cagaacacaga ttgaagagat ggagaagaag ctaaaagaaa tccaaactac 1560
tcagcagaat agaacacagtg atcagcaaga agacagcatg cctacaaagg agacaactaa 1620
actgcaaat gtctccgagt ctcagaaaat accaggaatg actctatcca gttctgtttg 1680
tcaagttaac tgtttgcga gagaaacttc acttgcggag aacatttggc aggaacaacc 1740
tcattctaaa ggtccagtg taocctttctc catttttgat gagtttcttc tttaaaaaaa 1800
gaagaataaa agtctctctg cagatccccc acgagtttta gctcaacgaa gaccocctgc 1860
agttctcaaa acctcagaaa gcatcacctc aaatgaagat gtgtctccag atgtttgtga 1920
tgaatttaca ggaattgaac ccttgagcga ggatgccatt atcacaggct tcagaaatgt 1980
aacaatttgt cctaaccag aagacacttg tgaatttggc agagcagctc gttttgtatc 2040
caotcctttt catgagataa tgtccttgaa ggatctccct totgatccctg agagaactgt 2100
accggaagaa gatctagatg taaagacctc tgaggacacg cagacagctt gtggcactat 2160
ctacagtcag actctcagca toaagaagct gagcccaatt attgaagaca gtgctgaagc 2220
cacacactoc tctgcttctc ctggttcttc tgccctcggtt gcaagcacct cctccactca 2280
atgtcttcaa attctctgaga aactagaact tactaatgag acttcagaaa accctactca 2340
gtccactcag tgttcacagt atcgcagaca gctactgaag tccctacacg agttaagtgc 2400
ctctgcagag ttgtgtatag aagacagacc aatgcctaag ttggaaattg agaaggaat 2460
tgaattaggt aatgagatt actgcattaa acgagaatac ctaatatgtg aagattacaa 2520
gttattttgg gtggggccaa gaaactttgc agaattaaac gtaataaagg tatcttctca 2580
acotgtccca tgggactttt atatcaacct caagttaaag gaactgttaa atgaagattt 2640
tgatcatttt tgcagctggt atcaatatca agatggctgt attgtttggc accaatatat 2700
aaactgcttc acccttcag atcttctcca acacagtgaa tataattacc atgaataaac 2760
agttgtgatt atttataacc ttttgacaa agtgagatg ctacacaag cagaaatagt 2820
ccatggtgac ttgagtccaa ggtgtctgat tctcagaaac agaattccag atccotatga 2880
ttgtacaag aaccaataag ctttgaagat agtggacttt tctcacagt ttgacagtat 2940
ggtgcagctg gatgttttta ccttcagcgg ctttoggact gtacagatcc tgggaaggaca 3000
aaagatcctg ctaactggtt cttctcccta ccaggtagac ctgtttggta cctcagattt 3060
agcacattta catattgttca aggaacacct acaggtcttc tgggagtggtt cattgtggaa 3120
acttagccaa aatatttctg agctaaaaga tggtagaatt tggaaataat tctttgtggc 3180
gattctgaat gccaatgatg aggccacagt gtctgttctt ggggagcttg cagcagaat 3240
gaatggggtt tttagaccta cattccaaag tcaactgaac aaagccttat ggaaggtag 3300
gaagttaact agtctgggg ctttgcctct tcagtgagct agggcaatcaa gtctcagaca 3360
ttgtgcctc agagcaatg ttgtattgtg gaocactgaa actgtatgtt cgttaattta 3420
atttaggaca cattttgatg cactaccatt gctgttctac tttttgttac aggtataatt 3480
tgaogtcact gatatttttt atacagtgat ataactact atggccttgt ctaacttttg 3540
tgaagaacta ttttattcta aacgactca ttacaaatg ttactctgtt atttaaccca 3600
tttgtctcta cttttccctg actttttccc atttgaatt tgtaaaatgt tctctataga 3660
tcaccatgta ttttgttaat aataaaatag tatctgttaa aaaaaaaaaa aaaaaaaaaa 3720
aaaaa 3725

```

```

<210> 7
<211> 567
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<222> (1)...(567)
<223> n = A,T,C or G

```

```

<400> 7
ggccaagaat tcggcacgag gacaacatac taaaggaggc aggcaatgac tgttggccag 60
ttctccacgg ggaaaaaacc actgttagga tggcatgaac atttctttag atctgtggna 120
gtccggagga atgtggcgctn caggctcttt gagagccatg ggtgcaccc ggccttaggc 180
tagtgaact cgcaccccat tgcagtcgac tttcttgact gtgtgtctgt 240

```

aacccgtgctg	aggetccaca	tagctcctgg	acctgtgtct	agtacatact	gaagcgatgg	300
tcagagtgtg	tagagtgaag	ttgctgtgcc	cacattgttt	gaactcgcgt	accccgtaga	360
tacattgtgc	aacgtttctc	tggtattccc	ttgaggtgg	aacttcogtat	gttcagttta	420
tgcgtagatt	gttgtaaatg	caatgcogta	gtttggatta	ataagtgat	gtgttttgtt	480
tctaaaaaga	aaaaaaaaat	cagtggtcac	ccttatagag	acatagtcaa	gttcagtgtg	540
ataataatca	aaggaattac	tctcttc				567

<210> 8

<211> 1365

<212> DNA

<213> Homo sapiens

<400> 8

acttcatgaa	cacggacaat	ttcacctccc	acggtctccc	ccacccctgg	tcgggcacgg	60
ggcaggtgg	ctacacacgt	tctatctact	ttacacaagt	ccagagccac	atcatcatca	120
gggtttgacct	gaagacagag	accatcctca	agaccccgag	cctggactat	gcgggttaca	180
acaacatgta	caactacgcc	tggggtggcc	actcggacat	cgacctcatg	tcgacagaga	240
cgccgggtgtg	ggccgtgtac	gccaccaacc	agaacgctgg	caacatcgtg	gtcagtaggc	300
tggaccctgt	gtccctgcag	accctgcaga	cctggaacac	gcgtaccccc	aagccgacgc	360
ccggggaggg	cttcatcctc	tgccggcacg	tgtagctcac	caacgggtac	tcagggggta	420
ccaaggtcca	ctatgcatac	cagaccaatg	cctccacctc	tgaatacatc	gacatcccat	480
tccagaacaa	atactcccac	atctccatgc	tggactacaa	ccccaaaggc	cgggcccctg	540
atgcctggaa	caacggccac	cagatcctct	acaaactgtac	cctcttccac	gtcatccctg	600
cgaacgagtt	gtagctccct	cctcctggaa	gccaaaggcc	cacgtcctca	ccacaaaggg	660
actcctgtga	aactgctgcc	aaaaagatac	caataacact	aacaataccg	atcttgaata	720
atcatcagca	gtgcggatcc	tgacatcgag	ggatggcatt	acctccgtgt	ttctcccttt	780
cgagccggcg	ggccacagac	gtcggaaaga	actcccgat	ttgcagctgg	aactgcagcc	840
caocggccgc	cggttttctc	ccccgcctg	tcctctctgt	gtcaaacac	atactaaaga	900
ggcgaggcaa	tgaactgttg	ccagttctca	ccggggaaaa	accactgttt	aggatggcat	960
gaacatttcc	ttagatctgt	gtcagctccg	aggaaatgtg	gcctcaggtc	ctttgagagc	1020
gactgggtgc	acccggccgt	aggctatgtt	aactcgcatc	ccattgcagt	gcggtttctt	1080
gactgtgttg	ctgtctctta	gattaacctg	gctgaggtcc	ccatagctcc	ctggacactgt	1140
gtctatgata	tactgaagcg	atggctcagag	tgtgtagagt	gaagttgctg	tgcgccacct	1200
gtttgaactc	gcgtaccocg	tagatacatt	gtgcaacgtt	cttctgttat	tcctctgagg	1260
tggttaactc	gtatgttcag	tttatgcgat	gattgtgtga	aatgcaatgc	cgtagtttgg	1320
attaataaagt	ggatggtttt	tgtttttaaa	aaaaaaaaaa	aaaaa		1365

<210> 9

<211> 1196

<212> DNA

<213> Homo sapiens

<400> 9

ctcagctcta	ggggaatgaa	ggctgttttg	ctggctgata	ctgaaataga	cctttttctc	60
acagacatcc	ctccctacaa	cgcagtgagc	ttcaactgaa	gatgctatct	caccaaatac	120
tgcaaatata	aactgaagga	catcgcatgt	ttaaaatgtg	gaacacattg	agtttatcat	180
gtgattgttc	catgtagtcc	ctgtctctct	tctctgaaca	acagacactc	ctggagtgtt	240
cacagccagg	cagttttatga	tattaaacaga	ctagaactcca	caggtgttaa	cgctcactat	300
cggggcaact	tgccagagat	agaagagagt	acagatgaag	atgtgttaaa	tatctcagca	360
gaggagtgtg	ttagataaat	ggaattatga	tatatatgat	atacaaaact	ttttctattt	420
aaaaatatat	taattgatca	actttaaaat	tgttagttgc	cagtgatctt	tttggaaaaa	480
caaaaatggg	gcatttgttg	atttatttat	tttctgtctc	taattagtta	cctcagtttg	540
attgaagcca	gtggagttgt	gcttttcttc	tactctctact	tcctctcccc	cactcttttc	600
tgcccaagtgt	aggtgtatcc	ttaaattccag	aogggaaagt	ttcttccact	atcactcagt	660
caccctccaa	tctgggggag	tttttcttac	aacttgatac	cagataccat	taattttaca	720
ttctctgaata	aaggcctagt	accacgcgat	atttcaacca	tgcatatata	aagttcaacy	780
gagttttaat	agggggttaa	aaaaacaagc	tgtaggtgtt	ccatgggcac	tggtttctcat	840
aggttctatt	ggtgataact	gctttaacat	ggagcaagag	tttgtgaatc	aggaatata	900

```

ataaattaaa attttaaata tatagaggaa tctctcttgat tgctcagcat gatgttagat 960
aaatgagttt gtcagaaaaa atcagttatc gctgtttacc aatgtttattt atttacattc 1020
ttctaaagcc attatggata ttgtattatg agagctaaac ctaaaataagt tatctcgttc 1080
cctaggacct tctctgtaaa tagtgaattt tagacagata gctctgccta aatctttaat 1140
agaaaaaaa actaaagcga ttgtcttaag ccattgtaca ttataaagag ctgttt 1196

```

```

<210> 10
<211> 1424
<212> DNA
<213> Homo sapiens

```

```

<400> 10
ctcagctcta ggggaatgaa ggctgttttg ctggctgata ctgaaataga ccttttctct 60
acagacatcc ctctaccocaa cgcagtggaac ttcaactggaa gatgctattt caccaaaatc 120
tgcaaatgta aactgaagga catcgcatgt ttaaaatgtg ggaacattgt agkttatcat 180
gtgattgttc catgttagttc ctgtctctct tctcgcaaca acagacactt ctgggatgtt 240
cacagccagg cagttttatga tattaacaga ctgaaactcc caggtgtaaa cgtctactat 300
cggggcaact tgccagagat agaagagagt acagatgaag atgtgtttaa tatctcagca 360
gaggagtgtg ttagataaat ggaattatga tatatatgat atacaaactt tttctatttt 420
aaaaatatat taatggatca actttaaaat tgttagttgc cagtgtattt tttkggaaaa 480
caaaaaatgg gcaattgttg atttatttat ttctgtcttc taattagtta cctcagtttg 540
attgaagcca gtggagttgt gcttttcttc tactctactt cctctctccc cacttttttc 600
tgcccaagtg aggtgtattc ttaaatccag acgggaagat tctttccact atcactcagt 660
tacctcccaa cctgggggag tttttcttac aacttgatac cagataccat taattttaca 720
ttctgaata aaggcctagt acccagcat atttcaacca tgcataatc aagtccaacy 780
gagttttaat aggggattaa aaaaacaagc tgtaggttt ccatgggcac tggttctcat 840
aggtttctat ggtgataact gctttaacat ggagcaagag tttgtgaatc aggaatataga 900
ataaattaaa attttaaata tatagaggaa tctctcttgat tgctcagcat gatgttagat 960
aatgagttt gtcagaaaaa atcagttatc gctgtttacc aatgtttattt atttacattc 1020
ttctaaagcc attatggata ttgtattatg agagctaaac ctaaaataagt tatctcgttc 1080
cctaggacct tctctgtaaa tagtgaattt tagacagata gctctgccta aatctttaat 1140
agaaaaaaa actaaagcga ttgtcttaag ccattgtaca ttataaagag ctgttttgtt 1200
ttgctttgtc ttgctttgtt ttgttttttt taaagctgca ttcagagcca caaaggaata 1260
ggaaagttag gtatgttttg attctgtttt tatgttaact taaaataaat gbatctcttt 1320
aatatctcag ttgtagggat tttgtcaata ccaagcaga ctgagttgtg gttttgttaa 1380
taaagtttt tctaaaaatg aaaaaaaag aaaaaaaa aaaa 1424

```

```

<210> 11
<211> 460
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> misc_feature
<222> (1)...(460)
<223> n = A,T,C or G

```

```

<400> 11
agacagngac gatatgaaaa gntcttaaca gatnatttaa atgacctcca gggctgcaat 60
gatnatgacg ccaatgggac tngggacttc tatggggaca ntgtgtttgt gaaccagatg 120
atgaaagtgg caaggccaaa caggatncaat ngccttagag nagaanaana agatgatgat 180
gacgatgcct atcagngatg tgtttgaatt ngaattttca gagacccccc tcttaccgtg 240
ttataacatc caagtatctg tggctcaggg gccacgaaac tggctactgc ttcggatgt 300
ccttaagaaa ttganaatgt cctccgcgat atttgcgtgc anttttccaa acngngaaat 360
tgtaccatt gcagaggcag aattttatcg gtaggtttct gcnagtctct tgnctctctg 420
ctccaaagac ctggcaagcc ttcaaccctt gaaagngaan 460

```

```

<210> 12

```

<211> 2206

<212> DNA

<213> Homo sapiens

<400> 12

```

cagaagacag atgtgctgtg tgcagacgaa gaagaggatt gccaggctgc ctccctgctg 60
cagaataaca cgcacaacag cgagaagcca tcggggaaga gactgtgcaa aaccaaacac 120
ttgatccctc aggagtcocag gcggggattg ccactgacag ggaataacta cgtggagaat 180
gcgcgatggca aggtgactgt ccggagattc agaaagcgcg ccgagccocag ttcggactat 240
gatctgtcac cagccaaagca ggagccaaag cccttcgacc gcttgcagca actgctacca 300
gcctccocagt ccacacagct gccatgtctca agttcccttc agggagaccac ccagtctcgc 360
cctatgcgcgc cggaaagcacg gagacttatt gtocagtaaga acgctggcga gaccttctgt 420
cagcgggcag ccaggccttg ctaggaggaa tgggtcctgt actgcttaga gaacaagatt 480
ttgtagttaa atcatcggga caacgcagggt tactgcgcgc tgcagtaagc ttgtgctagg 540
ggctggctca acattgtgcg acacctcctt gaatatggcg ctgatgtcaa ctgtagtgc 600
caggatggaa ccaggcctct gcacgatgct gttgagaacg atcacttgga aattgtccga 660
ctactctctc ctatgtgtgc tgacccacc ttggctacgt actcaggtag aaccatcatg 720
aaaatgaccc acagtgaact tatggaaagg ttcttaacag attattttaa tgacctccag 780
ggctcgcaatg atgatgacgc cagtggcact tgggaactct atggcagctc tgtttgtgaa 840
ccagatgatg aaagtggcta tgaatgttta gccaaacccc caggaccaga agaccaggat 900
gatgatgacg atgcctatag cgaatgtgtt gaatttgaat ttccagagac cccctcttta 960
ccgtgttata acatccaagt atctgtggct cagggggtgag catggctgtc atgtgattga 1020
aaactagctg agctgctctt gaggccacga aactggctac tgctttcgga tgtccttaag 1080
aaattgaaaa tgtctcccg catatttcgc tgcatttttc caaaocgtgga aactgcaacc 1140
attgocaggg cagaatttta tcggcagggt tctgcaagtc tcttgttctc ttgtcccaaa 1200
ccttcgaaag ccttcaaccc tgaaagtaag gatctgttag attcaggaac attcaggaac 1260
gaaattcaga ctctgctggy ctctctgta gatggctcc accccagtaga ttggtgctca 1320
gacaactact ggtgagcaag ctggaccac catgtacagt gtgttatgt gttaactctt 1380
gtgcatatgt gctataatac aactatttct gtaagaagag gacactatta catatgaaa 1440
tatctctctt ttatataaga gaaattactc cagtcagaa gaottgaaa catgtttttt 1500
tcctttttaa cttttaagtc agtttttat agttgttat atgttttct taactttcaa 1560
tgcaacatg ctctgggata ogtttgttt tacttggaa atttgtttct tttctttttt 1620
aaggagaaaa aaaaaatgag taaaaggagc tccacacttt gacttaatt catacaaagc 1680
tctgatgaca ggccatgact gtgagtggt cagaactgtg tggttggtt gagggagcga 1740
attcggggaa ggcacttggt gatataactt tgttttggt acagagtacc gtctcgggcc 1800
aggtaaatgc tattgtagt atccagtag tgtgtaata aaattcaaac catatccaca 1860
cacacaact aattgtatga aacttttata tctaattta aaagctgtga aattagtttt 1920
cagcgtacaa accggtattt ttatatgttt aaacatttta tgctcttatt taaagaagac 1980
tttagactat tttttctgt accctgtaaa atattgaaaa atgttgaggt atgttgaggt 2040
tgcttggaat tgtacataaa actaaaattt tctgaatcgt gtgtttatgt ttgaaatctg 2100
tgttttaact tgttaagtaa attctctgcc ttgttatata tattttacaa aattttctta 2160
aaaggcataa aactgttgag gaaaggagaa aaaaaaaaaa aaaaaa 2206

```

<210> 13

<211> 680

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<222> (1)...(680)

<223> n = A,T,C or G

<400> 13

```

ataagatccc agctttgcgg gaactcatgc actatctcag ggaggtgatg caggattacc 60
gagatgagct caaggacttc ttgcagttgc acaaacagct gccatcagag cttgagtatg 120
acatgaagaa gtaccaggaa cagctggtcc aggagcagga gctacgaaaa catgcagatg 180
tgcccgggac ggctggaggt gctgaggtgc cactctggcc acaggttgcc ctgtgttag 240

```

```

aaacagtgcc agttcctgct ggccaagaaa accctgccat gtcacctgcc gtgagccagc 300
cctgcacacc caggggcaagt gctggccatg tagcagtatc atctcctaca cctgaaacag 360
ggccatttga caggttgctg cccaaagcca ggcccatgtc cctgagcacc attgcaatcc 420
tgaattctgt caagaagacc gtggagtcaa agcagcggca gtggagtacc agtctgagg 480
tgctgccttt cactttaaat tctggaaacc cagaaaaaac gtgcagtcat ggtctctcat 540
acagttttag gcaagatgct aatggcgaga ttgagcacgt gaccaaagcg gcatcacga 600
cccccagaaa gagcatcagt gatgtcacgt ttggagcan gggccaagtt acatcgggac 660
accaogggac ttcctgtctc

```

<210> 14

<211> 5023

<212> DNA

<213> Homo sapiens

<400> 14

```

ggcggcgggc agccggtgcc ctgggatcat ggtggcggtt cggggccttg gtacggcct 60
cgagccctgg tgcctcgctg atcttagact cgaatggggt gacacagtgt ggaacttga 120
tttcacagag actgagcctt tggatccagc catagaagca gagatcatag aactggatt 180
ggctgcattc acaaaaactc atgaagacct ttacaccttt gctactggag aaactggatc 240
tatggagagt atctggaact tcttcattga gaacaatgtt tcccatagta cactggtggc 300
atgttctcat catcttggtc aaatagttca taagaagaat gtcagtgtac agtatcgaga 360
atatggcctt catggcgctg ggctttaact ttgtctaact gaagtaccag gcagtgtagc 420
caatcaagta ttccaccagc tgatgtttga caaatgcatt cagactctaa agaagagctg 480
gccccaggaa tctaacttga atcggaagaa aaagaaaagaa cagcctaaga gctctcaggc 540
taaccccggg aggcatagaa aaaggggaaa gccaccagag agagaagata ttgagatgga 600
tgaatttata gaagaacaag aagatgagaa tatttgtttt tctgcccggg accttctca 660
aattcgaaat gccatctttc accttttaaa gaatttttta aggctttctg caaagttttc 720
cttgaagaaa aagccacaat gtgtacagaa ttgtatagag gtctttgttt cattaactaa 780
ttttgagcca gttcttcatg aatgtcatgt tacacaagcc agagctctta accaagcaaa 840
atacatacca gaactggcct attatggatt gtatttgtct tgctctccca ttoactggga 900
aggagataag gtcacagttt gtgttttcca tcaaatgtct agtgaatat taatgttaga 960
agttggtgaa ggtaccatct gtgcccccct tgcgtttacc tcccaagtca tcaactgtga 1020
aaaccaggcg gtccagttta tgcgcgcctc ttgggatgaa ttaaaggaga gtatattccc 1080
agtgcctcgt atcttactgc agcacatctg tgccaaagtg gtatataaat cagagtatcg 1140
tacttttgca gccacgtccc tagtccagct gctcagtaaa ctctctgtgt gggaatacgc 1200
tatgtttcgt gctcggtctt acaaatctc cogaagtccc aagatccccc accgggtttt 1260
tactcttgat gttgttctag ctctgttaga actgcctgaa agagaggttg aataacacct 1320
ctccttgtag catcagaagt tcttaaaaga taagttcctg gtgcaggaaa ttatgtttga 1380
tcgttgctta gacaaggcgc ctactgtccg cagcaaggca ctgtccagct ttgcaacctg 1440
ctcggagttg actgtttacca gtgcgtcgga gagtatcctg gactcctga ttaacactgc 1500
taogttttct gtaatataga gtccacctggt acccttactg agaattcat cagotttttc 1560
ctaccaaggg cagacatcta accgttccga tgcctcaggg agactcaaca tagacagcag 1620
tggtgaacaa gtgggatctg gagaagatg tgcctatgga atgctgagaa ggaggatcag 1680
ggatgagaag accaacgtta ggaagtctgc actgcagta ttatgtagta ttttgaacaa 1740
ctgtgatctc tcaggcatga aggaagacct gtggattctg caggaccagt ttcgggaacc 1800
tgcagtgctt gtcgcgaagg aggccttcca gtctcttact gaactccta ttctcgagcc 1860
tagatgcgtg cagatccaga aagcctgtgt gcgggggggt gtcocggtgg tgatggaact 1920
cgagagcact gtgcagagga aggcctgtga gtctctggac gactgtctgc tgcagaacct 1980
ccggcatcac agtcattttc actctgggga gaggcgccag gtcctgcctc ggcgcctctc 2040
tactctcttc accaccgaaa gccagggaact gaggcgatat ttaataaagg ttcttcatat 2100
ctggtccaaag aaggaaaaaat tctcacccac ttttataaac cttaactatg ctcaactgtg 2160
cagggaacat tcggcacctg cctggatgct gctctccaaag attgctgggt cctcacccag 2220
gctggaactc agcagaataa tacaactctg ggagaaaatc agcagtcagc agaattccaa 2280
ttcaaacacc ttaggacata ttctctgtgt gattgggcat attgcaaagc atcttcttaa 2340
gagcaccggg gacaaaagtg ctgatgctgt caagtgtaa gtcagtgtgc atctctgttc 2400
cttagaggtg atcagttcag ctgttgacgc cttgcagagg ctttgtagag catctcgaga 2460
gacacacgga gaggagcagg aattgctgac cgaggtgtgt ggggatgtac tctccactgc 2520
cgagcaccgc ctctccaaca tcgttctcaa ggagaatgga acaggggaata tggacgaaga 2580

```



```

cctgttggtg aagtacattt ttaccttagg ggatatagcc cagctgtgtc cagccagggt 2640
ggagaagcgc atctctcttc tgattcagtc cgtctctggt tcgtctgtgt atgtcgacca 2700
ctcaccatca tctcaaggca gcagtgaggc cccagcgtct cagccacccc cccaggtcag 2760
aggttctgtc atgcccctctg tgatttagagc acatgccatc attaccttag ctaagctgtg 2820
cttcacagca gaggatctgg caaagaagag catccagacc ctggtgcgag agctcgaggt 2880
gtgtgaggac gtggctgtcc gcaacaacgt catcatgtta atgtgcgac tcctcatctg 2940
ctacaccatc atgtgtggaca agtatattcc caacatctcc atgtgtctga aggatctcga 3000
ccattcatcc cggaaagaga cactcatctt gctttaccaat cctttgcagc caggaattgt 3060
gaaatggaag ggctccctgt tcttcogatt tgtcagcact ctgatcgatt caccccaga 3120
cattgcagca ttccgggagt tttgcctggc tcacctgtta ctgaagagga accctgtcat 3180
gttcttccaa cacttcattg aatgtatttt tcactttaat aactatgaga agcatgaga 3240
gtacaacaag ttcccccagt cagagagaga gaagcggtg ttttcatgga agggaaagt 3300
aaacaagag agacgaatga atttcttcta gacacactca cagatgaaca 3360
gcatcctaac atcacttcca aaatctgcct tagtattttg gctgtctttg ctgatggcat 3420
cctaccctcg gacottggaag ccagtgaagt actctcagac acgtttgagg tctcagctc 3480
aaaggagatc aagctttttg caatgagatc taaaccagac aaagacctcc ttatggaaga 3540
agatgacatg gcccttggoaa atgtagtcat gcaggaagct cagaagaagc tcactctaca 3600
agttcagaag aggaatttca tagaaaaat tattccaatt atcatctccc tgaagactgt 3660
gctggagaaa aataagatcc cagctttgcg ggaactcatg cactatctca gggaggtgat 3720
cgaggattac cgagatgagc tcaaggactt ctttgcagtt gacaacacgc tggcatcaga 3780
gcttgagtat gacatgaaga agtaccagga acagctggtc caggagcagg agctagcaaa 3840
acatgcagat gtggccggga cggctggagg tgtcgaggtg gcacctgtgg cacaggtgtg 3900
cctgtgttta gaaacagtcg cagtctctgc tggccaagaa aacctgccca tgtcaactgc 3960
cgtgagccag ccctgcacac ccaggggcag gtctggccat gatcgagtat cactctcaca 4020
acctgaacac gggccatttc agaggttgtct gcccaagacc agggccatgt cctcgagcac 4080
cattgcaatc ctgaaattctg tcaagaaagc cgtggagtca acgagcagcg atcggagctg 4140
gagcttagga gtgtgccttt tcaactttaaa ttctggaagc ccagaaaaaa cgtcgactga 4200
gggtctctca tacagtttct agcaagagtc attgagcacg ttgagcacg tgaccaagcg 4260
ggccatcagc acccccagga agagcatcag ttagtgcacg tttggagcag gggtcagtta 4320
catcgggaca ccaaggactc cgtcgtcagc caaagagaaa attgaaggcc ggagtcagg 4380
aaatgacatc ttatgtttat cactgcctga taaaccgccc cccagcctc agcagtggaa 4440
tgtgoggctc cccgccagga ataaagacac tccagcctgc agcaggaggt cctctcgaaa 4500
gacccctctg aaaaacagcca actaaacagc gccctcccac agtgtccagg caggcaggag 4560
cccttgagga agcagctctcg tgtcctccgt gtgaaggcag ctggatcact tcccgagtc 4620
cttgggcagc gctttgctgt ggaacacgag agctcctcct caggggcctg cactcactct 4680
tctattctgt atgatgtatt tggttaaaaca ctgtcaaatat atagagatgt gccagatta 4740
gtatttctta cctaatctgt tttaatattg taactttatt cagtttgaaa gtgtcaagtc 4800
cattcagata agctataatc tggttcttaa ggaatacaac tttaaaactt cagctttct 4860
ttatatataa caagcctctg ttaacttgaa ttcttatag tacatatatt ccactctgta 4920
atgcocggaat tttgattcta atattttttc tattatttat aagtgcacaa tttttttaaa 4980
aggtgacagc tttcttaaaag taataaaggt ttgacataaa tac 5023

```

<210> 15

<211> 403

<212> DNA

<213> Homo sapiens

<400> 15

```

ccatcacggg gaattctgct gctgttatta cccatttcaa gttgacaact gaggcacgc 60
agactccagt ctccaataag aaaccagtg ttagctctaa agcaagtttg tctgtctccc 120
tcaactatga accacacaaa ggaagagctaa aacctctaaa gcaactctaa gaaaataatt 180
ctataataca acatgtcaac agaattaaact tctacaaaga aacttacaaa caaccccact 240
tccagacaaa ggaagagcaa cggaaagaac gcgagcaaga acctaaagag aagaagacaa 300
aggttttggg aatgcgaagg ggccctcatt tggctgaaga ttaataattt ttaaacatct 360
tgtaataatt cctgtattct caactttttt ccttttgtaa att 403

```

<210> 16

<211> 890

10

<212> DNA
 <213> Homo sapiens
 <220>
 <221> misc_feature
 <222> (1)...(890)
 <223> n = A,T,C or G

```

<400> 16
agcataagcg tntcactgac caagactoca gccagaaagt ctgcacatgt gaccgtgtct 60
gggggcaccc aaaaaggcga ggctgtgctt gggacacaca aattaaagac catcacgggg 120
aattctgctg ctgttattac ccattccaag ttgacaactg agggcaacgca gactccagtc 180
tccaataaga aaccagtggt tgatctttaa gcaagtttgt ctgctccctt caactatgaa 240
ccacacaaag gaaagctaaa accatggggg caatctaaag aaaataatta tctaatacaa 300
catgtcaaca gaattaactt ctacaagaaa acttacaaac aaccccatct ccagacaaa 360
gaagagcaac ggaagaaacg cgagcaagaa cgaagagaga agaagcaaaa ggttttggga 420
atgcgaaggg gcctcatttt ggctgaagat taataatttt ttaacatctt gtaaatattc 480
ctgtattctc aacttttttc cttttgtaa tttttttttt ttgtctgtca tccccattt 540
agtcacgaga tctttttctg ctactgttct atagtctgtg gtagtgtcca tgggtctctc 600
atgtgtgatg atctctgaaa agacgtttat accctaaagc tcaaatcttt tgggatggtt 660
tttacttaag tcatttaaca attcaggttt ctaacgagac ccactctaaa attctgttct 720
tagattttta atgtcaagtt cccaagtttc ccctgtgggt tctaataata acagaactgc 780
agtctcttgc tagccaatag catttacctg atggcagcta gttatgcag ctttagggag 840
aatttgaaca ttttccagga atgggggaag ctgggaaga aaggccaact 890

```

<210> 17
 <211> 371
 <212> DNA
 <213> Homo sapiens

```

<400> 17
ttggtctcagc aggacaatat ggtgggaaat gacaaagtaa ctctgtggc cctaggtcag 60
gttctcttga gaaaaacaaa aaggctggaa tgatacagct ctctgtaaac caggtgcctc 120
cagtgctctg ggttattccc aagtccacat ttgacagaca gggccctaaa atgtctagct 180
aggaagttcc tgagcctggt tttttaaaaa tctacacaca cacatgcaca cacacagca 240
cgtgtgcaca catgcggata tatacatcct caccctttct tgagattact gtcagaaga 300
aggcacattt ggtttggtct gcttaccagg tgctgaagt ggagggcgcc caagcttawt 360
tccttttagt g 371

```

<210> 18
 <211> 376
 <212> DNA
 <213> Homo sapiens

```

<400> 18
attctttggc tcagcaggac aatatggtgg gaaatgacaa agtaactcct gttggccctag 60
gtcaggttct cttgaggaaa acaaaaaggc tggaaatgata cagctctctg taaacacagt 120
gcctccagtg cctgcggtta ttcccaagtc cacatttttg agacagggcc ctaaaatgct 180
tagctaggaa gttctctgagc ctgttttttt aaaattctac acacacacat gcacacacac 240
acgcacgtgt gcacacatgc ggatatatac atcctcacct ttctttgaga ttactgctca 300
gaagaaggca catttgggtt ggtctgctta ccaggtgtctg aagtgaggag ggcgcgaagc 360
ttawttcctt ttatgt 376

```

<210> 19
 <211> 512
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <222> (1)...(512)
 <223> n = A,T,C or G

<400> 19
 ccatgttgata ctgtatgaac ctangtagnt tggagaagaaa agtaggggtt ttgtatacta 60
 gcttttgtat ttgaattaat taccattcca gctttttata tactatattt catttatgaa 120
 gaaattgatt ttcttttggg agncaactttt aatctgtaan tttaaaatac aagctcgaat 180
 attttagt ttgattcttaac tgtgcataana octagatata ccattatccc ttttatacct 240
 aanaaggcca tgctaataat taccactgtc aaagaggcaa aggnnggtgat ttttgnntat 300
 gaagttaagc ctccagnggag gctcatttgt tagtttttag cngganctaa ngntaaactc 360
 agggntccct gagctatatg cacactcaga cctctttgct ttaccocagng gcgttngtga 420
 gttgctcagc agtacaacact gcccttacct gacagagccc tgnctttgac ctgctcagcc 480
 ctgtgcgcta atcctctagt agcccaatca na 512

<210> 20
 <211> 3410
 <212> DNA
 <213> Homo sapiens

<400> 20
 gcaccaggcg cccagtggag ccgtttggga gaattgctcg gcggcagcag cggggccgga 60
 caggcggttaa ggaatcgatt aggccttcga acttgagttt gactgatgtc ttctgtgtgg 120
 tgcctcgctaa atccacagc atataaggatc agtcgacatt gttataaagt ttgcttctgg 180
 ctgggtgcgg tggctcagtc ctgtaatcca acattgggag gccaaaggcag gcggaccacc 240
 tgaagctcgg agcttgagtc cagccactgt ctgggtactg ccagccatcg gggccaggctc 300
 tctgggggttg tcttacgcga gtgagtacca cgcggtacta cagagaccgg ctgcccggtg 360
 gcccgccagg tggagccgcc gcatcagcgg octcggggaa tggagccgga gaaocggggc 420
 agctattccc ttcagcaagc tcaagctttt tatcgtttc cattcaaca actgatggct 480
 gaagctccta atatggcagt tgtgaatgaa cagcaaatgc cagaagaagt tccagcccca 540
 gctcctgctc aggaaccagt gcaagaggct ccaaaaggaa gaaaaggaaa acccagaaca 600
 acagaaccaa aacaaccagt ggaacccaaa aaacctgttg agtcaaaaaa atctggcaag 660
 tctgcaaaac caaaagaaaa acaagaaaaa attacagaca catttaaagt aaaaagaaaa 720
 gtgacgcgtt ttaatgggtg ttcagaagct gaacttctga ccaagaactct ccgcgatatt 780
 ttgaccttca atctggacat tgtcattatt ggcataaacc cgggactaat ggctgtctac 840
 aaagggcact attaccctgg acctggaaac catttttgga agtgtttgtt tatgtcaggg 900
 ctcaagttag tccagctgaa ccatatggat gatcacactc taccagggaa gtatggtatt 960
 ggattttacca acatgggtgga aaggaccacg cccggcagca aagatctcgc cagttaaaga 1020
 ttctgtgaag gaggacgtat tctagtacag aaattacaga aatatcagcc acgaatagca 1080
 gtgttttaag gaaaatgtat ttatgaaatt tttagtaag aagtttttgg agtaaaagtt 1140
 aagaacttgg aatttgggtc tcagccocat aagattccag acacagaagc tctctgctat 1200
 gttatgcoat catccagtg ccaagatgtct cagtttctct gagcccaaga caaagttcat 1260
 tactacataa aactgaagga ctttaagagat cagttgaaag cagttgaaag caaattggac 1320
 gttcaagag tgcaataaac atttgacctc cagcttggcc aagagagatgc aaagaagatt 1380
 gctgtttaag aagaaaaata tgatccaggt tatgaggcag catatgtgtg tgcttaccga 1440
 gaaaatccat gcagcagtg acccttgggc ttctcttcaa atgggctaat tgagagcgtg 1500
 gaggtaagag gagaatacgc ttctagtggc attcctaag ggcagtggt gaccagctca 1560
 ttacacagac aaattctctc cttttagtat cactgtgaa aacaagaaca ggaagaagaa 1620
 agccatctgt aagaattggg cttctcagct ctgcctaaat gctgcagttt taatgcagtt 1680
 gccaacagt agaacctcag ttgtctaact gaagtgtttt attgatattt taatgcagtt 1740
 gtgtaattgt aatgtgaaac agttgtgtgg tagtgtgaac cgtatgaacc taagtatttt 1800
 ggaagaaaaa gttaggtttt tgtatactag cttttgtatt tgaattaaat atcaattcca 1860
 ctttttatat acctaatctt atttatgaag aaattgattt tcttttggga gtcactttta 1920
 atctgtaatt tttaaaatga agtctgaata tttagattg attcttaact gtcataaacc 1980
 ctgatatata cattatacct tttataacct agaaggcact gctaataatt accactgtca 2040
 aagaagcaaa ggtgtgtgatt ttgtatata agttaagct ctagtgaggt cttattgttc 2100
 gtttttagtg gtaactaagg gtaaacctag ggttccctga gctatatgca cactcagacc 2160

12

tctttgcttt	accagtggtg	tttgtgagtt	gtcagtagtg	aaaaactggc	ccttacctga	2220
cagagccctg	gctttgacct	gctcagccct	gtgtgttaat	cctctagtag	ccaattaaat	2280
actctggggg	ggcaggttcc	agagaatcga	gtagaccttt	tgccactcat	ctgtgtttta	2340
cttgagacat	gtaaatatga	taggggaagg	actgaatttc	tcattcata	tttataacca	2400
ttctagtttt	atcttccctg	gctttaaagg	tgtgccatgg	aaagtgtata	gaatgaact	2460
tctaggctaa	gcacaaaagat	gctggagata	tttgatactc	tcatttaaac	tggtgcttta	2520
tgtaactag	atgtactaaa	ataagtata	tagaattttt	cttgctaggt	aaatccagta	2580
agccataaat	tttaaagatt	ctttatctgc	atcattctgt	tttgttacta	taaatataat	2640
gaacctcatg	gaaaggttga	ggtgtatacc	tttgtgattt	tctaatgagt	tttccatggt	2700
gctacaaata	atccagacta	ccaggtctgg	tagatattaa	agctgggtac	taagaaagt	2760
tatttgcatc	ctctcagtta	ctcctgaata	ttctgatttc	atacgtaccc	agggagcagt	2820
ctgtttttgc	aatcaatata	aaatatttat	gaggtctccc	ccacccccc	gaggttatat	2880
gattgctctt	ctctttataa	taagagaaac	aaattcttat	tgtgaatctt	aaactgcttt	2940
ttagctgttg	ctatgatgga	ttttattttt	tcctaggcca	agctgtgtaa	aagtcattta	3000
tgtttattaa	atgatgtact	gtactgctgt	ttacatggac	gttttgtcgc	ggtgctttga	3060
agtgcccttg	atcagggatt	aggagcaatt	aaattatttt	ttcacgggac	tgtgtaaagc	3120
atgtaactag	gtattgtctt	ggtatataac	tattgtagct	ttacaagaga	ttgtttttatt	3180
tgaattggga	aaataacctt	taaattatga	cggacatcca	ctagagatgg	gtttgaggat	3240
tttccaaagc	tgtaataaat	atgtttttcc	taacatgaca	gatgagtga	aaactgtgat	3300
atatcctata	catgacagtg	tgagactttt	tcattataata	atattgaaag	attttataat	3360
tcatttgaaa	gtctgatggc	ttttacaata	aaagatatta	agaattgtta		3410

<210> 21

<211> 627

<212> DNA

<213> Homo sapiens

<400> 21

ggccaagaat	tcggccgagg	ggtgccgagg	ccatggagaa	gcttagctcc	atcaaatctc	60
aaacaattta	tgagattatt	gataaattctc	aaggattcta	cgtttgtcca	gtggagcccc	120
aaaatagaag	caagatgaat	attccattcc	gcattggcaa	tgccaaaggc	gatgatgctt	180
tagaaaaaag	atttcttgat	aaagctcttg	aactcaatat	gtgtgccttg	aaagggcata	240
ggctctgtgg	aggcatccgg	gcctctctgt	ataatgctgt	cacaattgaa	gaacttcaga	300
agctggccgc	cttcgatgaa	aaatttttgg	agatgcata	gctatgaaca	catcctaacc	360
aggatatact	ctgttcttga	acaacataca	aagtttaaag	taacttgggg	atggctacaa	420
aaagtttaaca	cagttatttt	ctcaaatgaa	catgtttatt	gcagattctt	cttttttgaa	480
agaaacaacg	caaaaacatc	acaactctgt	aaagctgtgt	ggacctaatg	tcaoccttaa	540
tctgacttga	actggaagca	ttttaagaaa	tcttgttgct	tttctaaca	attcccgctg	600
attttgctt	tgctgctctt	tttctag				627

<210> 22

<211> 1065

<212> DNA

<213> Homo sapiens

<400> 22

ccttggtgta	ctcacgcccc	tgcgcgcgcg	accatggagc	ccccaggcca	ggtggtcaac	60
tttgggacct	gtcccgcaca	gctgcgcgac	tcagtgttgt	tagagatata	aaaggaatta	120
ttagactaca	aaggagttgg	cattagtgtt	cttgaataga	gcacagagtg	atcagatttt	180
gccaaagtta	ttacaacata	agagaactct	gtgcgggaat	tcagctgctg	tcacagaaac	240
tataaaggta	tttttctgca	aggaggtggg	tgccggccagt	tgatgtctgt	ccocctaaac	300
ctcattgctt	tgaagcagg	aaggtgtgcg	gactatgtgg	tgacaggagg	ttggtcagct	360
aaggccgcag	aagaagccaa	gaagtttggg	actataaata	tcgttcaccc	taaaccttggg	420
agttatataca	aaattccaga	tccaagcacc	tggaaacctca	accagatgac	ctctcactgt	480
tattatttgc	caaatgagac	ggtgcactgg	gtggagtgtg	actttatacc	cgatgtcaag	540
ggagcagtag	tggttttgta	catgtcctca	aacttctctg	ccaagccagt	ggatgtttcc	600
aagtttgggt	tgatttttgc	tgtgtgccag	aagaatgttg	gctctgctgg	ggtaacgctg	660
gtgattgtcc	gtgatgacct	gctgggggtt	gccttcagag	agtgcccttc	ggtcctggaa	720

13

```

tacaagggtgc aggcgtggaaa cagctccttg tacaacacgc ctccatgttt cagcatctac 780
gtcatggggtc tggttctgga gtggattaaa aacaatggag gtgccgcggc catggagaag 840
cttagcttcaa tcaaatctcaa aacaatttat gagattattg ataattctca aggattctac 900
gtgtctgtgg gaggcattccg ggctctcttg tataatgctg tcaaatgga agctattcag 960
aagctggcgct ccttcatgaa aaaatttttg gagatgcata agctatgaac acatcctaac 1020
caggatatac tctgttcttg acaacatac aaagtttaaa gtaac 1065

```

```

<210> 23
<211> 578
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<222> (1)...(578)
<223> n = A,T,C or G

```

```

<400> 23
gctcggggcc aagaattcgg cagcaggcca agttaaggaa ottgaagcta atgtacttgc 60
tacagccctc gacaaaaaaa gcagaatttg ctagaagaaa acgttagtgc tttcaaaaac 120
gaatangang ctgnggctga gaaagctggt aaagtagaag ctgaggttaa acgcttacac 180
aataccatcg tagaatacaa taatcataaa ctcaaggccc aacaagacaa acttgataaa 240
ataaataaag aattagatga atgtgctctc gctattacta aagcccaagt agcaatcaag 300
actcgtgaca gaaaccttca aaaggcaca gactctgtct tgcgtacaga gaaagaaaaa 360
aaagataactg agaaagaggt ggatgcacta acagcagagc tgaagagtct tgaggacaaa 420
gcagcagagg tcgtaagaaa tacaatgctc gcagagcagt tcttttcggt gtttaggaat 480
ccttacocaga gatccagaaa gaacatcgca atctgcttca agaattaaaa gttattcaag 540
aaaaagaaca tgctotctaa aaagatgcct tagtatta 578

```

```

<210> 24
<211> 3799
<212> DNA
<213> Homo sapiens

```

```

<400> 24
atagtaaaccc agaacttcaa atcctatgct ggggagaaaa ttctgggacc ttccataaag 60
cgcttttctc gtattatcgg gccaaatggc agtggcaaat ccaattgtat tgattotatg 120
ctttttgtgt ttggctatcg agcacaaaaa ataagatcta aaaaactctc agtattaata 180
cataattctg atgaacacaa ggacattcag agttgtacag tagaagtcca ttttcaaaag 240
ataattgata aggaagggga tgaatttgaa gtacttctta acagtaattt ctatgtatcc 300
agaagcgctc gcagagataa tacttctgtc tatcacataa gtggaaagaa aaagacattt 360
aaggatgttg gaaattctct tcgaagccat ggaattgact tggaccataa tactttttta 420
attttacagg gtgaagttag acaaatgtct atgatgaac caaaaggcca gactgaacac 480
gatgagggtta tgcctgaata tttagaagat ataattggtt gtggacggct aaatgaacct 540
attaagtgct tgbtccaag agttgaata ttaattgaac acagaggaga gaagttaaac 600
agggtaaaga tgggtgaaaa ggaagaggat gctttagaag gagagaaaaa catagotattc 660
gaatttctta ccttgaaaaa tgaatatatt agaaaaaaga atcatgtttg tcaattatat 720
atttatgagt tgcagaaacg aatttgtgaa atggaaaactc aaaaaggaaa aattcatgaa 780
gatacacaag aaattaatga gaagagcaat atactactca atgaatgaa agctaagatc 840
aaagatgtaa aagatcacaga aaagaaactg aataaaatta caaaatttat tgaggagaat 900
aaagaaaaaat ttacacacgt agatttgaaa gatgttcaag tttagagaaa gttaaaaact 960
gccacgagta aagccaaaaa actggagaaa caacttcaaa aagataaaga aaaggttgaa 1020
gaatttaaaa gtatacctgc caagagtaac aatatcatta atgaaacac aaccagaaac 1080
aatgcctcgc agaaagaaaa agagaaagaa gaaaaaaatc taaaggagat tatggatagc 1140
cttaaacagg aaacacaaag gcttcagaaa gaaaaagaaa gctgagagaa agaactatgt 1200
ggtttcagca aatcggttaa tgaagcaogt tcaaagatgg atgtagccca gtcagaacct 1260
gatatactat ctacgtgtca taatactgca gtgtctcaat taactaaggc taaggaaagt 1320
ctaattgcag tctctgagac tctcaagaaa aggaagcgtc caatcagaga tatagaagga 1380

```

```

aaactccctc aaactgaaca agaattaaag gagaagaaga aagaacttca aaaacttaca 1440
caagaagaaa caaactttaa aagtttggtt catgatctct tcacaaaagt tgaagaagca 1500
aagagctcat gcagcaatgaa ttctgagtag gggaaagctc ttgatgcaat aatcgaagaa 1560
aaaaactctg gcaggattcc aggaatatat ggaagatctg ggcacttagg agccattgat 1620
gaaaaatcac acgtggctat atcatcctgt tgcctgcatc tggactacat tgttgttgat 1680
tctattgata tagcccaaga atgtgttaac ttctttaaaa gacaaaatat tggagtgtga 1740
acctttatag gttttagata gatggctgta tgggcgaaaa agatgaccga aattcaaatc 1800
cctgaaaata ctctcgtttt atttgattta gtaaaagtga aagatgagaa aattcgccaa 1860
gctttttatt ttgctttacg agatacctta gtatgtgaca acttggatca agccacaaga 1920
gtagcatatc aaaaagatag aagatggaga gtggaactt tacagggaca aatcatagaa 1980
cagtcaggta caatgactgg tggtggaagc aaagtaatga aaggagaagt gggttcccta 2040
cttgtatttg aaatctctga agaagagtta aacaaaatgg aatcacagtt gccaaaacgc 2100
tctaaaaaag caatgcaaat ccaagaacag aaagtaacac ttgaagaaa agtagttaag 2160
ttacggcata gtgaacgaga aatgaggaaac acactagaaa aatttactgc aagcatccag 2220
cgttttaatg agcaagaaga atatttgaat gtccaagtta aggaacttga agctaagtta 2280
cttgctacag cccctgacaa aaaaaagcag aaattgctag aagaaaacgt tagtgttttc 2340
aaaaacgaat atgatgctgt ggctgagaaa gctggtaaa tagaagotga ggttaaacgc 2400
ttacacacata ccatgttaga aatcaataat cataaacctc agggcccaaca agacaaactt 2460
gataaaataa ataagcaatt agatgaatgt gcttctgcta ttactaaagc ccaagttagca 2520
atcaagactg ctgacagaaa ccttcaaaag gcacaagact ctgtcttgcg tacagagaaa 2580
gaaataaaag atactgagaa agaggtggat gacctaacag cagagctgaa aagctcttag 2640
gacaaagcag cagaggtcgt aaagaataca aatgctgcag aggaatcctt accagagatc 2700
cagaagaac atcgcaatct gcttcaagaa ttaaaagtta ttcaagaaaa tgaacatgct 2760
cttcaaaaag atgcacttag tatataagttg aaacttgaac aaatagatgt tcacatgct 2820
gaacataatt ctaaaataaa atattggcac aaagagattt caaaaatctc actgcatctc 2880
atagagata atctctattga agagatttgc gttctaagcc cagaggtctc tgaagcgtc 2940
aagaattccag attctataac aaatcaaaat gcacttttgc aagcccggtg tcatgaaatg 3000
aaacccaaccc tccgtgtccat cgcagagtat aaaaagaagg aagaattgta tttgcaacgg 3060
gtagcagaat tggacaaaat tacttatgaa agagacagtt agatagagc atatgaagg 3120
cttcgggaac aaaggcttaa tgaatttatg gcagggtttt atataatac aaataaatta 3180
aaggaaaatt accaaaatgct tactttggga aactcgagct tctgaacgc 3240
tttgatcctt tctctgaagg aatcatgttc agtgttcgac cacttaagaa aagttggaaa 3300
aagatcttca acotttgggg aggagagaaa acacttagtt cattggcttt agtatttgc 3360
cttcaccact acaagccocac tccctttac ttcatggatg agattgatgc agcccttgat 3420
tttaaaaatg tgtctattgt tgcattttat atatatgaac aaacaaaaaa tgcacagttc 3480
ataaatattt ctcttcgaaa taatatgttt gagatttcgg atagacttat tgaatttac 3540
aagacataca acataacaaa aagtggttgc ttaaatccaa aagaatttgc atctaaagga 3600
ctttgttgaa ctttatctga agtctcaagt tgattcaagt attactgatt tttttctat 3660
tgtaaaggat tatgagttgt ataaaaata tactccctaa actagatcat gaaactggt 3720
ctgtttttat gcagttgtca ttgttaaagt ctaataaaat actctctata attgctctta 3780
gattacaaaa atatgacaa 3799

```

```

<210> 25
<211> 429
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<222> (1)...(429)
<223> n = A, T, C or G

```

```

<400> 25
attgggaacaa agaagtattt taaaattata actactcatt ctttcttttg ccttagttaa 60
tttgagcaga agccacaaca agcaaacacc aataaattta taattggcag aaatccacat 120
taactcctct tcccaagttt ccacactact accatttaca gttgtagggt tgtaagtat 180
aattatgtaa tgcagaaact agctttgact tgtgtaacga tgcactgtca aagtaagcaa 240
agtaagaatt gaaattccac attcccagaa tttaaacact agctgctcct atgtacttaa 300

```

15

```

gttcctggggg ataatacatt aaccaacatt ggttgaaaca tacctgagta atcatatcag 360
gatgcgatgtt aagctgataa aacaataaga tcccaaaatg cagtactgca aaaaaaaaaa 420
aaaaaagggn                                     429

```

```

<210> 26
<211> 788
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> misc_feature
<222> (1)...(788)
<223> n = A,T,C or G

```

```

<400> 26
ncctttttttt tttttttttt gagctactgc attttgggat cttattgttt tatcagctta 60
acatgcatcc tgatatgatt actcagggtat gtttcaacca atgttggtta atgtattatc 120
cccaggaaact tattactaga ggagcagctg agtggttaaat tctgggaatg tgggaatttca 180
attcttacttt tgcttacttt gacagtgcat cgttacacaa gtcaaagcta gtttctgcat 240
tacataaata tacattacaa acctacaact gtaaatggtta gtagtgtgga aacttgggaa 300
gaggaggttaa tgtggatttc tgcoaatctt aaattttattg tggtttgctt gttgtggctt 360
ctgctcaaat taactaaggc taagaagaaga atgagtagtt ataattttaa aatacttctt 420
tgttcccatc tagcacccct taogcgctga gatgaaaaaa cactttttgt tgagactaag 480
agcttattac tcttcccaag attctctggc aattcagatt ccccaacttc catatcagcc 540
atthttcttt aataaaggaa ctactgatat tcttgggcaa attattacct cctctggctc 600
agttgttttg acctggggct aatgagccca gggcctgggg tttgattccc acgcatgcca 660
attagctttg cttgcttcca ccaaccagg ctgccctatt aaagcctgcc gctgtccga 720
agatgccacc acacatcttg cttatgagt cattgtgcat aaaaggggcc agctaagtga 780
tagggaaa                                     788

```

```

<210> 27
<211> 687
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> misc_feature
<222> (1)...(687)
<223> n = A,T,C or G

```

```

<400> 27
acatgggttg tgctttactc ttaaacatct ttaagtgct attattctat atctgtttgga 60
tgagtcatta tttttgaaat gataaaccta gcatgaactc tgatctatgg tgttggattc 120
tgtttcttaa ataactttaa aattaactgt tttccttgga gatttccctc tccatgtgag 180
gtatttgagc tattgttcta agtttaacctg taagtataaa ccttgggaga atctaaagtaa 240
acatatcttc aaaagcatag ttaccttctc attttctggc tottaacctc ttggagtatt 300
taaatgccca tttgccaaaa gcagacctga acatcaagcc tgttaattct tcaagaattt 360
taggtatttg tttcacogaa atgaagtgcac ttattagcca ttcagcgtat tagtattaca 420
gaggctcttg ccagccaca tccattcatt gatttttatg gctactcttc ccagttacat 480
tttatgcac tcgtagcttt ccttccttag caaaattgca ttcaaaaatg tgtaaaaatg 540
agtaataaca gaatatcact acagagactt gnatcctcan ggtaaatgga ttccacattg 600
nagaataaac agcaaanngt cttaggtttt caagtgaaaa ctttttgggt aatcacaaaa 660
atacctggac acataccagg ctttaaa                                     687

```

```

<210> 28
<211> 1529
<212> DNA
<213> Homo sapiens

```

<220>
 <221> misc_feature
 <222> (1)...(1529)
 <223> n = A,T,C or G

<400> 28
 gagatcatcg attttaggtg ctgcntaagt attactgatg tgccttaca tgcattagga 60
 aaaaactrcm cmtttwtwga gtgtgtogac ttttcagcta ctacaggtatc tgacagtggt 120
 gtgattgcac ttgttagtgg acctgtgcg aagaaattag aggagattca tatgggacat 180
 tgtgtaaatc tgaactgatgg ggctgtcgaa gctgtcctta ctactgttcc tcaaatacgt 240
 atattactct tccactggatg ccccttgata acagatcatt cccgagaagt gttggagcaa 300
 ttagtagggc caaacaact aaagcaagtg acatggactg tttattgatg cttttttgaa 360
 gatgatcaat gctaggaag cttatcaaaa ctactttccc aggaaacctat ctatagagat 420
 ttgcattcta cttaagtta acactatttt taattatttt attgtcttaa gttataactc 480
 tcaagagaatt agctaagtct tggatatatac atgggttggg ctttactctt aaacatcttt 540
 aaagtgcctat tattctawaw mtgttggatg agtcattatt tttgaaatga taactcctagc 600
 atgaactctg atctatgggt ttgattcttg tttcttaaat aactttaaaa ttaactgttt 660
 tcccttgaga tttcctctc ctatgtagggt atttgagcta ttgttctaa gttactctga 720
 agtataaac ctggggagaat ctaagtaaac atatttctaa aagcatagtt aoccttctat 780
 ttctggctct ttactctctt ggagtattta aatgcocatt tgccaaaagc agacctgaac 840
 atcaagcctg gtttaattctt caaagaattt agggkattkg tttcmccgga aatgragtga 900
 cttattagcc attcagogggt attagkawta cagaggctct tgcccagcca catccantyc 960
 attgattttt awggctactc ttcccagtta cattttatgc atctgtaagc tttcttccct 1020
 tagcaaaatt gcaattcaaaa atgtgtaaaa atgagtaaat acagaataat actacagaga 1080
 ctgttatcct caggttttatt gatttcacat tgtgaaataa acagcaaaagc ttttagtttt 1140
 caagtgaaaa ctttttggta atcacaaaaa taactgcac ataccaogct ttaaaccaac 1200
 ccccaattt agcatatttca ttttgccatg agccagctct gagattttct ttaaaagattt 1260
 cttattttgc ctctgatgta gtgaaaaaac gggtaagtat gctaactttc ttgtatatgt 1320
 tgggggggtac ttattcaact ccatttcttg tcttacaag atttataaat gtggtatgtt 1380
 tatagtgtgg atatatatgt tgccactgca aaggtggtgc atatgtatat atgtgcaaaa 1440
 tgggtaaggc ctgtttctaac tatgaaattt ttctaaagac aaattcaata aaatttaata 1500
 ctgaattatt aamcaagta aaaaaaaa 1529

<210> 29
 <211> 697
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <222> (1)...(697)
 <223> n = A,T,C or G

<400> 29
 aaaaagagaa gaaagacaag aaaaagaaaa aaaaagaaaa cacctttgtc tttgtacacg 60
 tcaacngggc tcccagagaa atgttctctc tctttttgtt ggcattgggca ctgtgggac 120
 tggngcattc cggctgacac tctcgtttat ttggactgta agtctgaacct ctatgaataa 180
 ttaacttcacg ccttgattgc tcccgtgcca agctccttgg ccaaaactttc accttagctt 240
 ctggttaagtc ttgggccaag ctaagcagca tctatcaatc atcccttcag ctctctgatt 300
 gtctctgggc aaaggcctgg gccaaagctga gccacacgtt tttcaagaca gccctgtgaa 360
 taggcacatt tcttctcctt cccagctcctt aaaaaccctg gaccagacct cgtagagggc 420
 accaacttca gacacctata tctgtctgga aagagctttc tctcttggct tcttaaaact 480
 tcaactcaac ctcaaccttg ngtttaacct ccttaattct cttagaggta gaacaaagaa 540
 ctctggatgg tatctcagac tacgagagac tgggtacatc ttgngcactg ctgagactat 600
 gacacttggg ttctttggag ttggactaaa tattttacat ggagggaagt aatcacagct 660
 tctnttttga ctggcntaat ttacttaacn aaaaag 697

<210> 30
 <211> 1165
 <212> DNA
 <213> Homo sapiens
 <220>
 <221> misc_feature
 <222> (1)...(1165)
 <223> n = A,T,C or G

```

<400> 30
aatgctaagt ccaaagtggt taagtgaact gcccaagctc tacaatgccc toctgaactc 60
ggatgtcttc atttcctgtg ccagactctt aaaaaaata aaaaataata aaaaaagaaa 120
gtacatctaa aaaagaaaga aagacaagaa aaagaaaaaa aaagaaaca cotttgtctt 180
tgtacagtca gtgggctccc aggaaaatgt tcctctcttt ttgttggca tgggcactgt 240
gggatctggt gcattccggt cgacactctc gtttatttgg actgtaagtc tgacctctat 300
gaataatfac ttcagccctt gattgctccc gtgccaagct ccttggccaa actttcaact 360
tagcttctgr taagtcttgg gccaaagctaa gcagcatcta tcaatcatcc ctccagctcc 420
tgattgrtcc ygggccaaag gcctgggcca aagctgagcc acacgttttt caagacagcc 480
tgtgaactag gcacatatcc ttcccttccc agtcoataaa aaccctggac ccagcctcgt 540
agaggcaacca ctttcagaca cctatctctg ctggcaaaaga gcttctctt ctgtctctt 600
aaactttcac tocaacotca cctttgtgtt yacrtctctt aatctcctta gaggtagaac 660
aaagaactct ggaatgttat tcagactacg agagactggt acatcttgtt gcactgctga 720
gactaygaca cttgtgtttc ttgagtttga ctaaatattt tacatgagtg taattawtac 780
agctttcctt tttagactgtc ttattttact taacagaatg ttttgaagga tttgtccyta 840
ttgttagtac ttttcaagat ttcoattatt ttaaggstgr atgctatccc acgtggattg 900
tacgtgccct gtttgcgtga tctactcatc ctttaagggtta catttgcttc caggtaacat 960
gtttgtgact aatactacaa atgtgcatat atctattcca tgttctgctt tggctctgtt 1020
ggggatattt ttccatacac tggattcagt accatggtgg taatcccttt gctnttgggt 1080
gnctccaatc cgggtgtagt gnaaggctcc ccccaaaatt aattggccca cggaccaagg 1140
tggccaanga aggcctcnac cccctt                                     1165

```

<210> 31
 <211> 557
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <222> (1)...(557)
 <223> n = A,T,C or G

```

<400> 31
cgcttagggc cctcgccggg ggcttgtggg tctctctccc cctcccaactg acaactgccc 60
caactgcctt tccgcgcccg gtcacagtga aaatgtagac ggggtcgttg tccgtacgac 120
tgtgcgccag ggctcgggga gggcgccctt ccgcgtgagc gcccccctgg gaattattga 180
cataatcacc tctcattcca gactatgtta ggtcttaatg gtgggaggac gcccgagtgc 240
tcggcccggtt tcaccccgag gaggaaggac actgggtcat gacgccatca gaggcgccca 300
gagcagggac cggacgcggg ttggagatgt tggactcgct gttggccttg gggcggtggt 360
gctgcttogg gattccgtgg agtggggagg gcgcagtcct ttgaaggcgc ctgtccaaaga 420
aagagagaga agccagagat agcctgatcc tgccttnnag ttcagttctt aaaaacagca 480
ggctctctg cggmctaggg canggcaggc taccagccac atctctatg agccagatgc 540
ttatgatgac ctgggacc                                     557

```

<210> 32
 <211> 527
 <212> DNA
 <213> Homo sapiens

```

<220>
<221> misc_feature
<222> (1)...(527)
<223> n = A,T,C or G

<400> 32
atccaggagagg agggagtctat ctccotcaagn ttgacaactc ctactottttg tggcggncaa 60
aatcagtccta ctacagagtc tattatacta gataaaaatg tnggtacaaa gtctgggagtc 120
taggggtggg cagaagatga catttaattt ggaaatttct ttttactttt gtggagcatt 180
agagtcacag tttaccttat tgatattggt ctgatggntt gtgaactcct gctgggaatc 240
aaaatttcct tgagactcct tagcattcat actttggggn taaaggagat tncctcagact 300
catccagccc ttgggtgctg accagcagag tccactagnng atgctgaagt tacatgagct 360
acatgtttaa tatttaaagt ctccaaaata aaacacccca acgttgacct taccgggctt 420
gatgggttagc ccctttgctg gctgctccat gtgccttatg agagcccgta agttacaggt 480
gtcctctaata ttgaaatcca taagntaaca ngctatatc agntgcn 527

<210> 33
<211> 934
<212> DNA
<213> Homo sapiens

<400> 33
gtaggccagc gatgacgacg agggaggaaga aggaacacatc ggttgtgaag agaaagccaa 60
aaggaatgcc aacaagccct tgctggatga gatgtgcctt gtgtccgacg ggaactgtcat 120
gaggtgtgtg atgctggcag ccactcaatat ccaaggagaga ggagtctatc tccctcaagtt 180
tgacaactcc tactctttgt ggcggtcaaa atcagctctac tacrgagtctc attatactag 240
ataaaaatgt tgtttacaaag tctggagctc wgggttgggc agaagatgac atttaatttg 300
gaaatttcct tttacttttg tggagcatta gagtcacagt ttaccttatt gattattggtc 360
tgatgtgttg tgaactcttg ctgggaatca aaatttccct gagactcttt agcattcata 420
ctttgggggtt aaaggagatt cctcagactc atccagccct tgggtgtctga ccagcagagt 480
cactagtggg tgctgaagtt acatgagcta catgttaaat atttaaaagtc tccaaaaataa 540
aacaccccaa cgttgacctt acccggtcga tgggttagccc cttgctgcct gctccatgtg 600
tcttatgaga gcccgtagtt acagtgctct ctaatttgaa atccataagt taacaagctc 660
atatcaggtg cagctggcct tgattaaagg ccatttttaa aacttaaaaa ctcaacacct 720
cacagattat aatagaaaaa mgaatggggc ctacagtttga tctccgttca gaatgaccca 780
gattgtttct gctttggggt gcagctgctt aagttcagag ttatattaca gagaattatt 840
tlyctggaga taactcttaa acctagaatg ktcaaaacc waattggata attggaagta 900
tccaagatac gtagaacacc cccggagaat tttc 934

<210> 34
<211> 758
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<222> (1)...(758)
<223> n = A,T,C or G

<400> 34
ggctttatag cccatcctca ttgcttactg ccacccctca gctggggctc aaggcagtac 60
tattcagttt attcaccaga cctgcctcca gacactctac tctttcaaaa attagtgttt 120
tccatcaagg agcatgttcc agagcatttc ccagagatgt cccaagaagc actgtccggt 180
gctgtggcgt acagttggca cagcatttaga ctaagtgga cctccagca ggtcgtttta 240
gaatccgctc atttgactag atacgatgta attggtgtc tttaaaaaac gcgcacacac 300
acacaactcg ataggcatat ctcatgccca tccaatatg aatgttctc gcttgctgaa 360
tttaagcctg tattttaagg ttttgtgtt cctcgccac aatgggtgat gtcactgata 420

```

19

```

gaacgaagct gagtttccaa gggtttgggg ctgtgcaaga gtaaacacta gagcttgagt 480
tggtatccag ctggcaagca cggaagtcct tgaagaatgt aatgtaaaaa gggaaaaaga 540
tgtaaagctt ttgtaccaa atgagagtgt gagccagcc acaaaatgct tttccctgtg 600
taaaagtctc tctggaaggg acattccatc tccatgggtc actctgaggg goactgtcaa 660
ctagagattg gcccatcca ggtgggagga accccttttg gatgngnagt atncaatctg 720
ctgngcattt tgacaggatc tctgaatggc taggtaat 758

```

```

<210> 35
<211> 1534
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> misc_feature
<222> (1)...(1534)
<223> n = A,T,C or G

```

```

<400> 35
ngaggttaaaa ggcaaggcag catttaataa gtacctgttg tatcctttta agtgtttgtt 60
gtggaatacc tcacaagac cggaagtcct tgaagaatgt aatgtaaaaa gggaaaaaga 120
tgaggaattt tgcttttcaa gtgcatatac actattaata ttttttcccc aagaggagca 180
ttctaagcta atttatgcag tgtgactgta ttaagcatta agcttccctc agagctggcc 240
tatcgaggat gctactgccc tctctacaga tgtgtctgaa atgctgccc aaggatggcc 300
cttagccagt taacagcttt atagccatc ctcatgtctt actgccccc ctccagctgg 360
gtccaaggca gtactattca gtttattcac cagacctgcc ccagacatc tacttcttcc 420
aaaaattagt gttttccatc aaggagcatg ttccagagca ttccocagag atgtccocaa 480
gaacactgtc cgtgtctgtg gcgtacagt gcaacagcat tagactaagt ggaacatccc 540
agcaggctgc tttagaatcc gctcaattga ctagatagca tgaattggc tgtcttttaa 600
aaacggcgca cacacacaca atctgatagg gcataatcca tgccattcca atatggaatg 660
ttcttcgctt gctgaattta agcctgtatt ttaaggtttt gtggttccct ggccacaatg 720
gggtgatgtc actgatagaa cgaagctgag ttccaaaggg ttgggggtg ttgaggagt 780
aaacactaga gcttgagtgt ttatccagct ggcaagcagg gaagtctttg aagaatgtaa 840
tgtaaaaagg gaaaagaatg taaagctttt tgtaccaaat gagagttgga gccagccaa 900
caaatgcttt tccctgtgta aaagtctctc tggaaaggac attccatctc catggtgca 960
tctgaggggc actgtcaact agagattggc cccatccagg tgggaggaa ccccttgrrr 1020
tggtgagtat ccaatctgct gtgcaattga caggatctct gaatggctag gtaatggatc 1080
ccaagcaggc tcacaaattt aaatgagggc tttgtgtgca gaaaggagaa taagtacaga 1140
ttattttcct accactagat ttttggggag agtcaoccat gaatgttgac aattacttaa 1200
aatattttta gctcccttgc tgaattcctg tctgtccctc gaggaaatcg atgtgcatc 1260
agccataggc accacaccca aatttcccta ggagttggag taatgctaga atgtgaagacc 1320
ttctgagtta agggctctct tgccttccca gaggcaggag aattttgcac tgggtgtggt 1380
aaatgtataa aaagctatat gtccaccagt ttactcattt ccaatgtgta atgaataaa 1440
atgtagtgtg caaattattt gaaaatccca gaagggaagt acttttccaa tacagttatt 1500
tttttaagca ataaacttac gattttttca gcaa 1534

```

```

<210> 36
<211> 125
<212> PRT
<213> Homo sapiens

```

```

<220>
<221> variant
<222> (1)...(125)
<223> Xaa = Any amino acid

```

```

<400> 36
Leu Ser Ser Arg Gly Met Iys Ala Val Leu Leu Ala Asp Thr Glu Ile
          5                      10                      15

```

Asp Leu Phe Ser Thr Asp Ile Pro Pro Thr Asn Ala Val Asp Phe Thr
 10 20 25 30

Gly Arg Cys Tyr Phe Thr Lys Ile Cys Lys Cys Lys Leu Lys Asp Ile
35 40 45

Ala Cys Leu Lys Cys Gly Asn Ile Val Xaa Tyr His Val Ile Val Pro
50 55 60

Cys Ser Ser Cys Leu Leu Ser Cys Asn Asn Arg His Phe Trp Met Phe
65 70 75 80

His Ser Gln Ala Val Tyr Asp Ile Asn Arg Leu Asp Ser Thr Gly Val
85 90 95

Asn Val Leu Leu Arg Gly Asn Leu Pro Glu Ile Glu Glu Ser Thr Asp
100 105 110

Glu Asp Val Leu Asn Ile Ser Ala Glu Glu Cys Ile Arg
115 120 125

<210> 37

<211> 448

<212> PRT

<213> Homo sapiens

 $\langle 220 \rangle$

<221> VARIANT

<222> (1) ... (448)

<223> Xaa = any amino acid

<400> 37

Met Ser Arg Arg Pro Cys Ser Cys Ala Leu Arg Pro Pro Arg Cys Ser
5 10 15

Cys Ser Ala Ser Pro Ser Ala Val Thr Ala Ala Gly Arg Pro Arg Pro
20 25 30

Ser Asp Ser Cys Lys Glu Glu Ser Ser Thr Leu Ser Val Lys Met Lys
35 40 45

Cys Asp Phe Asn Cys Asn His Val His Ser Gly Leu Lys Leu Val Lys
50 55 60

Pro Asp Asp Ile Gly Arg Leu Val Ser Tyr Thr Pro Ala Tyr Leu Glu
65 70 75 80

Gly Ser Cys Lys Asp Cys Ile Lys Asp Tyr Glu Arg Leu Ser Cys Ile
85 90 95

Gly Ser Pro Ile Val Ser Pro Arg Ile Val Gln Leu Glu Thr Glu Ser
100 105 110

Lys Arg Leu His Asn Lys Glu Asn Gln His Val Gln Gln Thr Leu Asn
115 120 125

Ser Thr Asn Glu Ile Glu Ala Leu Glu Thr Ser Arg Leu Tyr Glu Asp
 130 135 140
 Ser Gly Tyr Ser Ser Phe Ser Leu Gln Ser Gly Leu Ser Glu His Glu
 145 150 155 160
 Glu Gly Ser Leu Leu Glu Glu Asn Phe Gly Asp Ser Leu Gln Ser Cys
 165 170 175
 Leu Leu Gln Ile Gln Ser Pro Asp Gln Tyr Pro Asn Lys Asn Leu Leu
 180 185 190
 Pro Val Leu His Phe Glu Lys Val Val Cys Ser Thr Leu Lys Lys Asn
 195 200 205
 Ala Lys Arg Asn Pro Lys Val Asp Arg Glu Met Leu Lys Glu Ile Ile
 210 215 220
 Ala Arg Gly Asn Phe Arg Leu Gln Asn Ile Ile Gly Arg Lys Met Gly
 225 230 235 240
 Leu Glu Cys Val Asp Ile Leu Ser Glu Leu Phe Arg Arg Gly Leu Arg
 245 250 255
 His Val Leu Ala Thr Ile Leu Ala Gln Leu Ser Ser Asp Met Asp Leu Ile
 260 265 270
 Asn Val Ser Lys Val Ser Thr Thr Trp Lys Lys Ile Leu Glu Asp Asp
 275 280 285
 Lys Gly Ala Phe Gln Leu Tyr Ser Lys Ala Ile Gln Arg Val Thr Glu
 290 295 300
 Asn Asn Asn Lys Phe Ser Pro His Ala Ser Thr Arg Glu Tyr Val Met
 305 310 315 320
 Phe Arg Thr Pro Leu Ala Ser Val Gln Lys Ser Ala Ala Gln Thr Ser
 325 330 335
 Leu Lys Lys Asp Ala Gln Thr Lys Leu Ser Asn Gln Gly Asp Gln Lys
 340 345 350
 Gly Ser Thr Tyr Ser Arg His Asn Glu Phe Ser Glu Val Ala Lys Thr
 355 360 365
 Leu Lys Lys Asn Glu Ser Leu Lys Ala Cys Ile Arg Cys Asn Ser Pro
 370 375 380
 Ala Lys Tyr Asp Cys Tyr Leu Gln Arg Ala Thr Cys Lys Arg Glu Gly
 385 390 395 400
 Cys Gly Phe Asp Tyr Cys Thr Lys Cys Leu Cys Asn Tyr His Thr Thr
 405 410 415
 Lys Asp Cys Ser Asp Gly Lys Leu Leu Lys Ala Ser Cys Lys Ile Gly
 420 425 430
 Pro Leu Pro Gly Thr Lys Lys Ser Lys Lys Asn Leu Arg Arg Leu Xaa

435

440

445

```
<210> 38
<211> 1050
<212> PRT
<213> Homo sapiens
```

400> 38																	
Met	Ala	Ala	Val	Lys	Lys	Glu	Gly	Gly	Ala	Leu	Ser	Glu	Ala	Met	Ser		
				5					10					15			
Leu	Glu	Gly	Asp	Glu	Trp	Glu	Leu	Ser	Lys	Glu	Asn	Val	Gln	Pro	Leu		
			20					25					30				
Arg	Gln	Gly	Arg	Ile	Met	Ser	Thr	Leu	Gln	Gly	Ala	Leu	Ala	Gln	Glu		
		35					40					45					
Ser	Ala	Cys	Asn	Asn	Thr	Leu	Gln	Gln	Gln	Lys	Arg	Ala	Phe	Glu	Tyr		
	50					55					60						
Glu	Ile	Arg	Phe	Tyr	Thr	Gly	Asn	Asp	Pro	Leu	Asp	Val	Trp	Asp	Arg		
					70					75					80		
Tyr	Ile	Ser	Trp	Thr	Glu	Gln	Asn	Tyr	Pro	Gln	Gly	Gly	Lys	Glu	Ser		
				85					90				95				
Asn	Met	Ser	Thr	Leu	Leu	Glu	Arg	Ala	Val	Glu	Ala	Leu	Gln	Gly	Glu		
			100					105					110				
Lys	Arg	Tyr	Tyr	Ser	Asp	Pro	Arg	Phe	Leu	Asn	Leu	Trp	Leu	Lys	Leu		
		115					120					125					
Gly	Arg	Leu	Cys	Asn	Glu	Pro	Leu	Asp	Met	Tyr	Ser	Tyr	Leu	His	Asn		
		130				135					140						
Gln	Gly	Ile	Gly	Val	Ser	Leu	Ala	Gln	Phe	Tyr	Ile	Ser	Trp	Ala	Glu		
	145				150					155					160		
Glu	Tyr	Glu	Ala	Arg	Glu	Asn	Phe	Arg	Lys	Ala	Asp	Ala	Ile	Phe	Gln		
			165						170					175			
Glu	Gly	Ile	Gln	Gln	Lys	Ala	Glu	Pro	Leu	Glu	Arg	Leu	Gln	Ser	Gln		
		180						185					190				
His	Arg	Gln	Phe	Gln	Ala	Arg	Val	Ser	Arg	Gln	Thr	Leu	Leu	Ala	Leu		
		195					200					205					
Glu	Lys	Glu	Glu	Glu	Glu	Glu	Val	Phe	Glu	Ser	Ser	Val	Pro	Gln	Arg		
	210					215					220						
Ser	Thr	Leu	Ala	Glu	Leu	Lys	Ser	Lys	Gly	Lys	Lys	Thr	Ala	Arg	Ala		
	225				230				235					240			
Pro	Ile	Ile	Arg	Val	Gly	Gly	Ala	Leu	Lys	Ala	Pro	Ser	Gln	Asn	Arg		
			245					250						255			
Gly	Leu	Gln	Asn	Pro	Phe	Pro	Gln	Gln	Met	Gln	Asn	Asn	Ser	Arg	Ile		

260	265	270
Thr Val Phe Asp Glu Asn Ala Asp	Glu Ala Ser Thr	Ala Glu Leu Ser
275	280	285
Lys Pro Thr Val Gln Pro Trp Ile	Ala Pro Pro Met	Pro Arg Ala Lys
290	295	300
Glu Asn Glu Leu Gln Ala Gly Pro	Trp Asn Thr Gly Arg Ser	Leu Glu
305	310	315
His Arg Pro Arg Gly Asn Thr Ala	Ser Leu Ile Ala Val	Pro Ala Val
325	330	335
Leu Pro Ser Phe Thr Pro Tyr Val	Glu Glu Thr Ala Gln Gln	Pro Val
340	345	350
Met Thr Pro Cys Lys Ile Glu Pro	Ser Ile Asn His Ile Leu Ser Thr	
355	360	365
Arg Lys Pro Gly Lys Glu Glu Gly	Asp Pro Leu Gln Arg Val	Gln Ser
370	375	380
His Gln Gln Ala Ser Glu Glu Lys	Lys Glu Lys Met Met Tyr Cys Lys	
385	390	395
Glu Lys Ile Tyr Ala Gly Val Gly	Glu Phe Ser Phe Glu Glu Ile Arg	
405	410	415
Ala Glu Val Phe Arg Lys Lys Leu	Lys Glu Gln Arg Glu Ala Glu Leu	
420	425	430
Leu Thr Ser Ala Glu Lys Arg Ala	Glu Met Gln Lys Gln Ile Glu Glu	
435	440	445
Met Glu Lys Lys Leu Lys Glu Ile	Gln Thr Thr Gln Gln Glu Arg Thr	
450	455	460
Gly Asp Gln Gln Glu Glu Thr Met	Pro Thr Lys Glu Thr Thr Lys Leu	
465	470	475
Gln Ile Ala Ser Glu Ser Gln Lys	Ile Pro Gly Met Thr Leu Ser Ser	
485	490	495
Ser Val Cys Gln Val Asn Cys Cys	Ala Arg Glu Thr Ser Leu Ala Glu	
500	505	510
Asn Ile Trp Gln Glu Gln Pro His	Ser Lys Gly Pro Ser Val Pro Phe	
515	520	525
Ser Ile Phe Asp Glu Phe Leu Leu	Ser Glu Lys Lys Asn Lys Ser Pro	
530	535	540
Pro Ala Asp Pro Pro Arg Val Leu	Ala Gln Arg Arg Pro Leu Ala Val	
545	550	555
Leu Lys Thr Ser Glu Ser Ile Thr	Ser Asn Glu Asp Val Ser Pro Asp	
565	570	575

Val Cys Asp Glu Phe Thr Gly Ile Glu Pro Leu Ser Glu Asp Ala Ile
 580 585 590
 Ile Thr Gly Phe Arg Asn Val Thr Ile Cys Pro Asn Pro Glu Asp Thr
 595 600 605
 Cys Asp Phe Ala Arg Ala Ala Arg Phe Val Ser Thr Pro Phe His Glu
 610 615 620
 Ile Met Ser Leu Lys Asp Leu Pro Ser Asp Pro Glu Arg Leu Leu Pro
 625 630 635 640
 Glu Glu Asp Leu Asp Val Lys Thr Ser Glu Asp Gln Gln Thr Ala Cys
 645 650 655
 Gly Thr Ile Tyr Ser Gln Thr Leu Ser Ile Lys Lys Leu Ser Pro Ile
 660 665 670
 Ile Glu Asp Ser Arg Glu Ala Thr His Ser Ser Gly Phe Ser Gly Ser
 675 680 685
 Ser Ala Ser Val Ala Ser Thr Ser Ser Ile Lys Cys Leu Gln Ile Pro
 690 695 700
 Glu Lys Leu Glu Leu Thr Asn Glu Thr Ser Glu Asn Pro Thr Gln Ser
 705 710 715 720
 Pro Trp Cys Ser Gln Tyr Arg Arg Gln Leu Leu Lys Ser Leu Pro Glu
 725 730 735
 Leu Ser Ala Ser Ala Glu Leu Cys Ile Glu Asp Arg Pro Met Pro Lys
 740 745 750
 Leu Glu Ile Glu Lys Glu Ile Glu Leu Gly Asn Glu Asp Tyr Cys Ile
 755 760 765
 Lys Arg Glu Tyr Leu Ile Cys Glu Asp Tyr Lys Leu Phe Trp Val Ala
 770 775 780
 Pro Arg Asn Phe Ala Glu Leu Thr Val Ile Lys Val Ser Ser Gln Pro
 785 790 795 800
 Val Pro Trp Asp Phe Tyr Ile Asn Leu Lys Leu Lys Glu Arg Leu Asn
 805 810 815
 Glu Asp Phe Asp His Phe Cys Ser Cys Tyr Gln Tyr Gln Asp Gly Cys
 820 825 830
 Ile Val Trp His Gln Tyr Ile Asn Cys Phe Thr Leu Gln Asp Leu Leu
 835 840 845
 Gln His Ser Glu Tyr Ile Thr His Glu Ile Thr Val Leu Ile Ile Tyr
 850 855 860
 Asn Leu Leu Thr Ile Val Glu Met Leu His Lys Ala Glu Ile Val His
 865 870 875 880

Gly Asp Leu Ser Pro Arg Cys Leu Ile Leu Arg Asn Arg Ile His Asp
 885 890 895
 Pro Tyr Asp Cys Asn Lys Asn Asn Gln Ala Leu Lys Ile Val Asp Phe
 900 905 910
 Ser Tyr Ser Val Asp Leu Arg Val Gln Leu Asp Val Phe Thr Leu Ser
 915 920 925
 Gly Phe Arg Thr Val Gln Ile Leu Glu Gly Gln Lys Ile Leu Ala Asn
 930 935 940
 Cys Ser Ser Pro Tyr Gln Val Asp Leu Phe Gly Ile Ala Asp Leu Ala
 945 950 955 960
 His Leu Leu Leu Phe Lys Glu His Leu Gln Val Phe Trp Asp Gly Ser
 965 970 975
 Phe Trp Lys Leu Ser Gln Asn Ile Ser Glu Leu Lys Asp Gly Glu Leu
 980 985 990
 Trp Asn Lys Phe Phe Val Arg Ile Leu Asn Ala Asn Asp Glu Ala Thr
 995 1000 1005
 Val Ser Val Leu Leu Gly Glu Leu Ala Ala Glu Met Asn Gly Val Phe Asp
 1010 1015 1020
 Thr Thr Phe Gln Ser His Leu Asn Lys Ala Leu Trp Lys Val Gly Lys
 1025 1030 1035 1040
 Leu Thr Ser Pro Gly Ala Leu Leu Phe Gln
 1045 1050

 <210> 39
 <211> 258
 <212> PRT
 <213> Homo sapiens

 <400> 39
 Gly Lys Leu Thr Gly Ile Ser Asp Pro Val Thr Val Lys Thr Ser Gly
 5 10 15
 Ser Arg Phe Gly Ser Trp Met Thr Asp Pro Leu Ala Pro Glu Gly Asp
 20 25 30
 Asn Arg Val Trp Tyr Met Asp Gly Tyr His Asn Asn Arg Phe Val Arg
 35 40 45
 Glu Tyr Lys Ser Met Val Asp Phe Met Asn Thr Asp Asn Phe Thr Ser
 50 55 60
 His Arg Leu Pro His Pro Trp Ser Gly Thr Gly Gln Val Val Tyr Asn
 65 70 75 80
 Gly Ser Ile Tyr Phe Asn Lys Phe Gln Ser His Ile Ile Ile Arg Phe
 85 90 95

Asp Leu Lys Thr Glu Thr Ile Leu Lys Thr Arg Ser Leu Asp Tyr Ala
 100 105 110
 Gly Tyr Asn Asn Met Tyr His Tyr Ala Trp Gly Gly His Ser Asp Ile
 115 120 125
 Asp Leu Met Val Asp Glu Ser Gly Leu Trp Ala Val Tyr Ala Thr Asn
 130 135 140
 Gln Asn Ala Gly Asn Ile Val Val Ser Arg Leu Asp Pro Val Ser Leu
 145 150 155 160
 Gln Thr Leu Gln Thr Trp Asn Thr Ser Tyr Pro Lys Arg Ser Ala Gly
 165 170 175
 Glu Ala Phe Ile Ile Cys Gly Thr Leu Tyr Val Thr Asn Gly Tyr Ser
 180 185 190
 Gly Gly Thr Lys Val His Tyr Ala Tyr Gln Thr Asn Ala Ser Thr Tyr
 195 200 205
 Glu Tyr Ile Asp Ile Pro Phe Gln Asn Lys Tyr Ser His Ile Ser Met
 210 215 220
 Leu Asp Tyr Asn Pro Lys Asp Arg Ala Leu Tyr Ala Trp Asn Asn Gly
 225 230 235 240
 His Gln Ile Leu Tyr Asn Val Thr Leu Phe His Val Ile Arg Ser Asp
 245 250 255
 Glu Leu

<210> 40
 <211> 324
 <212> PRT
 <213> Homo sapiens

<400> 40
 Met Asp Ala Pro Arg Gln Val Val Asn Phe Gly Pro Gly Pro Ala Lys
 5 10 15
 Leu Pro His Ser Val Leu Leu Glu Ile Gln Lys Glu Leu Leu Asp Tyr
 20 25 30
 Lys Gly Val Gly Ile Ser Val Leu Glu Met Ser His Arg Ser Ser Asp
 35 40 45
 Phe Ala Lys Ile Ile Asn Asn Thr Glu Asn Leu Val Arg Glu Leu Leu
 50 55 60
 Ala Val Pro Asp Asn Tyr Lys Val Ile Phe Leu Gln Gly Gly Gly Cys
 65 70 75 80
 Gly Gln Phe Ser Ala Val Pro Leu Asn Leu Ile Gly Leu Lys Ala Gly
 85 90 95

Arg Cys Ala Asp Tyr Val Val Thr Gly Ala Trp Ser Ala Lys Ala Ala
 100 105 110
 Glu Glu Ala Lys Lys Phe Gly Thr Ile Asn Ile Val His Pro Lys Leu
 115 120 125
 Gly Ser Tyr Thr Lys Ile Pro Asp Pro Ser Thr Trp Asn Leu Asn Pro
 130 135 140
 Asp Ala Ser Tyr Val Tyr Tyr Cys Ala Asn Glu Thr Val His Gly Val
 145 150 155 160
 Glu Phe Asp Phe Ile Pro Asp Val Lys Gly Ala Val Leu Val Cys Asp
 165 170 175
 Met Ser Ser Asn Phe Leu Ser Lys Pro Val Asp Val Ser Lys Phe Gly
 180 185 190
 Val Ile Phe Ala Gly Ala Gln Lys Asn Val Gly Ser Ala Gly Val Thr
 195 200 205
 Val Val Ile Val Arg Asp Asp Leu Leu Gly Phe Ala Leu Arg Glu Cys
 210 215 220
 Pro Ser Val Leu Glu Tyr Lys Val Gln Ala Gly Asn Ser Ser Leu Tyr
 225 230 235 240
 Asn Thr Pro Pro Cys Phe Ser Ile Tyr Val Met Gly Leu Val Leu Glu
 245 250 255
 Trp Ile Lys Asn Asn Gly Gly Ala Ala Ala Met Glu Lys Leu Ser Ser
 260 265 270
 Ile Lys Ser Gln Thr Ile Tyr Glu Ile Ile Asp Asn Ser Gln Gly Phe
 275 280 285
 Tyr Val Ser Val Gly Gly Ile Arg Ala Ser Leu Tyr Asn Ala Val Thr
 290 295 300
 Ile Glu Asp Val Gln Lys Leu Ala Ala Phe Met Lys Lys Phe Leu Glu
 305 310 315 320
 Met His Gln Leu

<210> 41

<211> 410

<212> PRT

<213> Homo sapiens

<400> 41

Met Glu Ala Glu Asn Ala Gly Ser Tyr Ser Leu Gln Gln Ala Gln Ala
 5 10 15

Phe Tyr Thr Phe Pro Phe Gln Gln Leu Met Ala Glu Ala Pro Asn Met
 20 25 30

Ala Val Val Asn Glu Gln Gln Met Pro Glu Glu Val Pro Ala Pro Ala
 35 40 45
 Pro Ala Gln Glu Pro Val Gln Glu Ala Pro Lys Gly Arg Lys Arg Lys
 50 55 60
 Pro Arg Thr Thr Glu Pro Lys Gln Pro Val Glu Pro Lys Lys Pro Val
 65 70 75 80
 Glu Ser Lys Lys Ser Gly Lys Ser Ala Lys Pro Lys Glu Lys Gln Glu
 85 90 95
 Lys Ile Thr Asp Thr Phe Lys Val Lys Arg Lys Val Asp Arg Phe Asn
 100 105 110
 Gly Val Ser Glu Ala Glu Leu Leu Thr Lys Thr Leu Pro Asp Ile Leu
 115 120 125
 Thr Phe Asn Leu Asp Ile Val Ile Ile Gly Ile Asn Pro Gly Leu Met
 130 135 140
 Ala Ala Tyr Lys Gly His His Tyr Pro Gly Pro Gly Asn His Phe Trp
 145 150 155 160
 Lys Cys Leu Phe Met Ser Gly Leu Ser Glu Val Gln Leu Asn His Met
 165 170 175
 Asp Asp His Thr Leu Pro Gly Lys Tyr Gly Ile Gly Phe Thr Asn Met
 180 185 190
 Val Glu Arg Thr Thr Pro Gly Ser Lys Asp Leu Ser Ser Lys Glu Phe
 195 200 205
 Arg Glu Gly Gly Arg Ile Leu Val Gln Lys Leu Gln Lys Tyr Gln Pro
 210 215 220
 Arg Ile Ala Val Phe Asn Gly Lys Cys Ile Tyr Glu Ile Phe Ser Lys
 225 230 235 240
 Glu Val Phe Gly Val Lys Val Lys Asn Leu Glu Phe Gly Leu Gln Pro
 245 250 255
 His Lys Ile Pro Asp Thr Glu Thr Leu Cys Tyr Val Met Pro Ser Ser
 260 265 270
 Ser Ala Arg Cys Ala Gln Phe Pro Arg Ala Gln Asp Lys Val His Tyr
 275 280 285
 Tyr Ile Lys Leu Lys Asp Leu Arg Asp Gln Leu Lys Gly Ile Glu Arg
 290 295 300
 Asn Met Asp Val Gln Glu Val Gln Tyr Thr Phe Asp Leu Gln Leu Ala
 305 310 315 320
 Gln Glu Asp Ala Lys Lys Met Ala Val Lys Glu Glu Lys Tyr Asp Pro
 325 330 335
 Gly Tyr Glu Ala Ala Tyr Gly Gly Ala Tyr Gly Glu Asn Pro Cys Ser

340 345 350

Ser Glu Pro Cys Gly Phe Ser Ser Asn Gly Leu Ile Glu Ser Val Glu
355 360

Leu Arg Gly Glu Ser Ala Phe Ser Gly Ile Pro Asn Gly Gln Trp Met
370 375 380

Thr Gln Ser Phe Thr Asp Gln Ile Pro Ser Phe Ser Asn His Cys Gly
385 390 395 400

Thr Gln Glu Gln Glu Glu Ser His Ala
405 410

<210> 42
<211> 484
<212> DNA
<213> Homo sapiens

<400> 42
ttcacgtaag acttttttgggt ttgatcatct ttgttgaggt aggaactatca gttccctcta 60
aatgtatatg ttgattttatg agtaattgtt atttattctt tattttattta tattaattat 120
gaagattatg atattattttg attgcagatt tttttggcgc gctgccccct ccccaacctg 180
ccacttttga cattccactg tgcgttttag aagagagcct ttttctaagg ggaatcgtctt 240
aagattttta cttttatacc tatctgagtg aattacagac aacctatcat ttattctgct 300
tcagggttcc ccagggcctc tgtacaaccg acagctctta cttttaaatg caatctcttt 360
tctacataca ttatttttctt aattgttagc tatttataga aagcttcaat agaactgttt 420
caactgtata actatttact attcaataaa aatattttca aagtcacaaa aaaaaaaaaa 480
aaag 484

<210> 43
<211> 700
<212> DNA
<213> Homo sapiens

<400> 43
ctcacagta attccactcc catgaaactt tggctattgt tatgcattaa gtggggctta 60
tcttttggtt ggagttcatt tgaactcttg aaccttagtt tagtgaagat gaactgtctg 120
tctttaggta gaaacggtgt ttatttaaaa atcagtttta aaaaatgagc taccatattg 180
gctgtctatt ataaatggga caccacaaac aattttctat tacagtgttg tacttgcaaa 240
cattttgcta tacagtaact catagatgca tacaaatgag ctcaacttatt acaaaagacaa 300
acgtttaatt tgctaaatat tttaacaagt ttgttatata ttttatttaa tttaaaaaa 360
atctcttacc aacctacata tttattacta taatttgcta tgaactcag ttaatttatt 420
tgtgtttgca tagtttgagc aggatgtttt gtgaagtatg ttgttatata ttgtcctact 480
ttgtacttga tgtgttttgt aatgtgcaact gaatttggtt tcttttcaac tatgttaatg 540
atcaatactg taaattgggt cttttgtaaa caaaaaggca atgatgtatg catttttttt 600
aatttgaggt agttttgttg tatactgttt ctccaaacac ttaatatctt ttacatcaaa 660
gcaacaaaat tgtgttcagt gctgtacatt tgggtgatgg 700

<210> 44
<211> 672
<212> DNA
<213> Homo sapiens

<220>
 <221> misc_feature
 <222> (1)...(672)
 <223> n = A,T,C or G

<400> 44
 tttttgttta cataattgta aggaacagta attctagaaa cactagaaga aaaargcata 60
 gcaatgtcca cagttaaaaa aaaaagkgca cactactcgg tcacaatcac agtcattact 120
 tgaaaaacata tatgtaacaa gtatagataaga aatatcactg atgcctcctaaa ctcatgtcta 180
 aaaaactgaat gacataaatt ttacatgaaa taaggcaaat tcaggaatgc acaagaatt 240
 tgtaatccaa ccaaatctaa acaacagaaa aaagtgtgat aagaagcatg aactaaagta 300
 ottctcccta aatattttaa aaataggctt gtctcagtcg acaagaaaaa catcactcat 360
 gtgtatccca cactataaaa taagaaagaa gggtaaaagta tgggggatag gagggcacag 420
 ttcatgttaa gttgcagctg catccgtga gatgtcccta cattattttt agctagaact 480
 gaaaattata caaatcatat caggagatgt aatggctctt ttggaaacta ttctgaaaag 540
 aaatgaaaag aaaactacac acaagagtcg aaattttcag attgtcactt gcaacctctt 600
 aacattcagt catctacatc caggtgctgc tagagggatg cctggagaca gcagcggcaa 660
 tcaggaacga gc 672

<210> 45
 <211> 480
 <212> DNA
 <213> Homo sapiens

<400> 45
 tcagttccat gtatacaatt accagatgcc accgcagtcg cctgttgggg agcaaaaggag 60
 aaatctgtgg accgaagcat acaaatgggtg gtatctgttc tgtttaatcc agagaagaga 120
 ctgataaatt ccgttgttac tcaagatgac tgcttcaagg gtaaaagagt gcactcgttt 180
 agaagaagtt tggcagttat taaatctgtt ggatcctctc agctatctag ttctatggga 240
 agttgctggt ttgtgaattt aagctaaaaa ttttccacta ttacagaaat tctgaatttt 300
 ggtaaatcac actgaaactt tctgtataac ttgtattatt agactctcta tttttatctt 360
 aacactgaaa ctgttcttca ttgatgtttt atttagaacc tggttctgtg ttttaatatat 420
 agtttaaagt aacaataaat cgagactgaa agaattgttaa gatttatctg caaggatttt 480

<210> 46
 <211> 427
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <222> (1)...(427)
 <223> n = A,T,C or G

<400> 46
 tttttaaaaa taagttgtct actattgtat tatatatgta taogaaaact ttaaagctat 60
 ttgaaaaata tgagttctta gctttaatca tgaagtctga agtttgcttt cagtaattat 120
 tttaaaagtt gttttgggtc attgctttat aatattatt attgaatgcc aaacctgttc 180
 ttttttttac tgtgtccaat attctttcaa gcaaatgcaa tggctggaat ataattcaga 240
 attaactgaa acccagccag aagagggacc acctgtaaa caagtctctt caagtctcac 300
 tgcactatcc aaacctgtt acaaaaagag caactgctat attcacatta tgatattttt 360
 ctatcttaaa tttgtcaaaa taaagtatga gtctactaat taaaaaaaaa aaacctctck 420
 tscacaa 427

<210> 47

<211> 581
 <212> DNA
 <213> Homo sapiens

<400> 47
 tcttttgaaa aataaaggat ctaatgtctc cctaataagt ottttttcct tccaaactaaa 60
 tgacctaacac ggaacttttat tttcttgatc aaagagggtg ttattaaggga cttctggata 120
 actatacttt tactctattt ttaaagatca caaagtaatt ttaaattgta acaggttccc 180
 ataccatgaa tgcctggcctc accttctcta tcatccacat ttgaaatgc aaagaaagct 240
 cccttgtaag ccatacttcc ttcccccact ccatactagg atacttgccc agtgctcatt 300
 aggcatttct tattcagata gtccaaattt aggttattat gcttaatttg acacattaac 360
 taaatgcccc gttttaaatt atatccatca attcacgctg aaatgtgctt ctttgtgcta 420
 tcaaatggaa tagaatacac ttatttttta aacaatccca gaatactgta tgtagacttt 480
 tgttgtgctc aaataaagt ttacttatct tacaagctc aaatactgga ttgtaacctat 540
 gtgatgaagt tatctatggt gtacctaaaca tgcaaatat c 581

<210> 48
 <211> 491
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <222> (1)...(491)
 <223> n = A,T,C or G

<400> 48
 ccgggcccc cctcgagggy ttcaatggtc agatggaaca gttgaaaggc gcggtcgaaa 60
 ccctgcgcac cagcatcgcg caatctggca ttctggaatt cgtcacaaac atcgtcacog 120
 ccttgggcaa ctttgtcgat aagctcgccg aggtcagccc ggaactctg aagtgggta 180
 cgatcatcgg tggggtggcg gcggtgctag gtccggtggc gatccgcac cggccgctg 240
 tctctgcgct gggcgctttt ctccctgtca tctgtgctgt tgcgagcgcc atcggcgctg 300
 tctgttcggt catcacggcc ggtgccatcc cagccctggc cgggcttgtt gttgccctat 360
 cgctctgct cgtgccgctg gcggcggtgg ctgctgcagt cggcgccgct tatctggtgt 420
 ggaagaactg ggacatgac gggcccatcc tcgccaagct ttataacgga gtgaagacgt 480
 ggctggtcga t 491

<210> 49
 <211> 1929
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <222> (1)...(1929)
 <223> n = A,T,C or G

<400> 49
 ttaggctagt agaggctggt gttaatggcg cgagggcgcg tgtcagggtg gagtgcgccg 60
 ccggttcgcy cttggcgacg acaaatgctc gcgcactgct cgtgtggagt acgcgtgtcc 120
 cgaagtgcgc aaggtcttca gctgcccgcc caacctggcc tcgcaccgac gctgcacaaa 180
 accgcggccc gcgcgcgcgc ccgcgcgcgc gccggagcca gaagcagcag ccaggctgag 240
 gcgcgggagc caccgcggcg cggcagcgac cgggacacgc cgagcccgcg cggcgtgtcc 300
 gagtgcggct ccgaggacgg gctctacgag tgccatcaat gcgccaaaga gttccggcgc 360
 caggccctacc tacgcaagca cctgctggcg caccaccagg cctgcaggc caaggcgccg 420
 ccgctagcgc ccccgcccca ggacctactg gcttctgacc ccgggcccga ccgagaaggcg 480

```

ccccaggagg cggccggcga cggcgagggg gccggcgtgc ttgggcctga gtgcgtccga 540
cgagtgcac cctgtgccca gtgtgcggag agtcgttcgc cagcaaggsc gctcaggagc 600
rccrccctgc cctgtctgca cggccgscag gtgttcccc gcaagtactg sctccttgga 660
ccttctacag ctgcgccggc cttacgcggc acatcaacaa gtgccaccca tcgcaaaaaca 720
gacaggtgat cctcctgcag gtgcccgctgc gcccgccctg cttagagcgc cctcccaacc 780
cggcccccga actgtgcctt cgcttgagga ccacaaaaga gagtgcgcc ctcagccccc 840
gaaccggagt ccgcgctggg ggagcctcgc ccccgcccc accgggtgaa agtgctgtct 900
ccgctctctc cggtgtggcg tgacggtaac cctactctct ccttttact ccttttgaa 960
ccccactttt tacgttgtgt cctccgcctt ccccatgcg gcaacaggag tcagctctct 1020
tctgtacaag ggagaaaagc tgtacgcgtt tgtctcgtgg ttggaagcct cccctggcg 1080
gggagaagct ttttttctg ctagtattcg ctgtgttcac ggtctagaaa tgcgctctgg 1140
tctcgcctcg cctaccaatc tctgtctctc atgtatgtag cgtacgggtt gttttgggtg 1200
aatccttgagg aataaatgco tttatatttc acaggctgta aatggaactt cccacacgat 1260
tagctttatt atggcttgtg aactgctgga gtctggcctt accttttctg atgtgaacaa 1320
atcaaaattgc ttaaaaaaga gttttcttta gtatagccac aaatgccttg aactgttgtc 1380
tgggattgtt ttgtgggggg agggaaagga gtgttccgaa gatgctgtag taactgcctc 1440
agtgcttcac gtaagacttt ttggtttgat catcctttgtt gaggtaggac ttcagtgctc 1500
ctctaaatgt atatgttgat ttatgatgaa ttgttattta tcttttattt atttatata 1560
attatgaaga ttatgatatt attgattgc agattttttt ggcgcgctgc cccctcccca 1620
ccctgcactt cttgacattc cactgtgcgt tttagaagag agcctttttc taaaggyatc 1680
tgctttaaag ttttaacttt atacctatct gagtgaatta cagacaacct atcatttatt 1740
ctgcttcgag ggtccccagg gccctgtgac tcttactttt tctactttt atagaaagct tcaatgaaac 1800
tctttctcac atacattatt tcttaattgt ttactatttc aataaaatat ttgcaagctc 1860
tgtttcaact gtataactat ttactatttc aataaaatat aaaaaaaaac 1920
.aaaaaaaag

```

<210> 50
 <211> 6183
 <212> DNA
 <213> Homo sapiens

```

<400> 50
ctttttgtag ggagaagggc aggatgttt taactgaatg tgacctcagg ggaatactag 60
agaaaaaat aaaaattctg aatggggcag cgtggagaaa tctaagaga aatagcataa 120
gagcattttt gaacacatcc aggaaaagat aactttcgac acacctgtag acgttgcaca 180
ggttaaggag tgatggaaac totccagttc agatccagta cgttttaggg aaggaaacta 240
agttgtctgac ttaagtgtgaa gaagcatcta tttaatgtct ggtcaaatcc tacaagaana 300
acagaaatct atgattaaaa agctgagcac ttgtatata tgcaaaaggt agagaaggca 360
ggacggtaga aattttctgc aagaaagaa gaatttcagg atttatcac aaataagaca 420
aagtcattta tttagtcccc ctgacacagc agggcaaacg gagtgtgac acaaagttao 480
tggagaaaaa gagagcaatt ccaggacttc cctttcagcc taaaagaagg taccagatat 540
gtgcactggg gcgagtgtga agagactcgc ttattgcccc tgatgtaagc tccagttaag 600
aaagacgtga agtacaagta ctaggaaatc actttataca tctgtttata ggaatgacct 660
caggactttt tgttcatggt atagatggat gcagagcgct agataaaaa gctgcgtact 720
cgctotaaag gaaccagggt gccaatggat tcaactatcc aggagctcag tttgtcctat 780
gattgtccca tggcaagaaa gagaacagct gaagatcagg ctttgggggt tccagtaaac 840
aaaaggaaat cctgtctaag gaagccocga cactacagcc caaaagcaga ctgccaaaga 900
gaccgcagtg acaggacaga ggacgatggc ccttggaaa cacatggtca ctctaccgca 960
gaggaatca tgatgaatacc tatggatgaa agtcttcttt caactgcaca agaaaactcc 1020
agttaggaag aagacagata cctctgttat caagagctca tggcgaagtc ttaatgca 1080
ttgggggaat ttgaaaaaaa tgtatctggt cagactgtaa gtgaaaaatt aatgcagct 1140
ggcatccagt cttaaaaaagc agagagcgat agtgctttct tctgctcttc tgatgacatt 1200
gatgatggaa gagacaagat tgatgatctc cagccacctt tctgctcttc tgatgacatt 1260
gaaagttaac cttaaaagtgc agaaaatggc tggcagagtg gctccaaact ctcaagaaga 1320
accaaacac cttagagccc aaagtattgt ttaacagagc ataaaaaga cctattgaaa 1380
gttctcgaaa taaaaactga aggtgacaaa ttatccctt gtggaacag gttgtattct 1440
gaaacagaaa ggaagagacc gcagaatgct ctgcagaaac ccttggatgg caatgccag 1500

```


ccctcattcc	ctgcagcttga	ggaggaagat	agcgagagcc	tggcagtaat	gaocggaagag	1560
ggtagtgacc	tggaaaaagcg	caagggggaat	ttaagtttgc	tggagcagcc	aattgctctg	1620
cagcgtgagc	gaggttgtgt	ttccataac	acctacaaag	agctggatag	gttctctgctg	1680
gagcaccctag	caggggaaaag	gaggcaaaac	aaagttatcg	acatgggttg	aagacaatac	1740
tttaacaata	aacattcacc	aaggcctgaa	aagagggaga	ccaagtgcgc	gatccctgga	1800
tgtgatggca	ggggacacgt	gacagggctc	taccgccacc	acgcagcctc	ttcggggtgc	1860
ccccacaaa	tgggggttcc	cctggaaaatt	cttgccatgc	atgaaaaatgt	gtcacaagtgt	1920
ccacgcgcgg	gtgacacagg	aaggggctcat	gtgaacagca	acgcacaacac	ccacagcagt	1980
ctttctggtt	gtccaaattgc	tgcagctgaa	aaattggcaa	tgtccacagga	taaaaatcatg	2040
cttgattctc	cccaaatcgg	gcagtgctct	gaccaggccc	acaggacaag	tttgggtgaag	2100
caaatggaat	tcaatttccc	gtcacaaagg	atcacctctc	ccagagccac	agtgctcaaaa	2160
gaacaagaga	agtttggaag	agtaccattt	gattatgcca	gttttgatgc	ccaagttttc	2220
ggtaaacgcc	ctctctatac	aacagtgcaa	ggacgaaaaa	caccaccatt	tcttgaatca	2280
aagcattttc	caaatccagt	gaaatttcc	aatcgatgc	ctagtgcagg	gcgccacacc	2340
cagagccctc	gcggtgccag	ctcttatagc	tacggctcaat	gtagtgaaga	cacccacata	2400
gcagcagctg	gtgcctatct	gaacctttcc	accgctgca	gggaagccac	agacatctct	2460
tccaacagct	cacagagctc	gcagtccaag	ggagccgaaa	tagaagtggga	tgaaaaatggc	2520
acattggact	gaagcatgaa	aaaaaatcga	atcctggaca	agctcgcacc	cctaagcttc	2580
ttcaacactc	ctatttccac	tctctctctc	tccocattca	aaacaagcag	ctattctggct	2640
aatgcagcat	ctctatcagg	tctttgtgac	caagaggcgt	gggacactcc	tatcaactat	2700
agcaaaactc	acgggaagac	agaggagag	aaagagaag	accagtgag	ctcttagtaa	2760
aathtagagg	aaaaaaagt	tcttgagag	gcctctatac	caagccctaa	accgaagctt	2820
catgcaagag	atctcaaaaa	gaaactaatc	acctgtccaa	caccagatg	tgatgggaagt	2880
ggccacgtga	caggaaacta	tgcctctcat	cgcagtggtt	cttagctgtc	tttagcagat	2940
aagactctaa	aatccctcat	ggctgccaac	tctcaggagc	ttaagtgtcc	aaccocagcg	3000
tgcgactgct	cggggcagct	gactggaaac	tatgcttccc	acagaagctt	gtccggatgc	3060
ccctgtccaa	ggaaaggtgg	tgtcaaaatg	accctaccac	aggaagaaaa	agaagacctt	3120
gaactgaact	ctctgtgtgat	aggggtgtgat	ggccaaggct	acatatcagc	taaatcacaca	3180
tcacacgcga	cagctctctg	ctgtctctct	gctgccaaag	gacagaagga	gaatcctctc	3240
aatggaacct	ccctctctct	gaaactgaac	aaacaagagc	taccacattg	tcccttgcca	3300
ggctgcaatg	ggctggggcca	tgtaaataat	gtttttgtca	ccccccgaag	ottatctgga	3360
tgtctctctc	atgcacaagt	tatcaaaaa	ggcaaggttt	ctgaagaact	catgaccact	3420
aagctcaaa	caactggggg	aatagagagt	gatgaagaaa	ttaggcattt	ggatgaagaa	3480
ataaagagct	tgaatgaatc	caaccttaaa	attgaagcag	atatgatgaa	acttcagacc	3540
cagatcacat	ctatggagag	caacttaag	acgatagagg	aggagaacaa	actcatagaa	3600
cagaacaagt	aaagtctgct	gaaagagctg	gcaggtctaa	gccaaagctc	catttcaagc	3660
cttgcagaca	tccagcttcc	acagatggga	cctatcagtg	agcagaattt	tgaagcatat	3720
gtaaatatgc	tcacagatat	gtacagcaat	ctggaacggg	actattcccc	ggaaatgcaa	3780
gctctactgg	aaagtactaa	acaggcagtg	aaaggtatcc	attgtatgga	tcacagcgct	3840
gcggggcaac	agaagttaac	aacagcagta	aactccagat	ggatctgtta	gaggttcaatg	3900
tactgctaag	gcgtggaggt	tgccgtactg	catttacaat	ttgcaacatt	gcactaaatt	3960
tattttcccc	agctgatala	aaaaaggaag	aaaaactatg	atgactctct	tgttatataa	4020
gcaatgcagt	caattattag	atcttattta	ttttcattatg	ttttttcttt	atttctctac	4080
tgtactcttc	tttttgaaag	tatatgtaaa	ataaatgtga	catttttata	atttttttat	4140
tactaatctc	agagtttttt	atcttttaac	tgcattttga	agttctgcgt	attttttaca	4200
gtgtgtttat	taattttatt	tccaatagga	tttaaataga	aatgtctatc	tcaagctcatc	4260
tttctctgct	ggttttaaatg	aggaacacag	aaaggggtga	ggaaatccct	gtctaaggagc	4320
tgcaactatg	ttgagtttga	tttttattgc	acacttcttc	ccccactttt	cactgatttt	4380
tgtattttata	aatgaatttg	cggtaaagtg	agctgcacgg	aaggaaataag	aagacaaatg	4440
gcggccacta	tgggggaatc	cgcaactaca	aaagcacagg	atgctggaaa	acagcctgct	4500
cagaatttgt	tagcaataat	taaatatagc	aatcagcaaa	gtatctgact	tggttgagct	4560
gttttctgta	atatgaatta	tttatttgaa	atgtttttaa	gaaacataag	cccttttatgt	4620
gatgcagatt	tgtctgtttg	tttttcaagt	catatcagat	cgttgccaac	tcgatcccc	4680
agatgaaaaa	taagacttgg	tgtgaccagc	cagggtttcc	tgccatgatg	tggtacaata	4740
tcaacgtgac	aatattgggt	tagattttga	cttagcaaat	acaaacacat	ccaaatgaaa	4800
aattttgtag	ataccatatc	ccctgaaata	gcattttatc	tactgggttg	actctgaaag	4860
aatgaaaaat	atgtaaaaac	atgaaaaaat	gctactccaa	tctgaatgat	cactgcaaac	4920
actggcacct	tgggtctcac	ccaccatag	aaacaagaca	acattcaatt	tgatagaaat	4980

```

cttgccacaa aacttcaaat gctacaaaa atacacacac actcacacac acaggcatat 5040
tcacacacag acacacacac acacacacac acagactcat ccacacttca aattgagccc 5100
acaattctga attctctgaac ggatcagagt ttcatagttt ctatagttaa ggcaatgtct 5160
atttcaggga ttgtaaagta gttaagcatt gtttcaaaag tttttttata tttatttttt 5220
ttaaaggaaa ggtatagaca accagctaaa ctgccttttt ggtgtgcaca cacattcat 5280
gtgcagacgt gccctagtgt gctcatgtca atgaacttca tttgggctta tttttctgtg 5340
ctataaacaa aagtgtttat tttttattaa cctcatggat atttagatgg aaagtatggg 5400
cattcacaga ttctagtat tccactgtta tctactgtac ctgcacaaat gaaaacaaat 5460
actcaacagt aattccact ccatgaaact ttggtcattg ttatgcatta agtggggctt 5520
atcttttggt ttgaagtcat ttgaactctt gaaccttagt ttagtgaaga tgaactgtct 5580
gtctttaggt agaaacgggt tttattttaa aatcagtttt aaaaatgag ctaccatagt 5640
tgctgtctat tataaatggg acaccaaca aaattttcta ttacagttgt gtacttgcac 5700
acattttgct atacagtact tcataagatgc atacaaatga gctcacttat tacaagaca 5760
aacgtttaat ttgtcaata ttttaacaag ttgtttatat attttattta atttaaaaga 5820
aatctcttac caacctacat attttattact ataatttgc atgaacttcag gtttaatttat 5880
ttgtgtttgc atagtgttgag caggatgttt tgtgaagat gtttgtattt atttgctac 5940
tttgtacttg atgtgttttg taatgtgcac tgaatttgtt ttcttttcaa ctatgttaat 6000
gatcaatact gtaaatgtgg tctttgttaa acaaaaagcg aatgatgtat gcattttttt 6060
taattttgag tagttttgtt gtatactgtt tctccaaaca ctaatatatt ctacatcaa 6120
agcaacaaaa ttgtgttcag tgctgtacat ttggtgtatg taggaaata aaaattgata 6180
acg

```

```

<210> 51
<211> 1704
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<222> (1)... (1704)
<223> n = A,T,C or G

```

```

<400> 51
tcagaaaaaa taaaagatat ataggagcca caagtgtctt ggggaccata taaaacacog 60
tggtttgggtg cctattagaa tataacgttg ggctgtctcg ctgttacgag tgtacaatgc 120
ctcttcgcgcg gtttgttcaa tataccgcgc cgcgcgctat ctttcgcaag gttgtttaca 180
gccctacacc gcaggtttacc cagaggtaat cgggagagct taaaataacc gttactcctg 240
aaaaaagta tgtaaagagc gaattttctc agtcatagtt gaataatcaa tgaagtgtgc 300
ttgtctctta atgtccttac ccattctctg ataattcttt attagaatga atgttgagag 360
cctgggggagt cttaggatat tcttgagaaa taaatttgaa gtgccatttt ttgctaaaacg 420
taggttagaaa atggcgctttt agattttcaa aagtaaatgg ctaaaaatta agcattattac 480
ccttcagaaa gtttataaag tttagccatc atttttttaa cacagaaatc tgtttattaa 540
accaaacaaa acagagaaaa ttataccagc cctcaatttt tgaattttca tttaaaaaag 600
caaacctctaa atccacatct taaaagatgt ttgtgcagct atgtatttcc aaaaacttca 660
tattttcaata agatttttca catttatatt accaacagta tcacaaaagt tttttttttg 720
ttttttgttt acataattgt aaggaacagt aattctagaa acactagaag aaaaaagcat 780
agcaatgtcc agcgttacaa gaaaagtgc acatctactg gtccacaaat cagtattac 840
ttgaaaaact atatgtaaac agtagataag aaatatcact gatgcctcaa actcattgtc 900
aaaaactgaa tgaacataat ttacatgaa ttaaggcaaa ttacaggaatg cacaagaagt 960
ttgtaatcca accaaaacta acaacagaa aaaagtgtga taagaagcat gaactaaagt 1020
acttctccct aaatttttaa aaataggct tgtctcagtg cacaaagaaa acatcatca 1080
tgtgtatccc acactataaa ataagaaaga aggtgaaagt atgggggata ggagggcaca 1140
gttcaattga agttgcagct gcatccgctg agagttcctt acattatttt tagctagaac 1200
tgaaatttat acaaatcata tcaggagatg taatggctct ttgggaacct atttctgaaa 1260
gaaatgaaaa gaaaactaca cacaagagtg caaattttca gattgtcact tgaacctct 1320
taacattcag tcactctacat ccaggtgctg ctgagggatg gcctggagac agcagcgca 1380
atcaggaacg agcagctcta agaaaccaag gtgtgatttt ttttcaacaa catgtcttgt 1440

```

```

cattattaaaa aaaaaaatc tgggatgaaa actgctatga taaagtgca gtgttgagt 1500
gggttttttga gatcagcatg agagcagaaa tgcaggcttc tcttggaagt agttcctgat 1560
gtgacgattg aaagaacgta ggcaagggtt ttccagcat caagtgttat tttgtagaa 1620
agaatttgga aagaggagaa ggcaaggga tgtggaaaag gtaactacag tagtttctca 1680
aaacagtttt ctttttaggc ctat 1704

```

```

<210> 52
<211> 1886
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> misc_feature
<222> (1)...(1886)
<223> n = A,T,C or G

```

```

<400> 52
taaattccgt tgttactcaa gatgactgct tcaagggtaa aagagtgcac cgcttttagaa 60
gaagtttggc agtattttaaa tctgttgatg cctctcagct atctagtttc atgggaagtt 120
gctgtttttg aatatttaagc taaaagtttt ccaattttac agaaattctg aatttttgta 180
aatcacactg aacttttctg tataacttgt attatttagac tctctagttt tatcttaaca 240
ctgaaactgt tcttcttag agtttttttt agaacctggt tctgtgttta atataagtt 300
taaaagtaaa aataatcgag actgaaagaa tgttaagatt tatctgcaag gatttttttaa 360
aaatggaac ttgcatttta agtggttttaa agcaaatct gactttccaa aaagttttta 420
aaaactgatt tgaaagctaa caattttgat agtctgaaac caagcatttc acttctccaa 480
gaagtacctg tgaaacgtac aatatttcag tattgagctt tgcatttatg atttatctag 540
aaatttacct caaaagcaga atttttaaaa ctgcattttt aatcaagtga actcaatgta 600
tagttagctt tattgaagtc ttatccaaac ccagtaaaac agattctaa gaaacagttc 660
aatcagtgag tcataatggt tattcaaaag attttatctt ttatctagaa nccactatc 720
tatgtccaat ttgatnngga tagtagttag gataactaaa attctgggcc taatttttta 780
aagaatccaa gacaaactaa actttactgg gtatataaac ttctcaatga ggtaccact 840
ttttttataa aaaaaattgt tccctgaaat gctaaaacta atggctgtat gtgaaatttg 900
caaaactact gtatataaga acgctgcagc ttttttatgt caactcaagg ttaactggag 960
tatctgaaag gaattgtttt tataaaaaaca ttgaagtatt agttacttgc tataaataga 1020
ttttttttt tgttttttag cctgttatat ttctctctgt aaaaataaat atgtccagaa 1080
gaggaatggt gtttctagat taggtagtgt cctcatttta ttgtgtgacc caacagctag 1140
agcaacagag cctctttgct atactcacag tctgttttcc ccagcctctt ttactagtct 1200
ttcaggaggt ttgctcttag aactggtgat gtaagaatg gaagttagct ttagcagagt 1260
tccaggccca agcctgtgaa tggtagcaat gggatataat accctcttaa gggaaacatt 1320
tgtagcagta tcatttgatc tgcactggac atgtgtttta atgtggttcc ttggcctctt 1380
ttcaatggct tcttccctaa aacgtggaga ctctaagtta atgtcgttac tatgggcat 1440
attactaatg ccaactgggg tctatgattt ctcaaaattt tcattcgtaa tccgaagat 1500
acagtcttta aactttagaa ttcccaagaa ggctttatta caactcagaa attgaaagaa 1560
ccatgacttt gtccattaaa aaattatcca tagttttttt agtgcattta acattccag 1620
atacatcatt ctgtgtattaa atctccagat ctctgtaaat gatactacaa tctcaaaag 1680
ttaatcttaa ttattccgat atgaccttaa ggaaaagtaa aggaataaat tttgtcttt 1740
gttgaagtat ttaatagagt aaggtaaaga agatattaa tctcttcaa aatggaaat 1800
taattctaaa ctgagaaaaa tgttctact acctattgct gatactgtct ttgcataat 1860
gaataaaaat aaactttttt tcttca 1886

```

```

<210> 53
<211> 877
<212> DNA
<213> Homo sapiens

```

```

<220>

```

<221> misc_feature
 <222> (1)...(877)
 <223> n = A,T,C or G

<400> 53
 tttyggcacga ggaattttct aacawtkwt yttttaatag ttagactcat actttatatt 60
 gacaaatttta agatagaaaa atatcataat gtgaatatag cagttgtctct ttttgaaca 120
 tggtttgga tggtcagtgga aacttgaaa gacttgcctt acaggtggct cctottctgg 180
 ctgggtttca gttaattctg aatttatctc cagccattgc atttgcttga aagaattattg 240
 gacacagtaa aaaaaagaac aggtttggca tcaataata aattattata agcaatgaac 300
 caaaacaact tttaaaataa ttactgaaa caaacttcag acttcatgat taaagctaa 360
 aactcatatt ttcaaaatag ctttaacagt ttctatcaat atataatata atartaggac 420
 acttattttt aaaaaacaag tgagtagaat cagagtaaat atgatatctt agatgactat 480
 aaacagtaaa catcaattca atatatattt atatcattc agcaatatac tctktgcca 540
 gctggcgata aaaactgtag ttctatcatc aaaaaatgca tccctgaatg tcatctttga 600
 acttactaag tgctgtcatc atttctacac tccatcttgg gaggggggtg cttagggact 660
 cttggtacat gcagatattt agttatgggt ataagtacaa aaagttaagt tggcaggagt 720
 ctgaagcaga aacgttgcct tactttgtta agtagcttca cattcttttg tctctgtgat 780
 gctcaggtg aagtcacact aaataattca cacaggtgct aattttgtg cctctgtgca 840
 gtacctttca gctttttct ttcttctcct tccccac 877

<210> 54
 <211> 1364
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <222> (1)...(1364)
 <223> n = A,T,C or G

<400> 54
 tttttttttt tttttttgat tanattaagg ggctgccagc ccggagagaat acttaagata 60
 tgggtgagaa atccccagac ttttatcaaa aagatttoca ctttcaaatc aatgtcagta 120
 gacattgata aaagtatagc agcatcctct actgaggtga tttcatttat tccctgcagc 180
 ccaactgata atctctcact tctcccaaat agtatgtgga ctccagcata agcagaaaaac 240
 tattgtcatt caactgaaga agaggaagat aaaagattgt cttgtttcca tcaactgtatt 300
 acttgggtaa catgattaca taattcttat cctaagagaa agctttcata tttaaaaaaa 360
 agtcttttca gataaaatct gotttgtctc tgaataatat gaaatacaaa ctttcaacttt 420
 attttattgt aaattatcaa gagattattg tcttataata tatattgagt tagcttcaag 480
 ctctcctaaa tatgaagaga ttgttgtcta aagtcacata ttgacattga gctcagttgc 540
 ctgttctcat acgtatgtgc tgctacotgt acagcagaca tgcctgccta gtgacattta 600
 taatgacaga agcagggtga tggctttgtg ttgacatga tcaagttaga tcatagactt 660
 tccctgaact gtgatatta gctctgaatt ggggggaaag argactttga cacattttag 720
 ttattttaat aacagagatt tactcttttg aaaaaataag gtatctaatg tctccctaat 780
 aagtcctctt tctctccaac taatgacct acacggactt ttattttctt gatcaaaag 840
 gtgttttata aggacttctg gataactata cttttactct atttttaaag atcaaaaagt 900
 aatttttaaa gtgaacaggt tcccatacca tgaatgctgg cctcaccctc tctatcatcc 960
 acattttgaa atgcgaagaa agctcccttg taagccatc ttctctcccc actcccatcc 1020
 taggataact gccacgtgct cattaggcat ttctattcca gatagcccaa atttaggtta 1080
 ttatgcttaa ttgacacat taactaaatg occagtttta aaatatatcc atcaattcac 1140
 gctgaaatgt gcttctttgt gctatcaaat ggaatagaat acactatttt ttaataacat 1200
 ccagaaatc tggtgtgata cttttgttgt gctcaataaa atgtttactt atcttcaaaa 1260
 gctcaaatc tggattgtaa ccatgtgatg aagtattata tgtttacctt aacattgcaa 1320
 attaatcaat aaattctctgt tgcataaaaa aaaaaaaa aaaa 1364

Leu Phe Ala Pro Ala Glu Leu Lys Met Gly Thr Ala Phe Ser Ala Gly
 165 170 175
 Ala Glu Ala Ala Arg Gly Pro Gly Pro Gly Pro Pro Leu Pro Pro Ala
 180 185 190
 Ala Ala Leu Arg Pro Pro Gly Lys Arg Pro Pro Pro Pro Thr Ala Ala
 195 200 205
 Glu Pro Pro Ala Lys Ala Val Lys Ala Pro Gly Ala Lys Lys Pro Lys
 210 215 220
 Ala Ile Arg Lys Leu His Phe Glu Asp Glu Val Thr Thr Ser Pro Val
 225 230 235 240
 Leu Gly Leu Lys Ile Lys Glu Gly Pro Val Glu Ala Pro Arg Gly Arg
 245 250 255
 Ala Gly Gly Ala Ala Arg Pro Leu Gly Glu Phe Ile Cys Gln Leu Cys
 260 265 270
 Lys Glu Glu Tyr Ala Asp Pro Phe Ala Leu Ala Gln His Lys Cys Ser
 275 280 285
 Arg Ile Val Arg Val Glu Tyr Arg Cys Pro Glu Cys Ala Lys Val Phe
 290 295 300
 Ser Cys Pro Ala Asn Leu Ala Ser His Arg Arg Trp His Lys Pro Arg
 305 310 315 320
 Pro Ala Pro Ala Ala Ala Arg Ala Pro Glu Pro Glu Ala Ala Ala Arg
 325 330 335
 Ala Glu Ala Arg Glu Ala Pro Gly Gly Gly Ser Asp Arg Asp Thr Pro
 340 345 350
 Ser Pro Gly Gly Val Ser Glu Ser Gly Ser Glu Asp Gly Leu Tyr Glu
 355 360 365
 Cys His His Cys Ala Lys Lys Phe Arg Arg Gln Ala Tyr Leu Arg Lys
 370 375 380
 His Leu Leu Ala His His Gln Ala Leu Gln Ala Lys Gly Ala Pro Leu
 385 390 395 400
 Ala Pro Pro Ala Glu Asp Leu Leu Ala Leu Tyr Pro Gly Pro Asp Glu
 405 410 415
 Lys Ala Pro Gln Glu Ala Ala Gly Asp Gly Glu Gly Ala Gly Val Leu
 420 425 430
 Gly Leu Ser Ala Ser Ala Glu Cys His Leu Cys Pro Val Cys Gly Glu
 435 440 445
 Ser Phe Ala Ser Lys Gly Ala Gln Glu Arg His Leu Arg Leu Leu His
 450 455 460
 Ala Ala Gln Val Phe Pro Cys Lys Tyr Cys Pro Ala Thr Phe Tyr Ser

465 470 475 480
 Ser Pro Gly Leu Thr Arg His Ile Asn Lys Cys His Pro Ser Glu Asn
 485 490 495
 Arg Gln Val Ile Leu Leu Gln Val Pro Val Arg Pro Ala Cys
 500 505 510

 <210> 57
 <211> 1047
 <212> PRT
 <213> Homo sapiens

 <400> 57
 Met Asp Ala Glu Ala Glu Asp Lys Thr Leu Arg Thr Arg Ser Lys Gly
 5 10 15
 Thr Glu Val Pro Met Asp Ser Leu Ile Gln Glu Leu Ser Val Ala Tyr
 20 25 30
 Asp Cys Ser Met Ala Lys Lys Arg Thr Ala Glu Asp Gln Ala Leu Gly
 35 40 45
 Val Pro Val Asn Lys Arg Lys Ser Leu Leu Met Lys Pro Arg His Tyr
 50 55 60
 Ser Pro Lys Ala Asp Cys Gln Glu Asp Arg Ser Asp Arg Thr Glu Asp
 65 70 75 80
 Asp Gly Pro Leu Glu Thr His Gly His Ser Thr Ala Glu Glu Ile Met
 85 90 95
 Ile Lys Pro Met Asp Glu Ser Leu Leu Ser Thr Ala Gln Glu Asn Ser
 100 105 110
 Ser Arg Lys Glu Asp Arg Tyr Ser Cys Tyr Gln Glu Leu Met Val Lys
 115 120 125
 Ser Leu Met His Leu Gly Lys Phe Glu Lys Asn Val Ser Val Gln Thr
 130 135 140
 Val Ser Glu Asn Leu Asn Asp Ser Gly Ile Gln Ser Leu Lys Ala Glu
 145 150 155 160
 Ser Asp Glu Ala Asp Glu Cys Phe Leu Ile His Ser Asp Asp Gly Arg
 165 170 175
 Asp Lys Ile Asp Asp Ser Gln Pro Pro Phe Cys Ser Ser Asp Asp Asn
 180 185 190
 Glu Ser Asn Ser Glu Ser Ala Glu Asn Gly Trp Asp Ser Gly Ser Asn
 195 200 205
 Phe Ser Glu Glu Thr Lys Pro Pro Arg Val Pro Lys Tyr Val Leu Thr
 210 215 220
 Asp His Lys Lys Asp Leu Leu Glu Val Pro Glu Ile Lys Thr Glu Gly

225		230		235		240
Asp Lys Phe Ile Pro Cys Glu Asn Arg Cys Asp Ser Glu Thr Glu Arg						
		245		250		255
Lys Asp Pro Gln Asn Ala Leu Ala Glu Pro Leu Asp Gly Asn Ala Gln						
		260		265		270
Pro Ser Phe Pro Asp Val Glu Glu Asp Ser Glu Ser Leu Ala Val						
		275		280		285
Met Thr Glu Glu Gly Ser Asp Leu Glu Lys Ala Lys Gly Asn Leu Ser						
		290		295		300
Leu Leu Glu Gln Ala Ile Ala Leu Gln Ala Glu Arg Gly Cys Val Phe						
		305		310		315
His Asn Thr Tyr Lys Glu Leu Asp Arg Phe Leu Leu Glu His Leu Ala						
		325		330		335
Gly Glu Arg Arg Gln Thr Lys Val Ile Asp Met Gly Gly Arg Gln Ile						
		340		345		350
Phe Asn Asn Lys His Ser Pro Arg Pro Glu Lys Arg Glu Thr Lys Cys						
		355		360		365
Pro Ile Pro Gly Cys Asp Gly Thr Gly His Val Thr Gly Leu Tyr Pro						
		370		375		380
His His Arg Ser Leu Ser Gly Cys Pro His Lys Val Arg Val Pro Leu						
		385		390		395
Glu Ile Leu Ala Met His Glu Asn Val Leu Lys Cys Pro Thr Pro Gly						
		405		410		415
Cys Thr Gly Arg Gly His Val Asn Ser Asn Arg Asn Thr His Arg Ser						
		420		425		430
Leu Ser Gly Cys Pro Ile Ala Ala Ala Glu Lys Leu Ala Met Ser Gln						
		435		440		445
Asp Lys Asn Gln Leu Asp Ser Pro Gln Thr Gly Gln Cys Pro Asp Gln						
		450		455		460
Ala His Arg Thr Ser Leu Val Lys Gln Ile Glu Phe Asn Phe Pro Ser						
		465		470		475
Gln Ala Ile Thr Ser Pro Arg Ala Thr Val Ser Lys Glu Gln Glu Lys						
		485		490		495
Phe Gly Lys Val Pro Phe Asp Tyr Ala Ser Phe Asp Ala Gln Val Phe						
		500		505		510
Gly Lys Arg Pro Leu Ile Gln Thr Val Gln Gly Arg Lys Thr Pro Pro						
		515		520		525
Phe Pro Glu Ser Lys His Phe Pro Asn Pro Val Lys Phe Pro Asn Arg						
		530		535		540

Leu Pro Ser Ala Gly Ala His Thr Gln Ser Pro Gly Arg Ala Ser Ser
 545 550 555 560
 Tyr Ser Tyr Gly Gln Cys Ser Glu Asp Thr His Ile Ala Ala Ala Ala
 565 570 575
 Ala Ile Leu Asn Leu Ser Thr Arg Cys Arg Glu Ala Thr Asp Ile Leu
 580 585 590
 Ser Asn Lys Pro Gln Ser Leu His Ala Lys Gly Ala Glu Ile Glu Val
 595 600 605
 Asp Glu Asn Gly Thr Leu Asp Leu Ser Met Lys Lys Asn Arg Ile Leu
 610 615 620
 Asp Lys Ser Ala Pro Leu Thr Ser Ser Asn Thr Ser Ile Pro Thr Pro
 625 630 635 640
 Ser Ser Ser Pro Phe Lys Thr Ser Ser Ile Leu Val Asn Ala Ala Phe
 645 650 655
 Tyr Gln Ala Leu Cys Asp Gln Glu Gly Trp Asp Thr Pro Ile Asn Tyr
 660 665 670
 Ser Lys Thr His Gly Lys Thr Glu Glu Glu Lys Glu Lys Asp Pro Val
 675 680 685
 Ser Ser Leu Glu Asn Leu Glu Glu Lys Lys Phe Pro Gly Glu Ala Ser
 690 695 700
 Ile Pro Ser Pro Lys Pro Lys Leu His Ala Arg Asp Leu Lys Lys Glu
 705 710 715 720
 Leu Ile Thr Cys Pro Thr Pro Gly Cys Asp Gly Ser Gly His Val Thr
 725 730 735
 Gly Asn Tyr Ala Ser His Arg Ser Val Ser Gly Cys Pro Leu Ala Asp
 740 745 750
 Lys Thr Leu Lys Ser Leu Met Ala Ala Asn Ser Gln Glu Leu Lys Cys
 755 760 765
 Pro Thr Pro Gly Cys Asp Gly Ser Gly His Val Thr Gly Asn Tyr Ala
 770 775 780
 Ser His Arg Ser Leu Ser Gly Cys Pro Arg Ala Arg Lys Gly Gly Val
 785 790 795 800
 Lys Met Thr Pro Thr Lys Glu Glu Lys Glu Asp Pro Glu Leu Lys Cys
 805 810 815
 Pro Val Ile Gly Cys Asp Gly Gln Gly His Ile Ser Gly Lys Tyr Thr
 820 825 830
 Ser His Arg Thr Ala Ser Gly Cys Pro Leu Ala Ala Lys Arg Gln Lys
 835 840 845

Glu Asn Pro Leu Asn Gly Ala Ser Leu Ser Trp Lys Leu Asn Lys Gln
 850 855 860
 Glu Leu Pro His Cys Pro Leu Pro Gly Cys Asn Gly Leu Gly His Val
 865 870 875 880
 Asn Asn Val Phe Val Thr His Arg Ser Leu Ser Gly Cys Pro Leu Asn
 885 890 895
 Ala Gln Val Ile Lys Lys Gly Lys Val Ser Glu Glu Leu Met Thr Ile
 900 905 910
 Lys Leu Lys Ala Thr Gly Gly Ile Glu Ser Asp Glu Glu Ile Arg His
 915 920 925
 Leu Asp Glu Glu Ile Lys Glu Leu Asn Glu Ser Asn Leu Lys Ile Glu
 930 935 940
 Ala Asp Met Met Lys Leu Gln Thr Gln Ile Thr Ser Met Glu Ser Asn
 945 950 955 960
 Leu Lys Thr Ile Glu Glu Glu Asn Lys Leu Ile Glu Gln Asn Asn Glu
 965 970 975
 Ser Leu Leu Lys Glu Leu Ala Gly Leu Ser Gln Ala Leu Ile Ser Ser
 980 985 990
 Leu Ala Asp Ile Gln Leu Pro Gln Met Gly Pro Ile Ser Glu Gln Asn
 995 1000 1005
 Phe Glu Ala Tyr Val Asn Thr Leu Thr Asp Met Tyr Ser Asn Leu Glu
 1010 1015 1020
 Arg Asp Tyr Ser Pro Glu Cys Lys Ala Leu Leu Glu Ser Ile Lys Gln
 1025 1030 1035 1040
 Ala Val Lys Gly Ile His Val
 1045

<210> 58

<211> 2165

<212> DNA

<213> Homo sapiens

<400> 58

cgcacacgct ggggtcggcg aggcggcg gatcgggc ctgtgccggg gcccggtgct 60
 gggcatctcg gtggccatcg cgcacggggt ctctcgggc tccctcaaca tcttgctcaa 120
 gtctctcatc agccgctacc agttctcctt cctgacctg gtgcagtgc tgaccagctc 180
 caccgcgccg ctgagccctgg agctgctcgg cgcctcggg ctctcgcgc tgccccctt 240
 cggctctgagc ctggcgcgct ccttcggggg ggtgcgggtg ctctccacgc tgcagtcacg 300
 cctcacgctc tggctccctg cgggcctcag cctgcccatg tacgtggtct tcaagcgctg 360
 cctgccccctg gtcaccatgc tcatcggcgt cctggtgctc aagaacggcg cgcctcggc 420
 aggggtgctg cggcgcggtc tcatcaccac ctgcggcgcc gccctggcag gagccggcga 480
 ctgcacggcg gaccccatcg ggtacgtcac gggagtgctg gcggtgctgg tgcacgctgc 540
 ctacctggtg ctcatccaga aggcacgcgc agacaccgag caccggccgc tcaccgcgca 600
 gtacgtcatc gcgctctctg ccaccccgct cctggctcct tgctccttcg ccagcaccga 660
 ctccatccac gcttggaact tcccgggctg gaaggaccgc gccatggtct gcattcttgt 720

```

ggcctgcctc  ctgatcggtc  ggcgcctgaa  cttcaccacg  ctgcactgca  cctacatcaa  780
ttcgccgctg  accacctctc  tgttccattg  cgccgtggtg  gtgaacaccc  tgggctctat  840
catttactgt  gtggccaagt  tcatggagac  cagaagacaa  agcaactacg  aggcctctga  900
ggccacgctc  cggggagagg  aggcgcagct  aagtggagac  cagctgcctg  tctgatgga  960
ggagctgcct  ggggagggag  gaaatggcgg  gtcagaaggt  ggggaggcag  caggtggccc  1020
cgctcaggag  agcagggca  aggtcagggg  cagcccccga  ggagtcocgc  tgggtgctgg  1080
gagctctgaa  gaaggagaca  ggaggtcgtt  aaaagatgct  taacctcagg  tatggaggtt  1140
ggttaggagg  acaggttata  tgagaaggga  ttatttgata  gaaacagagg  gttaccocag  1200
tccctgagaa  ggaggtgcac  gtacgtacct  atgtgcatac  acttatttta  tatgttagaa  1260
atgacgtgtt  ttaattgag  gctcctccgt  ttattctttt  gaggagtggt  gaagggaaga  1320
aaagaagaa  gctgaaggt  actgacacag  agcaacaaaa  ttgacacctg  tgtgaattat  1380
ttaagtgtac  ttccactgag  gcatacaga  gacaaaagaa  tgtgaagcta  cttacaaaag  1440
taaggcaacg  ttctcgtctc  agactcctgg  cacatttact  ttttgcatt  ataaccataa  1500
ctaaatatct  gcattgacca  agagtcccta  agccaccccc  tccaagatg  gagttagaa  1560
atgatgacag  cacttagtaa  gttcaasgat  gacattcagg  gatgcatttt  ttgatgatag  1620
aactacagtt  ttatcgcca  gctgggcaaa  gattatattg  ctgaaatgat  atataaatat  1680
attgaattga  tgtttactgt  ttatagtcac  ctgaaatctc  atattacttc  tatgttctac  1740
cactgttttt  ttaaaaaata  gtgtctact  attgtattat  atattgatag  aaactgttaa  1800
agctattttt  aaaaattgag  ttcttagctt  taactcatgaa  gctgaaggtt  ggtttctagt  1860
aattattttt  aaagtgtttt  tggttcattg  ctttataata  ttttattatt  aatgccaata  1920
ctgttttttt  ttttactgtt  tccaattatt  ttccaagcaa  atgcaatggc  ggaataataa  1980
ttcagaatta  actgaacccc  agccagaaga  gggaccacct  gtaagcaatc  tcccttcaag  2040
tttccactga  catccgaaac  catgtttcaa  aaagagaac  tgcatatttc  acattatgat  2100
atttttctat  cttaaatttg  tcaaaaataa  gtatgagctc  aactattaaa  aaaaaaaa  2160
aaaaa  2165

```

<210> 59

<211> 1776

<212> DNA

<213> Homo sapiens

<400> 59

```

atgcgcgcagc  tgtgcggggg  cgcgctgctg  ggcactctgc  tggccatcgc  gcacgggggtc  60
ttctcggggtc  cctcaacat  ctgtctcaag  ttctctatca  ggcgctacca  gttctctctc  120
ctgacccctgg  tgcagctcct  gaccagctcc  accgoggcgc  tgagcctgga  gctgctgogg  180
cgctcggggc  tcatcgccgt  gccccccttc  ggtctgagcc  tggcgcgctc  ctgcgggggg  240
gtcgcgggtgc  tctccacgct  gcagtcacgc  ctccagctct  ggtccctgcg  cggtcccaag  300
ctgcccactgc  acgtggtctt  caagcgtgct  ctgcacctgc  tcaccatgct  catcgcgctc  360
ctggtgcttca  agaacggcgc  gccctgcgca  ggggtgctgg  cggcggtgct  catcaccacc  420
tcgcgcgccg  ccttgccagg  agccggcgac  ctgacggcgc  accccatgag  gtaagtcacg  480
ggagctctgg  cgggtgctgt  gcacgctgcc  taacctgtgc  taccacagaa  ggccacagaa  540
gacacccagc  acggcgccgt  caccgcccag  taagtcacgc  ccgtctctgc  caccocgctg  600
ctggtctact  gctcctctgc  cagcaccgac  tccatccagc  cctggacctg  cccgggctgg  660
aaggaccccg  ccatggtctg  catctctgtg  gctcgatcac  tgatcggtgc  cgccatgaac  720
ttcaccacgc  tgcactgcac  ctacatcaat  toggccgtga  ccacctctct  gttcattgoc  780
ggcgtggtgg  tgaacaccc  gggctctatc  atttactgtg  tggccaagt  catggagacc  840
agaaagcaaa  gcaactacga  ggacctggag  gccacgctc  ggggagagga  ggccagacta  900
agtggagacc  agctgcgctt  cgtgatggag  gagctgcocg  gggaggagag  aaatggccgg  960
tcagaaggtg  gggagggcgc  agtgggcccc  gctcaggaga  gcaggcaaga  ggtcaggggc  1020
agccccgag  gactcccgct  ggtggctggg  agctctgaag  aaggagacag  gaggtcgtta  1080
aaagatgctt  acctcgaggt  atggaggttg  gtaggggaa  ccagggtatat  gaagaaggat  1140
tatttgatag  aaaaacagga  gttaccacgt  ccttga  1176

```

<210> 60

<211> 1089

<212> DNA

<213> Homo sapiens

[illegible]

65		70		75		80
Val Ala Val Leu Ser	Thr Leu Gln Ser Ser Leu Thr Leu Trp Ser Leu					
	85		90			95
Arg Gly Leu Ser Leu Pro Met Tyr	Val Val Phe Lys Arg Cys Leu Pro					
	100		105			110
Leu Val Thr Met Leu Ile Gly Val Leu Val Leu Lys Asn Gly Ala Pro						
	115		120			125
Ser Pro Gly Val Leu Ala Ala Val Leu Ile Thr Thr Cys Gly Ala Ala						
	130		135			140
Leu Ala Gly Ala Gly Asp Leu Thr Gly Asp Pro Ile Gly Tyr Val Thr						
	145		150			155
Gly Val Leu Ala Val Leu Val His Ala Ala Tyr Leu Val Leu Ile Gln						
	165		170			175
Lys Ala Ser Ala Asp Thr Glu His Gly Pro Leu Thr Ala Gln Tyr Val						
	180		185			190
Ile Ala Val Ser Ala Thr Pro Leu Leu Val Ile Cys Ser Phe Ala Ser						
	195		200			205
Thr Asp Ser Ile His Ala Trp Thr Phe Pro Gly Trp Lys Asp Pro Ala						
	210		215			220
Met Val Cys Ile Phe Val Ala Cys Ile Leu Ile Gly Cys Ala Met Asn						
	225		230			235
Phe Thr Thr Leu His Cys Thr Tyr Ile Asn Ser Ala Val Thr Thr Ser						
	245		250			255
Leu Phe Ile Ala Gly Val Val Val Asn Thr Leu Gly Ser Ile Ile Tyr						
	260		265			270
Cys Val Ala Lys Phe Met Glu Thr Arg Lys Gln Ser Asn Tyr Glu Asp						
	275		280			285
Leu Glu Ala Gln Pro Arg Gly Glu Glu Ala Gln Leu Ser Gly Asp Gln						
	290		295			300
Leu Pro Phe Val Met Glu Glu Leu Pro Gly Glu Gly Gly Asn Gly Arg						
	305		310			315
Ser Glu Gly Gly Glu Ala Ala Gly Gly Pro Ala Gln Glu Ser Arg Gln						
	325		330			335
Glu Val Arg Gly Ser Pro Arg Gly Val Pro Leu Val Ala Gly Ser Ser						
	340		345			350
Glu Glu Gly Ser Arg Arg Ser Leu Lys Asp Ala Tyr Leu Glu Val Trp						
	355		360			365
Arg Leu Val Arg Gly Thr Arg Tyr Met Lys Lys Asp Tyr Leu Ile Glu						
	370		375			380

Asn Glu Glu Leu Pro Ser Pro
385 390

<210> 63
<211> 442
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<222> 220,391,428
<223> n = A,T,C or G

<400> 63
atagtaagca ctgattgtgtt tattcgtatga aataggggtg ggggtgtagc agccctagtc 60
ccacatttgc tgggctgggtg actgagtttaa cagcaaaagt ggatgcaaaa ggttctctgat 120
tgagagcccc cggattcggg ttctggatttt gctggccact tactctatga cttggggcat 180
gtcaactgtca tggcctcagt ttccccttct gcacagtgtn ttattggata gttccagctc 240
tgacatgcta ggattatgtg atactgtcaa tcaagactag ggttggccta agcacatggt 300
otgaaaacac ctccgggtccta tggacatat ttctccgcct ggggagtggt cagctcgtga 360
gtggcaaggc tgccctccaa agctgtccat nccacgcccc ggggtgctgtg ggtctctctt 420
ccctcgtngc cgaattcttg gg 442

<210> 64
<211> 456
<212> DNA
<213> Homo sapiens

<400> 64
cttcaacccat aaaaacaaag ggctctgatt gctttagggtg ataagtgtt taatatccac 60
aaacgtcccc actccaaaaa gtaactatat tctggatttc aacttttctt ctaattgtga 120
atctcttctgt tttttcttct taaggaggaa agttaaagga cactacaggt catcaaaaac 180
aagtgtggcca aggactcatt acttgtctta tatttttact gccactaaac tgccctgtatt 240
tctgtatgtc ctctctatcca aacagacggt cactgcccact tgtaaagtga aggatgtaaa 300
cgaggatata taactgttttc agtgaacaga ttttgtgaag tgccctctgt tttagcactt 360
taagtttatc acattttttt gactctctgac attccacttt cctagggttat aggaagagtc 420
tgtttatgta gtttgtttttt aaaatgtgcc aatgcc 456

<210> 65
<211> 654
<212> DNA
<213> Homo sapiens

<400> 65
aataaattcc agccttctct ttcttctctc ttctcagat attttctctc tttcttctcc 60
agtattcact ctcttctctg gattttgatg ggctgtttta tgtttttgca tgggtttctt 120
tctgtgtaat tttttctctc catatttctt atatgtctaaa ggtattccat atttagcggc 180
aggcctttgta attttctctg caggcataac agaaaatcag ttattgtctg aagctggctt 240
tttagctgtg ataggctgtg atccaaactt cgaaaaatgt tttagacaaa attctctctc 300
aataagctga gggagagaaa acttttcaat gcgtttggct ataaaaactt tctccaatat 360
ggagttgact gatgtctctat cctaggatt tcttttaaat aactgagaca ccaaaactgc 420
gagatcatag gaataatgca aagcacacag tggaaaagat ccagatatata tcttctagtac 480
caggtttttc atactgccag ctccaaaagc atgttttaagt gtacacagat cataaaggac 540
acccccaga gcccaaatgt ctttttatta ttgtaagttt gttttcacag atttcagggt 600

acaagtagat tggggccccc atcaagttcg gccccctctc cagtctttta gaac 654

<210> 66

<211> 592

<212> DNA

<213> Homo sapiens

<400> 66

```

tttttttttt tttttttatt gggaataaat ttatcaaaaa acatgtcatc caattccacc 60
aatgagacaa ttttaaatat agaataacact ctgttcatga atataaaato cccaggtgaa 120
agtcccttaa aacactatta tggttatggt tcttagaata attttataac tttttcagag 180
aattccttta aacttggttaa aataccttgt tgctagtgtc cagaacatct aggttcagtc 240
tttattttta agacagtatc tatcctaggc aaatgagagc ttgtttttat gtatttaaga 300
gtttcctott gtcatttcaa tgtcaaatgt atttgactca atttcatgat ttcattctgc 360
tcaaggccat caaccggtca gagccagagc ccttcaaaag ctgtatgtga gtatatgagg 420
gaaaactttc cacataattt tacatcattt ctatctcata gcagtttttag ttttctcata 480
gctatctcat acagagtttta gttttctcaa attctatgct gttttgttac tactgcagct 540
gaccaatcca aagccagttt acactcagca tgtgttattc tactttaaaa ta 592

```

<210> 67

<211> 469

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<222> 245,298,314,339,424,440,465

<223> n = A,T,C or G

<400> 67

```

gatgccaaaa atgctttccc aagtggctaa cattctgtat tccaccagc aatatatgag 60
agattaagtt gcttttccaa cocattttatg ctacgtattg tcaggttttg tttgtttctg 120
ggttctttat ttgttgggtt tcttttttat ttccagcatg ctaatagggtg tgatttgtgtg 180
tttaatttgc aattccctaa ctccataaat tagggaaacac agaacaacac tatgacacag 240
aaaantgcot ttgacctgat ttacttctct actattaaga aacagataaa attcatantg 300
tccttggaac acnntttttt tgttgcttta ttgtcatna catttaactc tttgttaagt 360
ggaaatggtc tcttcagata atttttttcc atttttaaac aggttggttg acctatacat 420
tgtngttttg agagttccan aaggtatccc gtattccaaa tccctncatt 469

```

<210> 68

<211> 510

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<222> 424,462

<223> n = A,T,C or G

<400> 68

```

tttttctcga gaatttaatt ttatttgctg tagattcaaa atgagggaagt ggtaaatgca 60
ttatttactc aaagcataaa gtcagccctta ggtaggagat gtaacaactc ctcaacttta 120
cactatccag ttaaagccaa tttttaaaac ctttttttct ctatgatga ccccttgagtc 180
atagaaaaat ttctatttta gaaaatgtta agcatgaaca caaaaagact acgataaacag 240
tgttataaac actcgtgtac ccaaggccca gctttaacat tcatacctta gcagtgttaa 300
ggtagtgctt aggttgaaat ttatattgtg tgatcagaaa taaagagcag ttcttgcaga 360
tagctagaat tacttcattt ttataggagt tttaggcata aactaacaag ggaactagg 420
ccnnttatag taaatatcct aaaagcattt taattttaca gnatttgaca gcggtatgcc 480

```


atggacattat tccatttgg tcaggggcaa

510

<210> 69

<211> 483

<212> DNA

<213> Homo sapiens

<400> 69

tgcatcagtt	aatgtaatac	gcccacagga	tggggattga	atggaagat	gccacgtacc	60
tttaagatat	gaagctggto	tgaagtacac	cttgaacaat	atatgtacag	ttcatcacac	120
actgtattta	tttctctctc	tgtaaatctc	cgagagaacag	aatttaagac	ttggggcaaaa	180
cagagtcctc	tttctctctc	aacttgaaaa	caagaaatag	attcccttc	caacacagtc	240
tgagtgaatt	ctgtggagct	atctgaagg	atgagcaatg	ggccaggaag	aacctgaggt	300
gatggaaag	gcagaaatac	agtagggcag	atgctttctt	gggaatgcg	agcagaaaat	360
gtgtgctgtc	caccagcgag	ctctgactac	tttaatggaa	ttgtgcatg	tgtgtttcaa	420
actgggatta	aatggcaatt	ttaggggaac	agtacaggtc	gctcatatg	ctccatcagt	480
ttc						483

<210> 70

<211> 481

<212> DNA

<213> Homo sapiens

<400> 70

gtactggaca	gacgtgagcg	aggaggccat	caagcagacc	tacctgaacc	agagcggggc	60
cgccgtgag	aacgtgtgca	tctccggcct	ggtctctccc	gacggcctcg	cctgcgactg	120
ggtgggcaag	aactgttact	ggacgggactc	agagaccaac	cgatctcgag	tgggcaacct	180
caatggcaac	tccgggaagg	tgtctttctg	gcaggacott	gaccagccga	gggccatcgc	240
cttgaccccc	gctcaagggt	acatgtactg	gacagactgg	ggtgagacgc	cccggaattga	300
gcggcgagg	atggatggca	gcaccgggaa	gatcattgtg	gactcggaac	tttactggcc	360
caatggactg	accatcgacc	tggaggagca	gaagctctac	tggtctgacg	ccaaagctcag	420
cttcacccac	cgtgcccaacc	tggacggctc	gttcggcagc	aagggtgttg	agggcagcct	480
g						481

<210> 71

<211> 341

<212> DNA

<213> Homo sapiens

<400> 71

cgcccgaggc	gaggctggag	aagtgtgct	ggccggggca	gtcgtccag	caggccgggg	60
acgcgggggc	ggcagggggc	gtggggcccg	gctctgggtg	gggtctctgg	gccgcacac	120
agctgcgaag	ggtgatgtcg	gcagagcccc	ctgactccag	tggtatgggg	tgtgtgtgga	180
agtggcgagg	catgtcaagc	acagactgga	accacagatg	ctgtacgtga	caactggcgt	240
ggcgtttcag	ggacaggcgc	atgtgcttgg	ccttgccctg	gaagttagag	gtcagcagct	300
actccccagg	cagagtctca	ctttggcgcg	ccctcgtgcc	g		341

<210> 72

<211> 283

<212> DNA

<213> Homo sapiens

<400> 72

tttttagatc	catccattta	ttccttcagc	caacattttc	tggtattcct	tgtgtgtctag	60
gctcgtgtcc	accatctgga	gatgcagaga	ggcgggagac	ccatgtggcc	tttgaggggc	120
tttcaggctc	gtgggggttc	aggcacagac	accaccaatc	tgaaccaggg	gactcgagga	180
tgtggggtta	ggggagagag	ggaatggctg	gtcggcctag	gggttctcta	ggaagtcttt	240
gggggtaagg	agagaactcc	tgaaggtaa	ggagagccgc	agg		283

<210> 73
 <211> 485
 <212> DNA
 <213> Homo sapiens

<400> 73
 ttttttttat ttttaggata ttttatttta atgcaaatga aatttctatc tatgtgaaac 60
 tggtaaaagg gagatatagg aactcctatt tttctctctg tcttctctctc tgtttcttctc 120
 tttttttatt atttttggat tatagatgct cctctcagtt gcaagttgca atgtccaca 180
 tctctcagcc agcacctggc tctgttcagc ggcttttagt gaggctctctc tgtcaaggca 240
 tgaataatac agcccttagg ctgttggcag actccaaatg aggcgtgcac acatcaggaa 300
 gcaagccctt gacttttagct ccagaacagc ctctctctgt gtcttgcata tttgccactg 360
 acatgaccac tgcogtcaca gccaggggtg ggacagctga acagctcttg tatggctggg 420
 tccacgggaa ctcgaaaccc ttgggacgcg gtgcgatgcc gcttctctc cgtgtgcac 480
 tccat 485

<210> 74
 <211> 338
 <212> DNA
 <213> Homo sapiens

<400> 74
 ttttttgatt atttcagaga tttattgcaa gtttaattgtc tgtgaagctg gatattcctt 60
 aacatgaagg taataaactt taacgttcca ctcaaaaaga caaaaaccaa acaacgaaaa 120
 ataagaaatt aaccagaaag ctatagcttg ttttcttact cagaaaaaaa gtataactga 180
 taaggtacaa tttctgtaac tggatatttt tcaaaattat aaggctttta gttctaanaag 240
 tataaagAAC tgtgatgcac tcttagtcaa cctaactctg ctagaagctt tatcaacact 300
 gcagcttcca atactttctc ttttgctatt atatagtc 338

<210> 75
 <211> 334
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <222> 265
 <223> n = A,T,C or G

<400> 75
 agcggcgcgc gcggagcagc aacagttcta cctgctcctg gaaacotgc tcagcccccga 60
 caatgtggtc cggaacacagc cagaggaaac ctatgagaat atcccaggcc agtcaaaagt 120
 cacattctctc ttacaagcca tcagaataac aacagctgct gaagaggcta gacaaatggc 180
 gcgcgttctc ctaagacgtc tctgtctctc tgcatttgat ggaagtctat ccagcaacttc 240
 cctcttgatg ttcagactgc catcnagagt gagctactca tgaattattc agatggaaac 300
 acaatctagc atgagaaaaa aagggtttgtg atat 334

<210> 76
 <211> 248
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <222> 32,33
 <223> n = A,T,C or G

```

<400> 76
gataggcata aacgtgttta ttaagtgaac cnnatccttt aaaaaataaa aaggggaagcc 60
tgtatataaaa tgaagttgtg gattcaacta gccagaattt attctgactt gcaccaaacc 120
acacaaaatc tttttaaagt ctagttagtc gtagtctaaa tggacactcc agagctctgtt 180
cttgaattcc attgcaagag ctccaacttc ctactttcag aaggggatggg gatcaagatg 240
agggttgt 248

<210> 77
<211> 515
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<222> 395,476
<223> n = A,T,C or G

<400> 77
atgtagaaac agcatcaagc tgtttctctc tacogtcttt gatagaaata aaaaaataaa 60
taaaaagtgt aatgtgcagaa aagctaagag gtttttagtt tttgtttttt gttttccttc 120
cacccagtca tatttggaaa ggatttagtg agtctgggtt atttttagctt caatctgggt 180
ttgtacacaa gcaaaaagca aatgttgta tttcaggtag accttcactgc agacattgcaa 240
aaccaactgt ctccgtgtgt aggagccatg gggagctctc cgaagggctt tccaggcagt 300
gggctaattg gcaaaatgac tactcagtg cctctgtgac cgtatggttaac ggtgtgccaa 360
ggatattcat cagccactct gagaatatga aacanagtgc tgagattcta cttaacctaa 420
taacaaagaa acgtaagca acacgaactg cagccagaag ggaacactgg aatggngggg 480
tgaatgttgt cctgattagc acccccaat ctccg 515

<210> 78
<211> 532
<212> DNA
<213> Homo sapiens

<400> 78
ctgttgttta tatagtttat tactgtcata gctaagaaaa ggcagtcgat ttcaacataa 60
tcoatatcta tgttcaaatt ctcaaaactat aggatatcta tgtttcaaat tgtaattttat 120
aacctggtaa gtatttctaaa caaaatatgt acaatccatt agctgacctt aaatcttatg 180
aagctgtatc atcagtttaa caaatacaca cgactttagc aaaaagtata acagatagta 240
ttataaatca ttaataatca ggcattgact aaaaaataca gataaaattg gagcaaat 300
aaagaggagt tgcattcaaa atattttttc catttgatat cattagaatt acaaaacagc 360
taataaaaaa atctaatgtt aaggcaatga caaataacaa agataaacagt tgcccaagga 420
ggcaggggtt gggagggtgaa tgcacaatca aggagggggc caaaacagcc ttcagggttaa 480
tttgttttat taagggggga gtcattggta gatagtcctt acatcttttt at 532

<210> 79
<211> 431
<212> DNA
<213> Homo sapiens

<400> 79
gggataagca aatgagtcac aacctttatt ctgataatag ccagtaaaatt tgcaaaagaga 60
ggagacaaac tgtaattgta tacataaaaa cacctagtc cactttaaaa ttttaataac 120
tatattatagt actgtattta atttttaaag atgaagacag caaaaattatt cacattaaaa 180
tatctttacag aaatcattat tcttctattc aagaaaacca attatactaa gttacacagg 240
aaaatttaac agagggaaatt ctccctggga cacttattga actgaggatt tcaactcata 300
gtttaaaaaa gtaaacaggt ctccaggtgtc tttttcatgg gtatggcacc ttatcaatct 360
gaattacagt tcatgggtgaa agctaacttt ttttgtgtga aataagttaa taatgccaat 420
tcagttttctt g 431

```

<210> 80
 <211> 431
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <222> 361,431
 <223> n = A,T,C or G

<400> 80
 acaaaccttc cggggggttc ctgagtggtc gctctcggaa aagcggatcc taaataaagc 60
 gggagggtta tagggcgacg tcgaggagag gacagggtcc gactcactgc tacagtttca 120
 ggtoactggg ctccgcagca gatcgtgttt ttccccgtgg ctcgagagct gcgctgggtt 180
 ctcatgcaaa ctccagagccg agctaagtac atgagcaact ttacttttta cacaagatga 240
 gcacgcgtgc cgaggcgctg ggcggcggtc gtgtgagttg gtggcccaga cgaacagctt 300
 gtgcgagact ctgggcattt cggtttctag atacaagatt tgcctaaatg tcacagtcca 360
 nagaagtga ttctagcat tgtagctact ggatgcacac aaagtaaaaa aaaaaaacct 420
 tcaactgccg n 431

<210> 81
 <211> 471
 <212> DNA
 <213> Homo sapiens

<400> 81
 aaggtcagat attgtttaac acttgaaatt ccaagagaaa aaaaatttcc caatgagtgc 60
 tctgttttcc atagagtaat tgctgaaata aaggaaacaca gaaaacaagg cttctgccag 120
 ttgtcactta caaaaacata cagaggatca taactagag acatggctaa ggctcaggt 180
 ggtttcattgc tcaagattga tgttttgcga gagagctgag ttgtggagtc ctgtttcggg 240
 agggcgctga tgggtgtgac ttcatctcca gctccttgct ttagggtctg ggcaagcttt 300
 tgaggctctg aacttgttga agacttgtgg acagagaatg gctgatatct cttaattttg 360
 tacagttgag gaacctgcag attgaagaag gaataactct gcttgatttg aactttctga 420
 gacttaattg ggaccagtc aaggccatca ggagccaact cgttggagtc c 471

<210> 82
 <211> 450
 <212> DNA
 <213> Homo sapiens

<400> 82
 tgcctaattt tgcaaatcaa agtgtatcat ttctccaatt ctactgatgc cagtttccaa 60
 gtccaattac tttttctacc ttctaatttt tottaatttc taagccaata tgttaaaaaa 120
 tattcttttg gotttccaaa tgttgcaatta tcttaactgc ctctgataat tccaacaatt 180
 catttggtct taaatgaac ttcttccatg taatgctctt tattaaatgt agatgtttcc 240
 ttaagaatga atctgcacca gacctttgct ctctccatg atttcaccta ctctcacaat 300
 ggtgatgggc attcccatgg cctgcacagc ttactgtatc tctttagcct gatctctccc 360
 tagaaatata atgttcatct gtgtttgtct gatgaggact gcctgatagc tgccaaatca 420
 acaaggataa aaccagaatt cacattccct 450

<210> 83
 <211> 540
 <212> DNA
 <213> Homo sapiens

<400> 83
 ttatacaaaa gcatttaaca agcttaaaaa atgaaactca atgaaaaaaa aaagaaggtt 60

tgaacacagt	caaataacct	gagaagtgc	agatggaaaa	gcaacagaat	gcaagcacct	120
tgtaaggtct	gtaattcttg	gatttactgt	gaaaagtttc	agaacatcat	agactcttac	180
tgccacattg	tccatagacc	ctggaaaata	acagtgaat	tcatatgtat	acacatatat	240
atgaatacac	actcatgcat	gcacactgtc	ttcacacacc	cctcctcacc	acttaaccgg	300
agttacataa	atgctttctc	gatatgtcat	tgcatattgt	tgttttctgc	atctcaacta	360
agttcagcgg	cttgccgctg	tgacattaat	tatgcaagat	tcaaacaccc	aagcaggcac	420
attttggggg	tgagttttaa	gaaatctgtg	acctgaaaga	aattctgtgg	ggactgtctg	480
ggttatccag	tttatccctg	gatttatatt	tgtttttagg	tcttgacctc	tttttaagct	540

<210> 84

<211> 559

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<222> 493,499,506,517,537,550,559

<223> n = A,T,C or G

<400> 84

gtgtgtgctg	ctgtttttac	tcggacaagt	cttattttac	agcggaattg	acaaataaag	60
ccttattttta	cacatccgaa	gaaacaccat	cacaggagggt	ttgtaggctg	gotgtgtgct	120
ttccaaaaca	gcaaaataga	ttcttcccat	ccaacccctc	ttctcttgtg	agagttaggt	180
gtggctctgt	gggctctgtc	tctctgcagg	cacagaaact	ggcagacctg	gtccctcctg	240
agcggggcct	gctcaaggga	atgggtgccag	attttgaaca	caggtaaaaca	ggctccttca	300
taacaacact	gtgcattttc	gtgtcatttt	gtttattgtc	cactgagtgt	ttggccacctc	360
agctcttgtt	ggaaaaacgt	gggtgtccag	aaattgctga	cacaagaaga	tggaattgct	420
atggtccggt	agggacacag	ggcagcccca	gccagatccc	actggtccat	gcagggcac	480
gcagtagaaa	ctnaacgttnc	cactngtaa	cagggtncaa	gacaccaatt	coggcancat	540
gggaaagaan	taaaccttn					559

<210> 85

<211> 2466

<212> DNA

<213> Homo sapiens

<400> 85

agttggtccg	agctgcggaa	aggtctggtc	gcagagacag	gaacgtgtaa	tcctcagcgt	60
gctccagccc	acagcttcgc	tcactgtctc	ggcagggcag	ctggcctctg	ggcaccggcg	120
gccctctctg	ctcgcgggaa	agcctgatga	agtcctccga	tattgatcag	gatttattca	180
cagacagtta	tcacaaagtg	tgcaagtgc	agctgatctc	cgaatcgcat	cgtgtggccc	240
actacgagag	tcgaaaacat	gcaagcaaa	tcgactgtta	ttacatgctt	caccocaggg	300
atggagggtg	tcctgcgaag	aggtcccggt	cagaaaatgg	aagtgatgcc	gacatggtgg	360
ataagaacaa	gtgctgcaca	ctctgaaaca	tgtoattcac	ttcagcgggt	tgggccgagt	420
cccattatca	aggcaaaatc	cacgccaaag	ggttaaaact	cttgctagga	gagaagaccc	480
cattaaagac	cacagcaaca	ccctcgagcc	cacttaagcc	cccacggatg	gacactgctc	540
cgtgtgtctg	atctccctat	caagaagag	atccagacag	atactgtggg	ctctgtgca	600
cctgcttttaa	taacccctctg	atggcccagc	aacattatga	tgccaagaaa	cacaaaaaga	660
atgcggcaag	aggttgccttg	ttagaacaac	tggggacaac	cctggatgat	gggggaactga	720
gaggtctgag	gcgcaattac	agatgtacca	tctgcagttg	ctccctaaac	tcaatagaac	780
agtatcatgc	ccatctgaaa	ggatctaacc	accagaccac	cctgaagaat	aagttagtga	840
agcatcaatc	aagacataag	aacaaaaacat	tagcattttc	ctgcogtgga	gaattgctta	900
tcacaaacca	gaggaggctt	ctttcttgaa	caataaacat	ttcttataag	gattccacga	960
ttccacatag	actgatcttg	atttttgaaa	atgaatgagg	ttctcttttt	tttttctctt	1020
tttttaattt	tggggtgaagt	tatgatatatt	ggatggattt	ttaaattctt	ccttgataac	1080
ataattagca	catgttctaa	attataatcc	tatagcaaac	agttggagca	ttattcaaac	1140
tgaaaatgga	aaaatttttaa	tttccaattt	attctagatt	tcctcagagc	ataattatcc	1200
tgttaaatcc	tcaatgagtg	tgatgtaaac	cacctctatc	cagaaaata	cattcttttc	1260

```

tcatcatggt ggacacagtt gagggtgaca tgcacagaa tggaacagat cactattagt 1320
ggaaaataacc aaatggacaa ataaatacca gtcgttttct ccgtttotcca agcacaggag 1380
ccagggtttac catctgaaca atgaagacga agggagtaaa taaaggaaga atttcatctc 1440
tttttctgga tcattcaaa aacagtttct caagggttaag ccaagtcctc cttgcaagtt 1500
gccaaataat agcttaggaa aagaatttagt ctgcctgcac gatgatcttc ttagggcaaaa 1560
acgtcttcac agcccttgac cttgggtgaat ttttttcccc aaaagcatcc aaaagaaaga 1620
ttataaacccc cagaacgaga tggaaataaa caagtatttt ttttttatga tgtttggcct 1680
gaaactgtgg ctttaattgg gggatactga tcgtttggaa agaagtgaaga aaattctgaa 1740
gaaatggcgg ctttgggcta ggcggggctcc cctatttctt ctgtttctca ctgaagtctc 1800
actgctgagc caagactcag tcaactctga aagagcatga ccgataaaga aaacagttcc 1860
tttctgatgg ggagcgtctg agtgcagatc atgaggctct tctctagagt ttaattcttt 1920
tccatggtga ccggacttgg tgtctgttag cctggttacg aagtgggacg ttgagcttct 1980
actgacgatg ccctgcattg accagctggg atctggctgg ggctgccctg tgtccctaac 2040
gaccataggg aatccatctt cttgtgtcag caatttctgg acaccactg ttttccacca 2100
agagcttgag tggcaacaac tcagtgcaga ataaacaaaa tgacacagaa atgcacagtg 2160
ttgttatgaa ggagcctggt tacctgtggt caaaactctg caccattccc ttgagcaggg 2220
ccgcctcagg agggaccagg tctgcccagt tctgtgctg cagagagacg aagcccccag 2280
atgccacccc tactctacaa gaggaaaggg ggttggatgg gaagaatcta ttttgcgtgt 2340
tgggaagaca cacagcgcac ctacaaacct cctgtgatgg tgtttcttcg gatgtgtaa 2400
ataagccttt atttgtcaat tccgcgtgaa aataagcatt gtccgagtaa aaacagcagc 2460
aacaac

```

```

<210> 86
<211> 408
<212> DNA
<213> Homo sapiens

```

```

<400> 86
ttttttggca tttaagtttt tcaccaattt attgctaaga gaaacatat aataatagc 60
tatagggtca taaaaccac ttgcaagcta tagaagcaag ttctgcctgt gctgtgtat 120
gtgtatgtat gacagtggac atgtaagtgt gaaactttaa acactattac agtaagaagt 180
ctttttgtga acttttgtta gtttgagagg ctgcaatgat ttttctcctt tcaaaatgct 240
gaaatagaac tcatcatctt gcttttcaaa ttgcaaacg gtagctgggt tggaaggctg 300
gagattgatt tctctccagc tagcaagctg tggggctcag tcaactgaagc atgtgggtga 360
tatgtgcaac caccacattg gcaaatattg aactatttta agtgcac 408

```

```

<210> 87
<211> 431
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> misc_feature
<222> 361,431
<223> n = A,T,C or G

```

```

<400> 87
acaaaccttc cgggggttgc ctgagtggtc gctctcgaaa aagcggatcc taaataaagc 60
gggagggtta tagggcgagc tcgaggagag gacaggtctc gagtcactgc tacagtttca 120
ggtcactggg ctccgcagca gatcgtgttt tctcccggtg ctcgagagct gctgtggttt 180
ctcatgcaaa ctacagagcc agctaattgac atgagcaact tttactttta cacaagatga 240
gcacgcgtgc cgaggcgctg ggcggcggtc gtgtgagttg gtggccagca cgaacagctt 300
gtcgagactc tgggcgctt cggtttctag atacaagatt tgcctaaagt tcacagttca 360
nagaagtgga tttagtcat tgtagctact ggatgcacac aaagtaaaaa aaaaaaaact 420
tcaactgcgc n

```

```

<210> 88
<211> 385

```

<212> DNA

<213> Homo sapiens

<400> 88

gaatattcag	tccacaaatt	ggcagacaat	gagatttaag	ccccctcttc	caaactcaga	60
cattggatgg	agagtagaat	ttcgacccat	ggagggtgcaa	ttaacagact	ttgagaactc	120
tgccatgtg	gtgtttgtgg	tactgtctcac	cagagtgtac	ctttctctaca	aattggattt	180
tctcattcca	ctgtcaaaagg	ttgatgagaa	catgaaggta	gcacagaaaa	gagatgctgt	240
cttgaggaga	atgttttatt	tcaggaaaga	tatttgcaaa	ggtggcaatg	cagtgggtga	300
tggttgctggc	aaggcccaga	acagcacgga	gctgctgtgca	gaggagtaca	cctcatgag	360
catagacacc	atcatcaatg	ggaac				385

<210> 89

<211> 272

<212> DNA

<213> Homo sapiens

<400> 89

tctttaaatt	acatacgaat	gtaaagagaa	aatggccaaa	acctcaaaac	tacgattgtt	60
gaaacaata	ttaaaggac	acaatctaaa	atcatgtctac	aaaaatagt	ttatctgttt	120
taactaaatg	tacatctttt	tttccaattc	catgattgac	aagagtgtct	atgcgacgca	180
tggaaggcac	cagaggtgaa	gtgattattt	gccttaaaat	atacaagaa	ttgcctactt	240
tgaaaaaagaa	atagtcatat	ttgtaaatga	at			272

<210> 90

<211> 504

<212> DNA

<213> Homo sapiens

<400> 90

gaagcagttt	attaccttaa	agcattttagc	aaacctaatg	tctgacctaa	tttcaaccaa	60
atgtctttat	tttccaata	atcttcaaaa	ctcttgattt	cccaaagcct	actaaagtca	120
tgctgtcaca	ggccattaga	cagcatgagc	agggcaggaa	agggctcttc	tcccaccac	180
caggaatgtt	gggtgatggc	tcagcagtta	tcacattgcc	tctctaaaag	tcatacatgt	240
gcacctaggg	tcaggaggac	gccatttctc	gatgggtccac	acctattgca	ctaaagtgtt	300
aattgaatgc	agatgccagg	gagatgcaac	ttccaggcca	aatgcattaa	gagacaaaac	360
ggcagagtat	gacctttccg	tggtcactcca	tgggaaaagg	gaagaaagcg	ttgggtgggc	420
atgtgtacaa	cttctctaac	acactgcagt	tgctcacctc	ccaaggatag	ggaggggcact	480
gtgcattgcg	gcagctcacc	ctaa				504

<210> 91

<211> 467

<212> DNA

<213> Homo sapiens

<400> 91

tttttttttt	tttttttttc	tttctcaaca	aatagtttac	tgggtggaac	ctaacagaa	60
taataattct	ttctgtccgt	aaataaaaaat	agatcatgct	tgaattgtct	actttgcccg	120
aactccccaa	gtcttcccg	atcttcagtt	cctccccctc	caacctgggt	tttatcagga	180
gaggggaaag	agcatttctt	gcctggcagg	aactcaagac	ctagaagaaa	gagggcctac	240
cctgcccaagg	aaacgacctt	cccttctctc	gcctctgtct	ctcttccctg	ttctgtctct	300
ttcttctctt	ttctctgggg	tttctctctc	ccgttaacta	tggggacaga	cacagctatt	360
cacaagtcgg	tctgggcagc	acactccgag	gtaaggcacg	aaggtcagga	gacaggttcc	420
cggtccccaa	atctctggaga	agatgagtta	aagctcttgc	cttccgat		467

<210> 92

<211> 229

<212> PRT

<213> Homo sapiens

<400> 92

```

Met Lys Ser Ser Asp Ile Asp Gln Asp Leu Phe Thr Asp Ser Tyr Cys
      5              10              15

Lys Val Cys Ser Ala Gln Leu Ile Ser Glu Ser Gln Arg Val Ala His
      20              25              30

Tyr Glu Ser Arg Lys His Ala Ser Lys Val Arg Leu Tyr Tyr Met Leu
      35              40              45

His Pro Arg Asp Gly Gly Cys Pro Ala Lys Arg Leu Arg Ser Glu Asn
      50              55              60

Gly Ser Asp Ala Asp Met Val Asp Lys Asn Lys Cys Cys Thr Leu Cys
      65              70              75              80

Asn Met Ser Phe Thr Ser Ala Val Val Ala Asp Ser His Tyr Gln Gly
      85              90              95

Lys Ile His Ala Lys Arg Leu Lys Leu Leu Leu Gly Glu Lys Thr Pro
      100             105             110

Leu Lys Thr Thr Ala Thr Pro Leu Ser Pro Leu Lys Pro Pro Arg Met
      115             120             125

Asp Thr Ala Pro Val Val Ala Ser Pro Tyr Gln Arg Arg Asp Ser Asp
      130             135             140

Arg Tyr Cys Gly Leu Cys Ala Ala Trp Phe Asn Asn Pro Leu Met Ala
      145             150             155             160

Gln Gln His Tyr Asp Gly Lys Lys His Lys Lys Asn Ala Ala Arg Val
      165             170             175

Ala Leu Leu Glu Gln Leu Gly Thr Thr Leu Asp Met Gly Glu Leu Arg
      180             185             190

Gly Leu Arg Arg Asn Tyr Arg Cys Thr Ile Cys Ser Val Ser Leu Asn
      195             200             205

Ser Ile Glu Gln Tyr His Ala His Leu Lys Gly Ser Lys His Gln Thr
      210             215             220

Asn Leu Lys Asn Lys
225

```

<210> 93

<211> 2327

<212> DNA

<213> Homo sapiens

<400> 93

```

gggagcgaaa accaacgtgt tcgggtgacag accccagcgc cgactgagcc tctaaagcga 60

```



```

cttcagctct gccccaccaa caccaccgcg cggccgggaa cagccgctcc ggggaagaaa 120
ctgaggggac tggcgggggc acgaggggaca gctgagggaa gggaggagcgc gagagaaaaa 180
gcgcaagcac gctgagggcc ggggggttgc aggagagggg cccgcggacc cgcagagcgg 240
aggaaagttcc gggagaaaag gggcggggac gaggagaatc cgggatcgcc tggcagaaaa 300
agagaaggga gtttctgaat cctgggaaga gggagcgctg gtagggatgc ttggcccgag 360
atccgacagc agggaaacgg agcgctccgg gggaggggct taatgctggg gaagggatgt 420
cttaaaagag gagaagcttt aaattagacg atcggagaag gctgagggaa tgcgtatgaa 480
ggggcgggag ctgaagtgtg gaggactcct ttgacagca gaaagggaaa cggctgtgaa 540
agtttccctc aaactccacc tgcctcctct ccaattcaaa ctccaactcc ttotocaaaa 600
gttaaaaaag aagccaagtt tgcacgcctc cctgttctct acttctctct 660
actccgccaac cgggaaaaaa gaaaaaaaaa actaatttcc ttcccaatat taggaactag 720
aaaaagctcta ggtccgcgaa cttgaatttt agcctagggg aatcaaaaaa gtaggagcat 780
taoctttgtt toctttttca aaatccaca cctcatcctt cctgcgacgc catgtotacc 840
aacatttgtg gtttcaagga caggtgogtg tocatcctgt gttgcaaat ctgtaaaaaa 900
gtgctcagct ctagggggat gaaggctgtt ttgctggctg atactgaaat agacottttc 960
tctacagaca tcoctcctac caacgcagtg gacttcaact gaagatgcta ttccacaaa 1020
atctgcaaat gtaaacgtga ggacatcgca tgtttaaaat gtgggaacat tttaggttat 1080
catgtgattt ttocatgtag ttctgtctct ctctcctgca acaacggaca ctctcggatg 1140
tttccagccc aggcagttta tgatattaac agactagact ccaacgggtg oacgctcta 1200
ctttggggca acttgccaga gatagaagag agtacagatg aagatgtgtt aaatatctca 1260
gcagaggagt gtattagata aatggaatta tgatatatat gatatacaaa cttttttcta 1320
tttaaaaaat tattaatgga tcaactttaa aattgttagt tgcagtgat cttttttgga 1380
aaacaaaaat ggggcatttg ttgatttatt tattttctgt ctctaattag ttaoctagat 1440
ttgattgaag ccagtgagtg ttgtcttttc ctctaactct ctactctctc ccccaacttt 1500
tttgcgccag ttagtggtga ttcttaaaat cagacgggaa gattotttca catatactc 1560
agtttaoctc caactggggg gagtttttct tacaacttga taccagatag cactaaat 1620
acattcttga ataaagcct agtaaccaag catatttcaa ccatgcatat atcaagttoa 1680
actgagtttt aataggggat taaaaaaaac agctgttagg ttccatggg cactggttct 1740
cataggttct attggtgata actgotttaa catggagcaa gaggtttga atcaggaat 1800
agaataaatt aaaaatttaa atatatagag gaatcctctt gatgtctcag catgatgtta 1860
gataaatgag ttgttcagaa aatatcagta tacgtctgtt acaatgttta tttatttaca 1920
tttctctaaa gccattatg atattgtatt atgagagcta aacctaaata agttatctg 1980
ttcctagga ccttctctgt aaatagttaa ttttagacga gtagtctgtc cttaatctta 2040
aatagaaaaa aaaaactaag ogatttgctt aagccatttg acattataaa gagctgtttt 2100
gttttgcttt gctttgcttt gttttgtttt ttttaagct gattcagagc cactgttttt 2160
ataggaaggt agggtagtgt tggattctgg ttttatgtaa ctctaaaaa aatgtatctc 2220
tttaatatct cagttgttagg gattttgtca atacaaaagc agactgagtt gtggtttgtg 2280
aaataaagtt ttttctaaaa atgaaaaaaa aagaaaaaaa aaaaaaaa 2327

```

```

<210> 94
<211> 2370
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<222> 741,1195,1683,2360
<223> n = A,T,C or G

```

```

<400> 94
gggagcgaaa accaacgtgt tcggtgacag accccagcgc cgactgagcc tctaaagcga 60
cttcagctct gccccaccaa caccaccgcg cggccgggaa cagccgctcc ggggaagaaa 120
ctgaggggac tggcgggggc acgaggggaca gctgagggaa gggaggagcgc gagagaaaaa 180
gcgcaagcac gctgagggcc ggggggttgc aggagagggg cccgcggacc cgcagagcgg 240
aggaaagttcc gggagaaaag gggcggggac gaggagaatc cgggatcgcc tggcagaaaa 300
agagaaggga gtttctgaat cctgggaaga gggagcgctg gtagggatgc ttggcccgag 360
atccgacagc agggaaacgg agcgctccgg gggaggggct taatgctggg gaagggatgt 420
cttaaaagag gagaagcttt aaattagacg atcggagaag gctgagggaa ttgctatgaa 480

```

```

ggggcgggag ctgaagtgtat gaggactcct ttagacagca gaaagggaaa gcccttgaga 540
agtccctctc aaactccacc tgccctcctc ccaattcaaa ctccactccc ttctccaaaa 600
gttaaaagga aagccaaagt tgccacgctc cctgttctct actcaataaa taactctctc 660
actccgcacc cgggaaaaaa gaaaaaaaaa actaatttcc ttcccaatat taggacttag 720
aaaagctcta ggtcccgcaa yttgaatttt agcctagggg aatcaaaaaa ttagggagcat 780
taactctgtt tctcttttca aaatcccaaa cctcatcctt cctgogacgc catgtctacc 840
aacaatttgtat gtttcaaggga cagggtgcgtg tccatcctgt gttgcaaat ctgtatacaaa 900
gtgctcagct ctaggggaaat gaaggctgtt ttgctggctg ataactgaaat agaccttttc 960
tctacagaca tccctctcac caacgcagtg gacttcaact gaagtgtcta ttccacaaaa 1020
atctgcaaat taaaactgaa ggacatcgca tgtttaaat gtgggaacat ttaggttat 1080
catgtgattg ttccatgtag ttctgtctct ctttctcgca acaacggaca cttctggatg 1140
tttcacagcc aggcagttta tgatattaac agactagact ccacaggtgt aaactctcta 1200
ctttggggcca acttgccaga gatagaagag agtacagatg aagatgtgtt aaatatctca 1260
gcagaggagt gtattagata aatggaatta tgatatatat gatatacaaa cttttttcta 1320
tttaaaaaa tattaatgga tcaactttaa aattgttagt tgccagtgat cttttttgga 1380
aaacaaaaat ggggcatttg ttgatttatt tattttccgt ctctaattag ttacctcagt 1440
ttgattgaag ccagttggagt tgtgcttttc ctctacttct acttctcttc ccccaacttt 1500
ttctgcccaag ttaggtgtta ttcttaaat cagacgggaa gattctttca catatacact 1560
agttacctcc caactggggg gagtttttct tacaacttga taccagatac catataattt 1620
acattctctga ataaaggcct agtaccacag catatttcaa caatgcatat atcaagttca 1680
acygagtgtt aataggggat taaaaaaaac agctgttagg ttccatggg cactggttct 1740
cataggttct atgtgtgata actgctttaa catggagcaa gagttgttga atcggaagt 1800
agaataaatt aaaattttaa atataatagag gaactcctct gattgtcag catgatgtta 1860
gataaatgag tttgtcagaa aatatcagta tacgtctgtt accaatgtta tttatttaca 1920
ttctctcaaa gccattatgg atattgtatt atgagagcta aaactaaaa agttatctctg 1980
ttctctagga cctctctctg aaatagtga ttttagacga ttagtctgtc ctaaaactta 2040
aatgaaaaaa aaaactaaag cgatttgctt aagccattgt acattataaa gagctgttt 2100
gttttgcttt gttttgcttt gttttgcttt ttttaagct gcaattcaag ccacaaagga 2160
ataggaaagt agggtagtgt tggattctcg ttttatgtaa cctacccta ctttctatt 2220
cctttgtctc ctgtaaacttt ttttacctat caatatgagt tgcgtgtctt cagtgtgtat 2280
tttttaagtt gctgggcatt acacttacca attaaagaat ttggaaatt caaaaaaaa 2340
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa

```

<210> 95
<211> 450
<212> DNA
<213> Homo sapiens

```

<400> 95
atgtctacca acattttagt tttcaaggac aggtgcgtgt ccatcctgtg ttgcaaatc 60
tgtaaacaag tgctcagctc taggggaagt aaggctgttt tgctggctga tactgaaata 120
gaacctttct ctacagacat cctctctacc aacgcagtgg acttcaactg aagatgctat 180
ttcccaaaaa gtgtcaaatg taaactgaag gacatcgcat gttttaaagt tgggaacatt 240
gtaggttatc atgtgattgt tccatgtagt tctctgtctc ttctctgcaa caacggagac 300
ttctggatgt ttccagagcca ggcagtttat gatattaaca gactagactc cacaggtgta 360
aacgtctcac tttggggcaa ctgcccagag atagaagaga gtacagatga agatgtgtta 420
aatctctcag cagaggagtg tattagataa

```

<210> 96
<211> 149
<212> PRT
<213> Homo sapiens

<400> 96
Met Ser Thr Asn Ile Cys Ser Phe Lys Asp Arg Cys Val Ser Ile Leu
5 10 15

Cys Cys Lys Phe Cys Lys Gln Val Leu Ser Ser Arg Gly Met Lys Ala
 20 25 30
 Val Leu Leu Ala Asp Thr Glu Ile Asp Leu Phe Ser Thr Asp Ile Pro
 35 40 45
 Pro Thr Asn Ala Val Asp Phe Thr Gly Arg Cys Tyr Phe Thr Lys Ile
 50 55 60
 Cys Lys Cys Lys Leu Lys Asp Ile Ala Cys Leu Lys Cys Gly Asn Ile
 65 70 75 80
 Val Gly Tyr His Val Ile Val Pro Cys Ser Ser Cys Leu Leu Ser Cys
 85 90 95
 Asn Asn Gly His Phe Trp Met Phe His Ser Gln Ala Val Tyr Asp Ile
 100 105 110
 Asn Arg Leu Asp Ser Thr Gly Val Asn Val Leu Leu Trp Gly Asn Leu
 115 120 125
 Pro Glu Ile Glu Glu Ser Thr Asp Glu Asp Val Leu Asn Ile Ser Ala
 130 135 140
 Glu Glu Cys Ile Arg
 145